

2016

Third Report - Addendum

ITALIAN ARTHROPLASTY REGISTRY PROJECT



Italian Arthroplasty Registry Project

**APPROACHING
DATA QUALITY**

Third Report 2016
Addendum

English version of Tables and Figures

Stefania Bellino
and Marina Torre

Executive Summary

Introduction

The *fil rouge* leading the 2016 RIAP Report is the word *quality*. In the previous reports, we focused on conceiving and developing the Registry and on the data collection that started in few Italian Regions, and furtherly extended to 13 out of 21. This third report points out that focusing on the quality of the collected data, both clinical and related to medical devices identification, is crucial. Therefore, starting from this year, a multidisciplinary Editorial Board will edit the RIAP Annual Report, to assure that results are presented taking into account the different points of view.

The total number of joint arthroplasties is steadily growing worldwide and represents an emerging challenge for the national health systems. Ageing, increase of obesity and arthritis prevalence, indication for surgery in patients even younger are only some of the leading causes of this trend. Policy makers and surgeons need reliable results to support the setting out of national health strategies and the selection of the most effective and safer devices. These will be available only if they are based on high quality data. Registries collecting high quality data will assist the implementation of an effective national medical devices vigilance and surveillance system.

The full report is published in Italian. In this addendum Tables and Figures of Chapters 4 and 5 (Statistical analyses on joint arthroplasty from national Hospital Discharge Records (HDR) database 2001-2014, and Data collected by the Registry in 2015) are translated into English for a comprehensive international reading.

Organization of the Report

The Report 2016 is arranged just like the previous editions: five Chapters and Appendices (seven, this year) including technical documents and information about the Project activities.

Chapter 1 briefly describes the Project, its structure and the data collection flow; Chapter 2 summarizes the procedures for the identification and characterization of medical devices through the RIAP-MD Library, a key tool of the Registry architecture implemented in cooperation with Manufacturers. Chapter 3 is an updating of the activities carried out by the participating Regions. Chapter 4 shows the analyses performed on the national HDR database for hip, knee and shoulder arthroplasties (Tables 4.1-4.31 and Figures 4.1-4.13). Finally, Chapter 5 shows the data collected by the 206 hospitals from the Regions and Autonomous Provinces participating in the RIAP project in 2015, eight and two respectively (hip and knee arthroplasties, Tables 5.1-5.20).

HDR data analysis

METHODS

The HDR national database was browsed and all the records referring to procedures coded with the following ICD-9-CM codes were selected:

- Hip: 81.51, 81.52, 81.53, 00.70, 00.71, 00.72, 00.73, 00.85, 00.86, 00.87
- Knee: 81.54, 81.55, 00.80, 00.81, 00.82, 00.83, 00.84
- Shoulder: 81.80, 81.81
- Other joints: 81.56, 81.57, 81.59, 81.73, 81.84, 81.97

The following analyses were performed:

- Time trends in hip, knee and shoulder arthroplasties – both main and secondary procedures – carried out from 2001 to 2014, and incidence rate
- Primary and revision joint replacements (main procedures) by patient characteristics, primary diagnosis, type of discharge, hospital length of stay and hospitalization burden
- Distributions of the hospitals by volume of interventions for primary total hip (THR), knee (TKR) and shoulder (TSR) replacements, for hip and knee revisions, and for partial shoulder replacements. Hospital was defined as low-volume hospital if

- performing less than or equal to 50 procedures/year (primary total hip¹ or primary total knee replacements), less than or equal to 10 procedures/year (hip or knee revision surgeries), less than or equal to 4 procedures/year (primary total or partial shoulder replacements)
- Inter-regional mobility measured by means of attraction and escape indexes for elective primary total replacements (hip, knee, shoulder)

RESULTS

In 2014, 175,290 arthroplasties were performed, 56.8% hip, 38.4% knee, 3.8% shoulder, and 1.0% other joints. Like in other countries, the total number of joint replacements unceasingly increased within the time range 2001-2014. Specifically, during the last year the number of primary and revision procedures increased by 2.2% for hip, 4.0% for knee, 12.6% for shoulder and 7.0% for other joints (Table 4.1 and Figures 4.1-4.4).

The volume of activity of every Region and the distribution of hospitals by class of volume are

¹ Glassou EN et al., Association between hospital procedure volume and risk of revision after total hip arthroplasty: a population-based study within the Nordic Arthroplasty Register Association database. *Osteoarthritis and Cartilage* 2016; 24:419-426.

shown for primary and revision hip replacements (Tables 4.2-4.4, Figures 4.5-4.6), primary and revision knee replacements (Tables 4.12-4.14, Figures 4.8-4.9), and shoulder replacements (Tables 4.22-4.24, Figures 4.11-4.12). The proportion of low-volume hospitals was 45% for THR and 54% for TKR.

The distribution of the procedures and the characteristics of the hospitals and patients are shown in Tables 4.5-4.11 (hip), 4.15-4.21 (knee), and 4.25-4.31 (shoulder).

Noticeably, the inter-regional mobility for elective primary interventions (hip, knee, and shoulder) is still high, although northern regions have an attraction index remarkably higher than the escape index (Figure 4.7, 4.10 and 4.13).

Registry data analysis

METHODS

Definitions

- *Regional participation rate:* Hospitals that submitted to the Registry at least one procedure/Hospitals in the Region that performed at least one procedure
- *Regional completeness rate:* Number of procedures submitted to the Registry/Number of procedures performed in the Region

- *Hospital completeness rate:* Number of procedures submitted by each hospital to the Registry/Number of procedures performed in each hospital

Data quality check

The quality of the collected data was checked out both syntactically and semantically. The syntactical check rejected the records in which information such as sex, age, type of hospital, joint, side, procedure, diagnosis, previous procedure, approach, fixation was missing. Instead, the semantical check rejected the records reporting procedures that included either values not accepted by the protocol or both revision and primary diagnoses, or if a primary intervention was paired with a revision diagnosis and vice versa. A crosscheck between diagnosis and previous procedure was performed and when resulted incompatible was classified as such.

Relating to the implanted medical devices, only primary procedures data were analyzed and the linked records included only if the following criteria were satisfied:

- The reference codes of the implanted devices were available in RIAP-MD Library
- No double components (for example two heads) were recorded

- Each procedure should have at least four components for a total hip arthroplasty, three components for a total knee arthroplasty.

RESULTS

The 52.009 procedures (Hip: 30,500, Knee: 21,509) recorded in 2015 by the National Registry represented about 30% of the national volume. The average completeness rate in the participating Regions was 61.9% (range 4.0%-98.6%) for hip and 59.0% (range 1.7%-100.0%) for knee; 48,135 procedures were included in the analyses (92.6%: 93.2% for hip and 93.7% for knee) (Table 5.1). Hospital completeness rates varied among Regions ranging from 4.0% to 100.0% for hip replacements and from 1.5% to 100.0% for knee replacements (Tables 5.2-5.3). Table 5.4 compares the distributions of the procedures recorded by RIAP in 2015 and by HDR in 2014, by joint and type of surgery. Tables 5.5-5.16 present the results of the analyses referred to the type of procedure (Tables 5.5 for hip and 5.11 for knee), type of provider (Tables 5.6 for hip and 5.12 for knee), patients' characteristics, and surgical practice, as well as diagnosis for primary and revision procedures (Hip: Tables 5.7-5.10; Knee: Tables 5.11-5.16).

The analysis of the implanted medical devices considered 27,880 primary procedures out of 38,704 satisfying the semantical and syntactical requirements: 12,611 hip (63.3%) and 15,269 knee (81.2%) (Tables 5.17-5.20).

Future perspectives

In 2017, a new informatics infrastructure will support the surgeons participating in the Registry, in order to reduce the burden connected to data input and to enhance data quality.

Some Regions are about to start, January 2017, the data collection for shoulder replacements; the list of the variables to be collected has been approved by the Italian Society of Shoulder and Elbow Surgery. In addition, a new cooperation has just started with the Italian Society of Ankle and Foot to define the information needed to set up the Ankle Arthroplasty National Registry.

Thanks to the collaboration with the Manufacturers, RIAP-MD Library currently counts about 55.000 reference codes. The cooperation with the International Consortium of Orthopaedic Registries provided useful information, regarding the classification adopted within the Global Library, which allows the undertaking of future steps such as linking each device to its technical attributes. Towards a European per-

spective, arrangements were recently agreed with the National Joint Registry of England and Wales to assess a possible cooperation to share RIAP-MD Library information within the joint component database organized by the National Joint Registry and the EndoProthesenRegister Deutschland.

Currently, the RIAP Working Group is increasing the dissemination activity to raise awareness of health professionals about the usefulness of the Registry and to include the largest possible number of Regions in the network. As soon as the law establishing the National Registry of the implanted prostheses is approved, the participation will be mandatory. Hence, it is hoped that both the participation and completeness rates in every Region will increase

to levels close to 100%. In addition, given its characteristics, the RIAP model based on HDR integrated by a specific additional Minimum Data Set, could be applied to other implanted medical devices.

The association between low-volume activity and negative outcomes is known. The proportion of low-volume hospitals (45% for THR and 54% for TKR) is quite large and spread throughout the country. Hopefully, the full implementation of the registry in all the Regions, providing reliable data on outcomes, will also result in the decrease of this proportion. In the meantime, if the responsible authorities decided to make data of hospitals volume public, the patients would be offered useful information to choose the hospital where to be treated.

Index of Tables

CHAPTER 4 - Joint arthroplasties: Hospital Discharge Records analysis

- Table 4.1 Joint arthroplasty in Italy (main and secondary procedures)
- Table 4.2 Hip. Arthroplasties performed in the Italian regions (main and secondary procedures)
- Table 4.3 Hip. Primary total replacement: number of hospitals by region and class of volume (main and secondary procedures)
- Table 4.4 Hip. Revision: number of hospitals by region and class of volume (main and secondary procedures)
- Table 4.5 Hip. Distribution of arthroplasties (main procedure)
- Table 4.6 Hip. Demographic characteristics by type of procedure
- Table 4.7 Hip. Arthroplasties by main diagnosis (ICD-9-CM)
- Table 4.8 Hip. Primary and revision procedures by type of discharge
- Table 4.9 Hip. Length of stay (days) by procedure and region
- Table 4.10 Hip. Primary and revision procedures by hospitalization burden
- Table 4.11 Hip. Primary and revision procedures by hospitalization burden. Details of the *Other* value of Table 4.10
- Table 4.12 Knee. Arthroplasties performed in the Italian regions (main and secondary procedures)
- Table 4.13 Knee. Primary total replacement: number of hospitals by region and class of volume (main and secondary procedures)
- Table 4.14 Knee. Revision: number of hospitals by region and class of volume (main and secondary procedures)
- Table 4.15 Knee. Distribution of arthroplasties (main procedure)
- Table 4.16 Knee. Demographic characteristics by type of procedure
- Table 4.17 Knee. Arthroplasties by main diagnosis (ICD-9-CM)
- Table 4.18 Knee. Primary and revision procedures by type of discharge
- Table 4.19 Knee. Length of stay (days) by procedure and region
- Table 4.20 Knee. Primary and revision procedures by hospitalization burden
- Table 4.21 Knee. Primary and revision procedures by hospitalization burden. Details of the *Other* value of Table 4.20
- Table 4.22 Shoulder. Arthroplasties performed in the Italian regions (main and secondary procedures)
- Table 4.23 Shoulder. Primary total replacement: number of hospitals by region and class of volume (main and secondary procedures)

- Table 4.24 Shoulder. Primary partial replacement: number of hospitals by region and class of volume (main and secondary procedures)
- Table 4.25 Shoulder. Distribution of arthroplasties (main procedure)
- Table 4.26 Shoulder. Demographic characteristics by type of procedure
- Table 4.27 Shoulder. Arthroplasties by main diagnosis (ICD-9-CM)
- Table 4.28 Shoulder. Primary procedures by type of discharge
- Table 4.29 Shoulder. Length of stay (days) by procedure and region
- Table 4.30 Shoulder. Primary procedures by hospitalization burden
- Table 4.31 Shoulder. Primary procedures by hospitalization burden. Details of the *Other* value of Table 4.30

CHAPTER 5 - Joint arthroplasties: RIAP data analysis

- Table 5.1 Participation rate and completeness of the regions involved in the RIAP project (2015)
- Table 5.2 Hip. Completeness of the hospitals participating in the RIAP project (2015)
- Table 5.3 Knee. Completeness of the hospitals participating in the RIAP project (2015)
- Table 5.4 Procedures distribution. Comparison between 2014 HDR data and 2015 evaluable RIAP data
- Table 5.5 Hip. Type of procedure
- Table 5.6 Hip. Procedures by type of provider
- Table 5.7 Hip. Demographic characteristics by type of procedure
- Table 5.8 Hip. Surgical practice for primary procedures
- Table 5.9 Hip. Primary replacements by diagnosis and previous surgery
- Table 5.10 Hip. Revision procedures by diagnosis and previous surgery
- Table 5.11 Knee. Type of procedure
- Table 5.12 Knee. Procedures by type of provider
- Table 5.13 Knee. Demographic characteristics by type of procedure
- Table 5.14 Knee. Surgical practice for primary procedures
- Table 5.15 Knee. Primary replacements by diagnosis and previous surgery
- Table 5.16 Knee. Revision procedures by diagnosis and previous surgery
- Table 5.17 Hip. Primary total replacement: type of implanted medical device
- Table 5.18 Hip. Primary total replacement: bearing surface
- Table 5.19 Knee. Primary total replacement: type of implanted medical device
- Table 5.20 Knee. Primary total replacement: implant of patella

Index of Figures

CHAPTER 4 - Joint arthroplasties: Hospital Discharge Records analysis

- Figure 4.1 Temporal trend of elective primary joint replacements in Italy
- Figure 4.2 Hip. Incidence rates per 100,000 population by type of procedure
- Figure 4.3 Knee. Incidence rates per 100,000 population by type of procedure
- Figure 4.4 Shoulder. Incidence rates per 100,000 population by type of procedure
- Figure 4.5 Hip. Primary total replacement: percentage of hospitals by region and volume of activity
- Figure 4.6 Hip. Revision: percentage of hospitals by region and volume of activity
- Figure 4.7 Hip. Elective primary total replacement: inter-regional mobility index (%),
a) attraction index, b) escape index
- Figure 4.8 Knee. Primary total replacement: percentage of hospitals by region and volume of activity
- Figure 4.9 Knee. Revision: percentage of hospitals by region and volume of activity
- Figure 4.10 Knee. Primary total replacement: inter-regional mobility index (%),
a) attraction index, b) escape index
- Figure 4.11 Shoulder. Primary total replacement: percentage of hospitals by region and volume of activity
- Figure 4.12 Shoulder. Primary partial replacement: percentage of hospitals by region and volume of activity
- Figure 4.13 Shoulder. Elective primary total replacement: inter-regional mobility index (%),
a) attraction index, b) escape index

Table 4.1. Joint arthroplasties in Italy (main and secondary procedures)

| ICD-9-CM Codes | Procedures | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | % (*) | |
|----------------|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------|------------|
| | Hip | 72.575 | 76.653 | 78.859 | 82.844 | 84.961 | 87.260 | 88.249 | 89.074 | 89.979 | 92.112 | 92.943 | 95.081 | 97.399 | 99.497 | 2.5 | |
| 81.51 | Total hip replacement | 45,792 | 48,793 | 51,311 | 54,442 | 55,599 | 57,612 | 58,650 | 58,786 | 59,528 | 59,764 | 60,712 | 62,361 | 64,056 | 66,045 | 2.9 | |
| | Total hip replacement (elective) | 39,144 | 41,396 | 43,419 | 45,764 | 46,561 | 48,157 | 49,104 | 49,289 | 49,923 | 50,394 | 51,422 | 52,940 | 54,624 | 56,561 | 2.9 | |
| 81.52 | Partial hip replacement | 20,768 | 21,358 | 21,020 | 21,657 | 22,402 | 22,418 | 22,326 | 23,069 | 22,542 | 23,953 | 24,177 | 24,324 | 24,998 | 25,313 | 1.5 | |
| | Hip resurfacing(*) | - | - | - | - | - | - | - | - | - | 303 | 476 | 157 | 94 | 96 | 70 | |
| 81.53 | Revision of hip replacement(**) | 6,015 | 6,502 | 6,528 | 6,745 | 6,960 | 7,230 | 7,273 | 7,219 | 7,606 | 7,919 | 7,897 | 8,302 | 8,249 | 8,069 | 2.3 | |
| | Knee | 28.056 | 32.704 | 37.703 | 43.093 | 46.257 | 50.651 | 55.123 | 57.706 | 58.628 | 60.761 | 60.973 | 63.214 | 64.763 | 67.365 | 7.0 | |
| 81.54 | Total knee replacement | 26,787 | 31,039 | 35,799 | 40,904 | 43,785 | 47,986 | 52,116 | 54,395 | 54,778 | 56,808 | 56,977 | 58,979 | 60,261 | 62,886 | 6.8 | |
| 81.55 | Revision of knee replacement(***) | 1,269 | 1,665 | 1,904 | 2,189 | 2,472 | 2,665 | 3,007 | 3,311 | 3,850 | 3,953 | 3,996 | 4,235 | 4,502 | 4,479 | 10.2 | |
| | Shoulder | 1.539 | 1.673 | 1.851 | 2.259 | 2.506 | 2.879 | 3.239 | 3.409 | 3.757 | 4.298 | 4.655 | 5.145 | 5.853 | 6.588 | 11.8 | |
| 81.80 | Total shoulder replacement | 695 | 798 | 934 | 1,239 | 1,455 | 1,688 | 2,036 | 2,175 | 2,515 | 2,965 | 3,444 | 3,793 | 4,421 | 5,307 | 16.9 | |
| | Total shoulder replacement (elective) | 405 | 503 | 634 | 868 | 1,080 | 1,331 | 1,620 | 1,773 | 2,073 | 2,355 | 2,784 | 3,011 | 3,464 | 4,089 | 19.5 | |
| 81.81 | Partial shoulder replacement | 844 | 875 | 917 | 1,020 | 1,051 | 1,191 | 1,203 | 1,234 | 1,242 | 1,333 | 1,211 | 1,352 | 1,432 | 1,281 | 3.3 | |
| | Other joints | 871 | 988 | 1,062 | 1,394 | 1,940 | 2,041 | 1,915 | | 1,696 | 1,665 | 1,623 | 1,739 | 1,674 | 1,720 | 1,840 | 5.9 |
| 81.56 | Total ankle replacement | 96 | 116 | 150 | 175 | 178 | 256 | 267 | 282 | 254 | 251 | 296 | 313 | 330 | 389 | 11.4 | |
| 81.57 | Replacement of joint of foot and toe | 336 | 409 | 435 | 489 | 649 | 673 | 736 | 720 | 571 | 614 | 608 | 683 | 507 | 526 | 3.5 | |
| 81.59 | Revision of joint replacement of lower extremity, not elsewhere classified | 219 | 189 | 183 | 363 | 707 | 599 | 383 | 153 | 201 | 133 | 111 | 90 | 103 | 103 | -5.6 | |
| 81.73 | Total wrist replacement | 45 | 45 | 46 | 50 | 63 | 86 | 74 | 75 | 61 | 54 | 76 | 84 | 69 | 54 | 1.4 | |
| 81.84 | Total elbow replacement | 92 | 147 | 163 | 214 | 250 | 321 | 320 | 316 | 411 | 404 | 438 | 451 | 478 | 499 | 13.9 | |
| 81.97 | Revision of joint replacement of upper extremity | 83 | 82 | 85 | 103 | 93 | 106 | 135 | 150 | 167 | 167 | 210 | 253 | 233 | 269 | 9.5 | |
| | Total | 103.041 | 112.018 | 119.475 | 129.590 | 135.664 | 142.831 | 148.526 | 151.885 | 154.029 | 158.794 | 160.310 | 165.114 | 169.735 | 175.290 | 4.2 | |

^(*)Annual growth rate^(*)ICD-9-CM codes introduced on 1 January 2009: 00.85, 00.86, 00.87^(**)ICD-9-CM codes introduced on 1 January 2009: 00.70, 00.71, 00.72, 00.73^(***)ICD-9-CM codes introduced on 1 January 2009: 00.80, 00.81, 00.82, 00.83, 00.84

Table 4.2. Hip. Arthroplasties performed in the Italian regions (main and secondary procedures)

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision | | TOTAL | |
|-----------------------|------------------------------|-------------|-------------------------------|------------|---------------------|-------------|--------------|------------|---------------|--------------|
| | N | % | N | % | N | % | N | % | N | % |
| Italy | 56,631 | 56.9 | 9,484 | 9.5 | 25,313 | 25.4 | 8,069 | 8.1 | 99,497 | 100.0 |
| Piedmont | 5,178 | 9.1 | 843 | 8.9 | 1,979 | 7.8 | 706 | 8.8 | 8,706 | 8.8 |
| Valle d'Aosta | 150 | 0.3 | 10 | 0.1 | 85 | 0.3 | 12 | 0.2 | 257 | 0.3 |
| Lombardy | 13,188 | 23.3 | 1,118 | 11.8 | 5,199 | 20.5 | 1,844 | 22.9 | 21,349 | 21.5 |
| AP Bolzano | 1,065 | 1.9 | 86 | 0.9 | 179 | 0.7 | 124 | 1.5 | 1,454 | 1.5 |
| AP Trento | 591 | 1.0 | 52 | 0.6 | 329 | 1.3 | 63 | 0.8 | 1,035 | 1.0 |
| Veneto | 5,946 | 10.5 | 776 | 8.2 | 2,227 | 8.8 | 775 | 9.6 | 9,724 | 9.8 |
| Friuli Venezia Giulia | 1,770 | 3.1 | 178 | 1.9 | 843 | 3.3 | 240 | 3.0 | 3,031 | 3.1 |
| Liguria | 1,700 | 3.0 | 723 | 7.6 | 574 | 2.3 | 341 | 4.2 | 3,338 | 3.4 |
| Emilia-Romagna | 4,928 | 8.7 | 523 | 5.5 | 1,653 | 6.5 | 784 | 9.7 | 7,888 | 7.9 |
| Tuscany | 4,963 | 8.8 | 747 | 7.9 | 2,074 | 8.2 | 744 | 9.2 | 8,528 | 8.6 |
| Umbria | 836 | 1.5 | 127 | 1.3 | 566 | 2.2 | 104 | 1.3 | 1,633 | 1.6 |
| Marche | 1,218 | 2.1 | 341 | 3.6 | 659 | 2.6 | 196 | 2.4 | 2,414 | 2.4 |
| Lazio | 4,343 | 7.7 | 1,032 | 10.9 | 2,060 | 8.1 | 628 | 7.8 | 8,063 | 8.1 |
| Abruzzi | 1,272 | 2.2 | 286 | 3.0 | 648 | 2.6 | 133 | 1.7 | 2,339 | 2.4 |
| Molise | 179 | 0.3 | 18 | 0.2 | 156 | 0.6 | 24 | 0.3 | 377 | 0.4 |
| Campania | 2,779 | 4.9 | 874 | 9.2 | 1,490 | 5.9 | 394 | 4.9 | 5,537 | 5.6 |
| Apulia | 2,132 | 3.8 | 536 | 5.7 | 1,507 | 6.0 | 333 | 4.1 | 4,508 | 4.5 |
| Basilicata | 290 | 0.5 | 45 | 0.5 | 274 | 1.1 | 33 | 0.4 | 642 | 0.7 |
| Calabria | 687 | 1.2 | 275 | 2.9 | 628 | 2.5 | 83 | 1.0 | 1,673 | 1.7 |
| Sicily | 2,519 | 4.4 | 752 | 7.9 | 1,643 | 6.5 | 391 | 4.9 | 5,305 | 5.3 |
| Sardinia | 897 | 1.6 | 142 | 1.5 | 540 | 2.1 | 117 | 1.5 | 1,696 | 1.7 |

Table 4.3. Hip. Primary total replacement: number of hospitals by region and class of volume (main and secondary procedures)

| | Class of volume of total replacements | | | | | |
|-----------------------|---------------------------------------|------------|------------|-----------|-----------|------------|
| | 1-50 | 51-100 | 101-200 | 201-300 | >300 | TOTAL |
| Italy | 343 | 204 | 135 | 49 | 23 | 754 |
| Piedmont | 11 | 20 | 15 | 5 | 2 | 53 |
| Valle d'Aosta | 0 | 2 | 0 | 0 | 0 | 2 |
| Lombardy | 38 | 34 | 18 | 11 | 9 | 110 |
| AP Bolzano | 2 | 2 | 5 | 1 | 0 | 10 |
| AP Trento | 4 | 2 | 2 | 0 | 0 | 8 |
| Veneto | 11 | 15 | 13 | 8 | 3 | 50 |
| Friuli Venezia Giulia | 1 | 6 | 6 | 2 | 0 | 15 |
| Liguria | 6 | 4 | 4 | 1 | 2 | 17 |
| Emilia-Romagna | 25 | 16 | 12 | 3 | 2 | 58 |
| Tuscany | 17 | 12 | 12 | 6 | 4 | 51 |
| Umbria | 6 | 6 | 3 | 0 | 0 | 15 |
| Marche | 8 | 6 | 5 | 1 | 0 | 20 |
| Lazio | 48 | 17 | 13 | 3 | 1 | 82 |
| Abruzzi | 10 | 6 | 4 | 1 | 0 | 21 |
| Molise | 3 | 2 | 0 | 0 | 0 | 5 |
| Campania | 47 | 16 | 8 | 2 | 0 | 73 |
| Apulia | 20 | 12 | 7 | 2 | 0 | 41 |
| Basilicata | 6 | 1 | 1 | 0 | 0 | 8 |
| Calabria | 15 | 2 | 2 | 1 | 0 | 20 |
| Sicily | 52 | 17 | 3 | 2 | 0 | 74 |
| Sardinia | 13 | 6 | 2 | 0 | 0 | 21 |

Table 4.4. Hip. Revision: number of hospitals by region and class of volume (main and secondary procedures)

| | Class of volume of revisions | | | | |
|-----------------------|------------------------------|------------|-----------|-----------|------------|
| | 1-10 | 11-25 | 26-50 | >50 | TOTAL |
| Italy | 447 | 158 | 39 | 24 | 668 |
| Piedmont | 25 | 18 | 4 | 2 | 49 |
| Valle d'Aosta | 1 | 1 | 0 | 0 | 2 |
| Lombardy | 59 | 29 | 4 | 10 | 102 |
| AP Bolzano | 4 | 3 | 0 | 1 | 8 |
| AP Trento | 2 | 4 | 0 | 0 | 6 |
| Veneto | 20 | 19 | 7 | 1 | 47 |
| Friuli Venezia Giulia | 5 | 8 | 2 | 0 | 15 |
| Liguria | 7 | 7 | 0 | 2 | 16 |
| Emilia-Romagna | 33 | 12 | 5 | 1 | 51 |
| Tuscany | 24 | 12 | 5 | 4 | 45 |
| Umbria | 8 | 3 | 1 | 0 | 12 |
| Marche | 12 | 6 | 1 | 0 | 19 |
| Lazio | 58 | 8 | 4 | 2 | 72 |
| Abruzzi | 16 | 3 | 0 | 0 | 19 |
| Molise | 4 | 0 | 0 | 0 | 4 |
| Campania | 49 | 6 | 2 | 1 | 58 |
| Apulia | 31 | 6 | 3 | 0 | 40 |
| Basilicata | 6 | 0 | 0 | 0 | 6 |
| Calabria | 14 | 2 | 0 | 0 | 16 |
| Sicily | 53 | 8 | 1 | 0 | 62 |
| Sardinia | 16 | 3 | 0 | 0 | 19 |

Table 4.5. Hip. Distribution of arthroplasties (main procedure)

| ICD-9-CM codes | | N | % |
|-----------------------|--|---------------|----------|
| | Arthroplasty | 96,147 | |
| 81.51 | Total hip replacement | 64,666 | |
| | Total hip replacement (elective) | 55,461 | 85.8 |
| | Total hip replacement (emergency) | 9,205 | 14.2 |
| 81.52 | Partial hip replacement | 24,613 | |
| | Revision | 6,868 | |
| 81.53 | Revision of hip replacement, not otherwise specified | 3,013 | 43.9 |
| 00.70 | Revision of hip replacement, both acetabular and femoral components | 1,090 | 15.9 |
| 00.71 | Revision of hip replacement, acetabular component | 1,397 | 20.3 |
| 00.72 | Revision of hip replacement, femoral component | 759 | 11.1 |
| 00.73 | Revision of hip replacement, acetabular liner and/or femoral head only | 609 | 8.9 |

Table 4.6. Hip. Demographic characteristics by type of procedure

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision | | TOTAL | |
|----------------------|------------------------------|------|-------------------------------|------|---------------------|------|----------|------|--------|------|
| | N | % | N | % | N | % | N | % | N | % |
| Patients | 54,671 | | 9,107 | | 24,364 | | 5,850 | | 93,992 | |
| Procedures | 55,461 | | 9,205 | | 24,613 | | 6,868 | | 96,147 | |
| Gender | | | | | | | | | | |
| Males | 25,255 | 45.5 | 2,426 | 26.4 | 6,480 | 26.3 | 2,591 | 37.7 | 36,752 | 38.2 |
| Females | 30,206 | 54.5 | 6,779 | 73.6 | 18,133 | 73.7 | 4,277 | 62.3 | 59,395 | 61.8 |
| Age by gender | | | | | | | | | | |
| Males | | | | | | | | | | |
| Mean age | 65.4 | | 72.2 | | 82.4 | | 69.6 | | 69.2 | |
| Standard deviation | 11.9 | | 12.0 | | 9.6 | | 12.3 | | 13.2 | |
| Median age | 67 | | 74 | | 84 | | 72 | | 71 | |
| Interquartile range | 58-74 | | 65-81 | | 79-88 | | 63-78 | | 61-79 | |
| Females | | | | | | | | | | |
| Mean age | 69.5 | | 74.7 | | 83.9 | | 73.4 | | 74.8 | |
| Standard deviation | 10.8 | | 9.9 | | 7.3 | | 10.7 | | 11.6 | |
| Median age | 71 | | 75 | | 85 | | 75 | | 76 | |
| Interquartile range | 64-77 | | 69-82 | | 80-89 | | 68-81 | | 68-83 | |
| Class of age | | | | | | | | | | |
| Males | | | | | | | | | | |
| <45 | 1,426 | 5.7 | 51 | 2.1 | 53 | 0.8 | 114 | 4.4 | 1,644 | 4.5 |
| 45 - 54 | 3,212 | 12.7 | 163 | 6.7 | 82 | 1.3 | 208 | 8.0 | 3,665 | 10.0 |
| 55 - 64 | 5,815 | 23.0 | 364 | 15.0 | 127 | 2.0 | 408 | 15.8 | 6,714 | 18.3 |
| 65 - 74 | 8,598 | 34.0 | 716 | 29.5 | 602 | 9.3 | 802 | 31.0 | 10,718 | 29.2 |
| 75 - 84 | 5,745 | 22.8 | 781 | 32.2 | 2,594 | 40.0 | 896 | 34.6 | 10,016 | 27.3 |
| ≥ 85 | 459 | 1.8 | 351 | 14.5 | 3,022 | 46.6 | 163 | 6.3 | 3,995 | 10.9 |
| Females | | | | | | | | | | |
| <45 | 768 | 2.5 | 23 | 0.3 | 21 | 0.1 | 66 | 1.5 | 878 | 1.5 |
| 45 - 54 | 2,160 | 7.2 | 194 | 2.9 | 68 | 0.4 | 188 | 4.4 | 2,610 | 4.4 |
| 55 - 64 | 5,373 | 17.8 | 771 | 11.4 | 227 | 1.3 | 494 | 11.6 | 6,865 | 11.6 |
| 65 - 74 | 10,804 | 35.8 | 2,174 | 32.1 | 1,203 | 6.6 | 1,283 | 30.0 | 15,464 | 26.0 |
| 75 - 84 | 9,907 | 32.8 | 2,507 | 37.0 | 7,424 | 40.9 | 1,734 | 40.5 | 21,572 | 36.3 |
| ≥ 85 | 1,194 | 4.0 | 1,110 | 16.4 | 9,190 | 50.7 | 512 | 12.0 | 12,006 | 20.2 |

Table 4.7. Hip. Arthroplasties by main diagnosis (ICD-9-CM)

| | Males | | Females | | TOTAL | |
|---|---------------|------|---------------|------|---------------|------|
| | N | % | N | % | N | % |
| Primary total replacement | 27,681 | | 36,985 | | 64,666 | |
| Osteoarthritis | 23,871 | 86.2 | 28,442 | 76.9 | 52,313 | 80.9 |
| Fracture of neck of femur | 2,390 | 8.6 | 6,731 | 18.2 | 9,121 | 14.1 |
| Other disorders of bone and cartilage | 896 | 3.2 | 987 | 2.7 | 1,883 | 2.9 |
| Fracture of other and unspecified parts of femur | 90 | 0.3 | 168 | 0.5 | 258 | 0.4 |
| Other and unspecified disorders of joint | 76 | 0.3 | 103 | 0.3 | 179 | 0.3 |
| Complications peculiar to certain specified procedures | 54 | 0.2 | 132 | 0.4 | 186 | 0.3 |
| Late effects of musculoskeletal and connective tissue injuries | 53 | 0.2 | 104 | 0.3 | 157 | 0.2 |
| Other congenital anomalies of limbs | 46 | 0.2 | 83 | 0.2 | 129 | 0.2 |
| Other and unspecified arthropathies | 46 | 0.2 | 30 | 0.1 | 76 | 0.1 |
| Other orthopedic aftercare | 15 | 0.1 | 26 | 0.1 | 41 | 0.1 |
| Fracture of pelvis | 19 | 0.1 | 20 | 0.1 | 39 | 0.1 |
| Other derangement of joint | 16 | 0.1 | 15 | 0.0 | 31 | 0.0 |
| Other diagnoses (N. 75, freq <0.1% on total) | 109 | 0.4 | 144 | 0.4 | 253 | 0.4 |
| Primary partial replacement | 6,480 | | 18,133 | | 24,613 | |
| Fracture of neck of femur | 5,976 | 92.2 | 17,148 | 94.6 | 23,124 | 94.0 |
| Fracture of other and unspecified parts of femur | 122 | 1.9 | 365 | 2.0 | 487 | 2.0 |
| Other disorders of bone and cartilage | 97 | 1.5 | 220 | 1.2 | 317 | 1.3 |
| Osteoarthritis | 137 | 2.1 | 127 | 0.7 | 264 | 1.1 |
| Complications peculiar to certain specified procedures | 31 | 0.5 | 59 | 0.3 | 90 | 0.4 |
| Late effects of musculoskeletal and connective tissue injuries | 11 | 0.2 | 55 | 0.3 | 66 | 0.3 |
| Secondary malignant neoplasm of other specified sites | 15 | 0.2 | 19 | 0.1 | 34 | 0.1 |
| Fracture of humerus | 4 | 0.1 | 13 | 0.1 | 17 | 0.1 |
| Other diseases of lung | 8 | 0.1 | 7 | 0.0 | 15 | 0.1 |
| Multiple fractures involving both lower limbs, lower with upper limb, and lower limb(s) with rib(s) and sternum | 0 | 0.0 | 13 | 0.1 | 13 | 0.1 |
| Other diagnoses (N. 80, freq <0.1% on total) | 79 | 1.2 | 107 | 0.6 | 186 | 0.8 |

Follow

Table 4.7. Follow

| | Males | | Females | | TOTAL | |
|--|-------|------|---------|------|-------|------|
| | N | % | N | % | N | % |
| Revision | 2,591 | | 4,277 | | 6,868 | |
| Complications peculiar to certain specified procedures | 2,308 | 89.1 | 3,771 | 88.2 | 6,079 | 88.5 |
| Fracture of other and unspecified parts of femur | 43 | 1.7 | 92 | 2.2 | 135 | 2.0 |
| Fracture of neck of femur | 29 | 1.1 | 93 | 2.2 | 122 | 1.8 |
| Osteoarthritis | 40 | 1.5 | 72 | 1.7 | 112 | 1.6 |
| Organ or tissue replaced by other means | 41 | 1.6 | 70 | 1.6 | 111 | 1.6 |
| Other orthopedic aftercare | 22 | 0.9 | 29 | 0.7 | 51 | 0.7 |
| Developmental dislocation of joint, pelvic region and thigh | 14 | 0.5 | 24 | 0.6 | 38 | 0.6 |
| Other and unspecified disorders of joint | 16 | 0.6 | 17 | 0.4 | 33 | 0.5 |
| Other derangement of joint | 8 | 0.3 | 17 | 0.4 | 25 | 0.4 |
| Other disorders of bone and cartilage | 5 | 0.2 | 19 | 0.4 | 24 | 0.4 |
| Late effects of musculoskeletal and connective tissue injuries | 6 | 0.2 | 16 | 0.4 | 22 | 0.3 |
| Other complications of procedures not elsewhere classified | 12 | 0.5 | 7 | 0.2 | 19 | 0.3 |
| Disorders of muscle, ligament, and fascia | 5 | 0.2 | 10 | 0.2 | 15 | 0.2 |
| Fracture of pelvis | 4 | 0.2 | 6 | 0.1 | 10 | 0.2 |
| Bacterial infection in conditions classified elsewhere and of unspecified site | 3 | 0.1 | 4 | 0.1 | 7 | 0.1 |
| Malignant neoplasm of bone and articular cartilage | 4 | 0.2 | 3 | 0.1 | 7 | 0.1 |
| Arthropathy associated with infections | 3 | 0.1 | 4 | 0.1 | 7 | 0.1 |
| Osteomyelitis, periostitis, and other infections involving bone | 3 | 0.1 | 4 | 0.1 | 7 | 0.1 |
| Other and unspecified arthropathies | 3 | 0.1 | 3 | 0.1 | 6 | 0.1 |
| Secondary malignant neoplasm of other specified sites | 4 | 0.2 | 1 | 0.0 | 5 | 0.1 |
| Other acquired deformities of limbs | 2 | 0.1 | 2 | 0.1 | 4 | 0.1 |
| Symptoms involving cardiovascular system | 4 | 0.2 | 0 | 0.0 | 4 | 0.1 |
| Other diagnoses (N. 20, freq <0.1% on total) | 12 | 0.6 | 13 | 0.3 | 25 | 0.3 |

Table 4.8. Hip. Primary and revision procedures by type of discharge

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision | | TOTAL | |
|---|------------------------------|------|-------------------------------|------|---------------------|------|----------|------|--------|-------|
| | N | % | N | % | N | % | N | % | N | % |
| Type of discharge | 55,461 | 57.7 | 9,205 | 9.6 | 24,612 | 25.6 | 6,868 | 7.1 | 96,109 | 100.0 |
| Dead | 67 | 0.1 | 138 | 1.5 | 711 | 2.9 | 68 | 1.0 | 981 | 1.0 |
| Ordinary discharge | 29,218 | 52.7 | 5,118 | 55.6 | 13,204 | 53.7 | 3,744 | 54.5 | 51,266 | 53.3 |
| Discharge to a residential health care | 761 | 1.4 | 507 | 5.5 | 1,868 | 7.6 | 185 | 2.7 | 3,320 | 3.5 |
| Discharge to hospital at home | 84 | 0.2 | 30 | 0.3 | 71 | 0.3 | 16 | 0.2 | 201 | 0.2 |
| Discharge against medical advice | 70 | 0.1 | 45 | 0.5 | 107 | 0.4 | 19 | 0.3 | 241 | 0.3 |
| Transfer to an acute admission unit of a different hospital | 1,253 | 2.3 | 280 | 3.0 | 885 | 3.6 | 210 | 3.1 | 2,626 | 2.7 |
| Transfer in the same hospital | 13,093 | 23.6 | 925 | 10.1 | 1,853 | 7.5 | 1,299 | 18.9 | 17,167 | 17.9 |
| Transfer to an inpatient rehabilitation hospital | 10,720 | 19.3 | 2,021 | 22.0 | 5,417 | 22.0 | 1,269 | 18.5 | 19,418 | 20.2 |
| Discharge to a nursing home | 195 | 0.4 | 141 | 1.5 | 496 | 2.0 | 58 | 0.8 | 889 | 0.9 |

Table 4.9. Hip. Length of stay (days) by procedure and region

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision | |
|-----------------------|------------------------------|------------|-------------------------------|------------|---------------------|------------|-------------|-------------|
| | Mean | S.D. | Mean | S.D. | Mean | S.D. | Mean | S.D. |
| Arthroplasty | 55,461 | | 9,205 | | 24,613 | | 6,868 | |
| Italy | 8.3 | 4.7 | 12.3 | 6.2 | 13.0 | 7.4 | 13.4 | 11.2 |
| Piedmont | 8.3 | 4.2 | 13.2 | 6.5 | 14.3 | 8.2 | 14.3 | 11.0 |
| Valle d'Aosta | 7.4 | 5.4 | 13.4 | 2.9 | 13.5 | 8.9 | 16.3 | 11.7 |
| Lombardy | 7.4 | 4.0 | 11.7 | 5.9 | 12.8 | 7.9 | 11.6 | 10.7 |
| AP Bolzano | 8.5 | 4.0 | 11.4 | 6.8 | 11.4 | 7.2 | 13.4 | 9.9 |
| AP Trento | 6.7 | 2.8 | 10.3 | 5.9 | 11.3 | 5.6 | 10.2 | 6.4 |
| Veneto | 9.4 | 4.7 | 13.8 | 6.9 | 15.9 | 8.8 | 15.0 | 10.5 |
| Friuli Venezia Giulia | 8.8 | 3.7 | 13.6 | 6.8 | 15.4 | 7.9 | 14.6 | 12.1 |
| Liguria | 7.5 | 6.7 | 13.8 | 7.1 | 14.8 | 8.7 | 13.2 | 11.1 |
| Emilia-Romagna | 9.3 | 4.2 | 12.1 | 6.2 | 12.5 | 6.5 | 12.6 | 9.1 |
| Tuscany | 7.9 | 2.8 | 10.2 | 4.0 | 11.2 | 5.0 | 11.7 | 11.4 |
| Umbria | 8.3 | 3.4 | 11.7 | 7.6 | 11.2 | 6.5 | 12.0 | 5.6 |
| Marche | 10.0 | 5.0 | 14.9 | 6.1 | 15.0 | 8.7 | 15.9 | 12.9 |
| Lazio | 8.8 | 6.7 | 10.7 | 5.9 | 12.5 | 7.9 | 15.0 | 13.0 |
| Abruzzi | 7.9 | 4.5 | 11.7 | 5.1 | 12.0 | 5.6 | 12.8 | 9.8 |
| Molise | 9.8 | 8.2 | 15.5 | 6.4 | 15.7 | 6.1 | 17.2 | 13.1 |
| Campania | 9.0 | 4.7 | 13.0 | 6.2 | 12.5 | 5.7 | 14.3 | 11.0 |
| Apulia | 8.4 | 4.3 | 11.8 | 5.9 | 12.5 | 7.1 | 15.2 | 13.8 |
| Basilicata | 8.4 | 2.8 | 10.0 | 4.0 | 9.0 | 3.4 | 11.5 | 6.5 |
| Calabria | 8.6 | 4.5 | 11.5 | 5.5 | 11.0 | 4.7 | 15.7 | 11.6 |
| Sicily | 7.8 | 5.0 | 11.5 | 6.1 | 11.9 | 6.1 | 13.8 | 11.6 |
| Sardinia | 10.1 | 7.5 | 15.2 | 6.9 | 14.7 | 7.2 | 21.0 | 16.5 |

Table 4.10. Hip. Primary and revision procedures by hospitalization burden

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision | | TOTAL | |
|-------------------------------|------------------------------|------|-------------------------------|------|---------------------|------|--------------|------|---------------|------|
| | N | % | N | % | N | % | N | % | N | % |
| Hospitalization burden | 55,461 | | 9,205 | | 24,612 | | 6,868 | | 96,146 | |
| National Health System | 52,840 | 95.3 | 8,986 | 97.6 | 24,382 | 99.1 | 6,577 | 95.8 | 92,785 | 96.5 |
| Other | 2,621 | 4.7 | 219 | 2.4 | 230 | 0.9 | 291 | 4.2 | 3,361 | 3.5 |

Table 4.11. Hip. Primary and revision procedures by hospitalization burden. Details of the *Other* value of Table 4.10

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision | | TOTAL | |
|--|------------------------------|------|-------------------------------|------|---------------------|------|------------|------|--------------|------|
| | N | % | N | % | N | % | N | % | N | % |
| Hospitalization burden | 2,621 | | 219 | | 230 | | 291 | | 3,361 | |
| Patient's contribution for the accommodation payment | 913 | 34.8 | 30 | 13.7 | 63 | 27.4 | 77 | 26.5 | 1,083 | 32.2 |
| Reimbursement | 112 | 4.3 | 9 | 4.1 | 3 | 1.3 | 15 | 5.2 | 139 | 4.1 |
| Solvent | 1,102 | 42.0 | 82 | 37.4 | 84 | 36.5 | 142 | 48.8 | 1,410 | 42.0 |
| Surgeon's fee charged to patient | 183 | 7.0 | 17 | 7.8 | 27 | 11.7 | 13 | 4.5 | 240 | 7.1 |
| Surgeon's fee and accommodation charged to patient | 259 | 9.9 | 16 | 7.3 | 19 | 8.3 | 30 | 10.3 | 324 | 9.6 |
| Foreigners charged to NHS | 25 | 1.0 | 54 | 24.7 | 30 | 13.0 | 3 | 1.0 | 112 | 3.3 |
| Indigent foreigners charged to NHS | 24 | 0.9 | 8 | 3.7 | 1 | 0.4 | 2 | 0.7 | 35 | 1.0 |
| Other | 3 | 0.1 | 3 | 1.4 | 3 | 1.3 | 9 | 3.1 | 18 | 0.5 |

Table 4.12. Knee. Arthroplasties performed in the Italian regions (main and secondary procedures)

| | Total replacement | | Revision | | TOTAL | |
|-----------------------|-------------------|-------------|--------------|------------|---------------|--------------|
| | N | % | N | % | N | % |
| Italy | 62,886 | 93.4 | 4,479 | 6.6 | 67,365 | 100.0 |
| Piedmont | 4,509 | 7.2 | 320 | 7.1 | 4,829 | 7.2 |
| Valle d'Aosta | 121 | 0.2 | 1 | 0.0 | 122 | 0.2 |
| Lombardy | 13,586 | 21.6 | 1,160 | 25.9 | 14,746 | 21.9 |
| AP Bolzano | 800 | 1.3 | 58 | 1.3 | 858 | 1.3 |
| AP Trento | 564 | 0.9 | 31 | 0.7 | 595 | 0.9 |
| Veneto | 6,937 | 11.0 | 443 | 9.9 | 7,380 | 11.0 |
| Friuli Venezia Giulia | 1,640 | 2.6 | 98 | 2.2 | 1,738 | 2.6 |
| Liguria | 1,649 | 2.6 | 131 | 2.9 | 1,780 | 2.6 |
| Emilia-Romagna | 4,167 | 6.6 | 424 | 9.5 | 4,591 | 6.8 |
| Tuscany | 6,004 | 9.6 | 503 | 11.2 | 6,507 | 9.7 |
| Umbria | 1,291 | 2.1 | 95 | 2.1 | 1,386 | 2.1 |
| Marche | 1,617 | 2.6 | 76 | 1.7 | 1,693 | 2.5 |
| Lazio | 4,337 | 6.9 | 296 | 6.6 | 4,633 | 6.9 |
| Abruzzi | 1,739 | 2.8 | 85 | 1.9 | 1,824 | 2.7 |
| Molise | 229 | 0.4 | 8 | 0.2 | 237 | 0.4 |
| Campania | 2,963 | 4.7 | 160 | 3.6 | 3,123 | 4.6 |
| Apulia | 3,067 | 4.9 | 158 | 3.5 | 3,225 | 4.8 |
| Basilicata | 376 | 0.6 | 25 | 0.6 | 401 | 0.6 |
| Calabria | 1,135 | 1.8 | 53 | 1.2 | 1,188 | 1.8 |
| Sicily | 4,671 | 7.4 | 309 | 6.9 | 4,980 | 7.4 |
| Sardinia | 1,484 | 2.4 | 45 | 1.0 | 1,529 | 2.3 |

Table 4.13. Knee. Primary total replacement: number of hospitals by region and class of volume (main and secondary procedures)

| | Class of volume of total replacements | | | | | |
|-----------------------|---------------------------------------|------------|------------|-----------|-----------|------------|
| | 1-50 | 51-100 | 101-200 | 201-300 | >300 | TOTAL |
| Italy | 404 | 154 | 119 | 37 | 33 | 747 |
| Piedmont | 26 | 10 | 12 | 3 | 2 | 53 |
| Valle d'Aosta | 1 | 1 | 0 | 0 | 0 | 2 |
| Lombardy | 53 | 24 | 18 | 7 | 10 | 112 |
| AP Bolzano | 4 | 2 | 3 | 1 | 0 | 10 |
| AP Trento | 3 | 2 | 3 | 0 | 0 | 8 |
| Veneto | 20 | 12 | 10 | 2 | 5 | 49 |
| Friuli Venezia Giulia | 2 | 6 | 5 | 2 | 0 | 15 |
| Liguria | 7 | 4 | 3 | 0 | 2 | 16 |
| Emilia-Romagna | 34 | 16 | 5 | 0 | 4 | 59 |
| Tuscany | 19 | 12 | 10 | 5 | 5 | 51 |
| Umbria | 6 | 5 | 4 | 1 | 0 | 16 |
| Marche | 10 | 6 | 3 | 2 | 0 | 21 |
| Lazio | 50 | 15 | 12 | 1 | 1 | 79 |
| Abruzzi | 12 | 5 | 1 | 1 | 2 | 21 |
| Molise | 4 | 0 | 1 | 0 | 0 | 5 |
| Campania | 51 | 12 | 6 | 1 | 0 | 70 |
| Apulia | 26 | 4 | 7 | 4 | 0 | 41 |
| Basilicata | 5 | 1 | 2 | 0 | 0 | 8 |
| Calabria | 15 | 2 | 0 | 2 | 1 | 20 |
| Sicily | 43 | 12 | 11 | 3 | 1 | 70 |
| Sardinia | 13 | 3 | 3 | 2 | 0 | 21 |

Table 4.14. Knee. Revision: number of hospitals by region and class of volume (main and secondary procedures)

| | Class of volume of revisions | | | | |
|-----------------------|------------------------------|-----------|-----------|----------|------------|
| | 1-10 | 11-25 | 26-50 | >50 | TOTAL |
| Italy | 444 | 72 | 29 | 6 | 551 |
| Piedmont | 30 | 11 | 1 | 0 | 42 |
| Valle d'Aosta | 1 | 0 | 0 | 0 | 1 |
| Lombardy | 64 | 9 | 12 | 3 | 88 |
| AP Bolzano | 6 | 1 | 0 | 0 | 7 |
| AP Trento | 7 | 0 | 0 | 0 | 7 |
| Veneto | 30 | 10 | 2 | 1 | 43 |
| Friuli Venezia Giulia | 10 | 4 | 0 | 0 | 14 |
| Liguria | 9 | 1 | 2 | 0 | 12 |
| Emilia-Romagna | 37 | 5 | 2 | 1 | 45 |
| Tuscany | 29 | 6 | 6 | 1 | 42 |
| Umbria | 12 | 0 | 1 | 0 | 13 |
| Marche | 13 | 1 | 0 | 0 | 14 |
| Lazio | 43 | 5 | 2 | 0 | 50 |
| Abruzzi | 10 | 3 | 0 | 0 | 13 |
| Molise | 4 | 0 | 0 | 0 | 4 |
| Campania | 38 | 3 | 0 | 0 | 41 |
| Apulia | 28 | 4 | 0 | 0 | 32 |
| Basilicata | 6 | 0 | 0 | 0 | 6 |
| Calabria | 11 | 1 | 0 | 0 | 12 |
| Sicily | 43 | 8 | 1 | 0 | 52 |
| Sardinia | 13 | 0 | 0 | 0 | 13 |

Table 4.15. Knee. Distribution of arthroplasties (main procedure)

| ICD-9-CM codes | Arthroplasty | N | % |
|-----------------------|---|---------------|----------|
| 81.54 | Totale knee replacement | 61,923 | |
| | Revision | 3,884 | |
| 81.55 | Revision of knee replacement, not otherwise specified | 1,668 | 42.9 |
| 00.80 | Revision of knee replacement, total (all components) | 1,493 | 38.4 |
| 00.81 | Revision of knee replacement, tibial component | 191 | 4.9 |
| 00.82 | Revision of knee replacement, femoral component | 96 | 2.5 |
| 00.83 | Revision of knee replacement, patellar component | 238 | 6.1 |
| 00.84 | Revision of total knee replacement, tibial insert | 198 | 5.1 |

Table 4.16. Knee. Demographic characteristics by type of procedure

| | Total replacement | | Revision | | TOTAL | |
|----------------------|-------------------|------|----------|------|--------|------|
| | N | % | N | % | N | % |
| Patients | 60,855 | | 3,276 | | 64,131 | |
| Procedures | 61,923 | | 3,884 | | 65,807 | |
| Gender | | | | | | |
| Males | 19,574 | 31.6 | 1,155 | 29.7 | 20,729 | 31.5 |
| Females | 42,349 | 68.4 | 2,729 | 70.3 | 45,078 | 68.5 |
| Age by gender | | | | | | |
| Males | | | | | | |
| Mean age | 69.5 | | 67.9 | | 69.4 | |
| Standard deviation | 9.3 | | 11.3 | | 9.4 | |
| Median age | 71 | | 70 | | 71 | |
| Interquartile range | 65-76 | | 63-76 | | 65-76 | |
| Females | | | | | | |
| Mean age | 70.7 | | 70.2 | | 70.7 | |
| Standard deviation | 8.0 | | 9.5 | | 8.1 | |
| Median age | 72 | | 71 | | 72 | |
| Interquartile range | 66-76 | | 65-77 | | 66-76 | |
| Class of age | | | | | | |
| Males | | | | | | |
| <45 | 314 | 1.6 | 49 | 4.2 | 363 | 1.8 |
| 45 - 54 | 940 | 4.8 | 78 | 6.8 | 1,018 | 4.9 |
| 55 - 64 | 3,409 | 17.4 | 211 | 18.3 | 3,620 | 17.5 |
| 65 - 74 | 8,647 | 44.2 | 481 | 41.7 | 9,128 | 44.0 |
| 75 - 84 | 5,992 | 30.6 | 309 | 26.8 | 6,301 | 30.4 |
| ≥ 85 | 272 | 1.4 | 27 | 2.3 | 299 | 1.4 |
| Females | | | | | | |
| <45 | 217 | 0.5 | 42 | 1.5 | 259 | 0.6 |
| 45 - 54 | 1,243 | 2.9 | 111 | 4.1 | 1,354 | 3.0 |
| 55 - 64 | 6,888 | 16.3 | 467 | 17.1 | 7,355 | 16.3 |
| 65 - 74 | 19,383 | 45.8 | 1,153 | 42.3 | 20,536 | 45.6 |
| 75 - 84 | 13,925 | 32.9 | 879 | 32.2 | 14,804 | 32.8 |
| ≥ 85 | 693 | 1.6 | 77 | 2.8 | 770 | 1.7 |

Table 4.17. Knee. Arthroplasties by main diagnosis (ICD-9-CM)

| | Males | | Females | | TOTAL | |
|---|---------------|------|---------------|------|---------------|------|
| | N | % | N | % | N | % |
| Total replacement | 19,574 | | 42,349 | | 61,923 | |
| Osteoarthritis | 18,867 | 96.4 | 41,103 | 97.1 | 59,970 | 96.9 |
| Other acquired deformities of limbs | 259 | 1.3 | 578 | 1.4 | 837 | 1.4 |
| Other disorders of bone and cartilage | 81 | 0.4 | 166 | 0.4 | 247 | 0.4 |
| Complications peculiar to certain specified procedures | 48 | 0.3 | 106 | 0.3 | 154 | 0.3 |
| Other and unspecified disorders of joint | 39 | 0.2 | 108 | 0.3 | 147 | 0.2 |
| Internal derangement of knee | 74 | 0.4 | 52 | 0.1 | 126 | 0.2 |
| Other and unspecified arthropathies | 25 | 0.1 | 41 | 0.1 | 66 | 0.1 |
| Osteochondropathies | 39 | 0.2 | 23 | 0.1 | 62 | 0.1 |
| Malignant neoplasm of bone and articular cartilage | 27 | 0.1 | 20 | 0.1 | 47 | 0.1 |
| Other derangement of joint | 27 | 0.1 | 13 | 0.0 | 40 | 0.1 |
| Other diagnoses (N, 40, freq <0,1% on total) | 88 | 0.5 | 139 | 0.3 | 227 | 0.4 |
| Revision | 1,155 | | 2,729 | | 3,884 | |
| Complications peculiar to certain specified procedures | 938 | 81.2 | 2,251 | 82.5 | 3,189 | 82.1 |
| Osteoarthritis | 66 | 5.7 | 184 | 6.7 | 250 | 6.4 |
| Organ or tissue replaced by other means | 46 | 4.0 | 83 | 3.0 | 129 | 3.3 |
| Other disorders of bone and cartilage | 22 | 1.9 | 45 | 1.7 | 67 | 1.7 |
| Other orthopedic aftercare | 20 | 1.7 | 38 | 1.4 | 58 | 1.5 |
| Other derangement of joint | 11 | 1.0 | 27 | 1.0 | 38 | 1.0 |
| Internal derangement of knee | 9 | 0.8 | 23 | 0.8 | 32 | 0.8 |
| Arthropathy associated with infections | 11 | 1.0 | 6 | 0.2 | 17 | 0.4 |
| Other and unspecified arthropathies | 6 | 0.5 | 10 | 0.4 | 16 | 0.4 |
| Fracture of other and unspecified parts of femur | 2 | 0.2 | 10 | 0.4 | 12 | 0.3 |
| Other acquired deformities of limbs | 1 | 0.1 | 10 | 0.4 | 11 | 0.3 |
| Osteomyelitis, periostitis, and other infections involving bone | 7 | 0.6 | 4 | 0.2 | 11 | 0.3 |
| Other disorders of bone and cartilage | 1 | 0.1 | 5 | 0.2 | 6 | 0.2 |
| Fracture of tibia and fibula | 1 | 0.1 | 5 | 0.2 | 6 | 0.2 |

Follow

Table 4.17. Follow

| | Males | | Females | | TOTAL | |
|---|-------|-----|---------|-----|-------|-----|
| | N | % | N | % | N | % |
| Dislocation of knee | 1 | 0.1 | 5 | 0.2 | 6 | 0.2 |
| Malignant neoplasm of bone and articular cartilage | 1 | 0.1 | 4 | 0.2 | 5 | 0.1 |
| Bacterial infection in conditions classified elsewhere and of unspecified | 0 | 0.0 | 4 | 0.2 | 4 | 0.1 |
| Osteochondropathies | 0 | 0.0 | 4 | 0.2 | 4 | 0.1 |
| Sprains and strains of knee and leg | 2 | 0.2 | 1 | 0.0 | 3 | 0.1 |
| Other disorders of synovium, tendon, and bursa | 0 | 0.0 | 2 | 0.1 | 2 | 0.1 |
| Fitting and adjustment of other device | 1 | 0.1 | 1 | 0.0 | 2 | 0.1 |
| Fracture of patella | 0 | 0.0 | 2 | 0.1 | 2 | 0.1 |
| Late effects of musculoskeletal and connective tissue injuries | 1 | 0.1 | 1 | 0.0 | 2 | 0.1 |
| Septicemia | 1 | 0.1 | 1 | 0.0 | 2 | 0.1 |
| Other diagnoses (N, 10, freq <0,1% on total) | 7 | 0.6 | 3 | 0.1 | 10 | 0.3 |

Table 4.18. Knee. Primary and revision procedures by type of discharge

| | Total replacement | | Revision | | TOTAL | |
|---|-------------------|------|----------|------|--------|-------|
| | N | % | N | % | N | % |
| Type of discharge | 55,461 | 57.7 | 6,868 | 7.1 | 96,109 | 100.0 |
| Dead | 33 | 0.1 | 7 | 0.2 | 40 | 0.1 |
| Ordinary discharge | 32,601 | 52.7 | 2,162 | 55.7 | 34,763 | 52.8 |
| Discharge to a residential health care | 542 | 0.9 | 38 | 1.0 | 580 | 0.9 |
| Discharge to hospital at home | 83 | 0.1 | 2 | 0.1 | 85 | 0.1 |
| Discharge against medical advice | 65 | 0.1 | 4 | 0.1 | 69 | 0.1 |
| Transfer to an acute admission unit of a different hospital | 1,060 | 1.7 | 74 | 1.9 | 1,134 | 1.7 |
| Transfer in the same hospital | 15,784 | 25.5 | 1,016 | 26.2 | 16,800 | 25.5 |
| Transfer to an inpatient rehabilitation hospital | 11,577 | 18.7 | 570 | 14.7 | 12,147 | 18.5 |
| Discharge to a nursing home | 178 | 0.3 | 11 | 0.3 | 189 | 0.3 |

Table 4.19. Knee. Length of stay (days) by procedure and region

| Arthroplasty | Total replacement | | Revision | |
|-----------------------|-------------------|------------|------------|------------|
| | Mean | S.D. | Mean | S.D. |
| Italy | 7.7 | 3.7 | 9.8 | 7.6 |
| Piedmont | 7.5 | 3.8 | 10.1 | 8.3 |
| Valle d'Aosta | 5.1 | 3.3 | 4.0 | - |
| Lombardy | 6.8 | 3.3 | 7.6 | 5.2 |
| AP Bolzano | 9.3 | 3.2 | 12.0 | 7.3 |
| AP Trento | 7.2 | 3.9 | 11.0 | 8.8 |
| Veneto | 8.2 | 4.0 | 11.1 | 10.2 |
| Friuli Venezia Giulia | 8.8 | 3.8 | 11.9 | 7.4 |
| Liguria | 6.8 | 3.5 | 12.0 | 10.8 |
| Emilia-Romagna | 9.0 | 3.5 | 10.9 | 7.5 |
| Tuscany | 7.9 | 2.5 | 9.6 | 4.7 |
| Umbria | 7.3 | 3.3 | 10.3 | 7.3 |
| Marche | 9.3 | 3.5 | 11.6 | 7.7 |
| Lazio | 8.1 | 4.9 | 9.9 | 6.1 |
| Abruzzi | 7.1 | 3.0 | 8.1 | 5.5 |
| Molise | 6.6 | 2.7 | 13.7 | 6.2 |
| Campania | 8.5 | 3.8 | 10.7 | 6.2 |
| Apulia | 7.9 | 3.2 | 11.4 | 10.3 |
| Basilicata | 8.6 | 3.2 | 15.3 | 20.2 |
| Calabria | 8.1 | 3.7 | 9.8 | 11.1 |
| Sicily | 7.0 | 3.4 | 9.3 | 6.8 |
| Sardinia | 8.4 | 4.4 | 14.7 | 17.0 |

Table 4.20. Knee. Primary and revision procedures by hospitalization burden

| | Total replacement | | Revision | | TOTAL | |
|-------------------------------|-------------------|------|----------|------|---------------|------|
| | N | % | N | % | N | % |
| Hospitalization burden | | | | | 65,807 | |
| National Health System | 60,127 | 97.1 | 3,780 | 97.3 | 63,907 | 97.1 |
| Other | 1,796 | 2.9 | 104 | 2.7 | 1,900 | 2.9 |

Table 4.21. Knee. Primary and revision procedures by hospitalization burden. Details of the *Other* value of Table 4.20

| | Total replacement | | Revision | | TOTAL | |
|--|-------------------|------|------------|------|--------------|------|
| | N | % | N | % | N | % |
| Hospitalization burden | 1,796 | | 104 | | 1,900 | |
| Patient's contribution for the accommodation payment | 879 | 48.9 | 67 | 64.4 | 946 | 49.8 |
| Reimbursement | 75 | 4.2 | 2 | 1.9 | 77 | 4.1 |
| Solvent | 664 | 37.0 | 25 | 24.0 | 689 | 36.3 |
| Surgeon's fee charged to patient | 72 | 4.0 | 3 | 2.9 | 75 | 3.9 |
| Surgeon's fee and accomodation charged to patient | 78 | 4.3 | 2 | 1.9 | 80 | 4.2 |
| Foreigners charged to NHS | 17 | 1.0 | 4 | 3.9 | 21 | 1.1 |
| Indigent foreigners charged to NHS | 10 | 0.6 | 1 | 1.0 | 11 | 0.6 |
| Other | 1 | 0.1 | 0 | 0.0 | 1 | 0.1 |

Table 4.22. Shoulder. Arthroplasties performed in the Italian regions (main and secondary procedures)

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | TOTAL | |
|-----------------------|------------------------------|-------------|-------------------------------|-------------|---------------------|-------------|--------------|--------------|
| | N | % | N | % | N | % | N | % |
| Italy | 4,089 | 62.1 | 1,218 | 18.5 | 1,281 | 19.4 | 6,588 | 100.0 |
| Piedmont | 332 | 8.1 | 107 | 8.8 | 59 | 4.6 | 498 | 7.6 |
| Valle d'Aosta | 3 | 0.1 | 3 | 0.3 | 0 | 0.0 | 6 | 0.1 |
| Lombardy | 789 | 19.3 | 232 | 19.1 | 271 | 21.2 | 1,292 | 19.6 |
| AP Bolzano | 21 | 0.5 | 4 | 0.3 | 21 | 1.6 | 46 | 0.7 |
| AP Trento | 42 | 1.0 | 12 | 1.0 | 20 | 1.6 | 74 | 1.1 |
| Veneto | 432 | 10.6 | 173 | 14.2 | 240 | 18.7 | 845 | 12.8 |
| Friuli Venezia Giulia | 116 | 2.8 | 26 | 2.1 | 25 | 2.0 | 167 | 2.5 |
| Liguria | 99 | 2.4 | 32 | 2.6 | 19 | 1.5 | 150 | 2.3 |
| Emilia-Romagna | 550 | 13.5 | 113 | 9.3 | 101 | 7.9 | 764 | 11.6 |
| Tuscany | 418 | 10.2 | 73 | 6.0 | 104 | 8.1 | 595 | 9.0 |
| Umbria | 92 | 2.3 | 23 | 1.9 | 25 | 2.0 | 140 | 2.1 |
| Marche | 117 | 2.9 | 51 | 4.2 | 21 | 1.6 | 189 | 2.9 |
| Lazio | 338 | 8.3 | 75 | 6.2 | 104 | 8.1 | 517 | 7.9 |
| Abruzzi | 92 | 2.3 | 32 | 2.6 | 46 | 3.6 | 170 | 2.6 |
| Molise | 4 | 0.1 | 5 | 0.4 | 4 | 0.3 | 13 | 0.2 |
| Campania | 170 | 4.2 | 35 | 2.9 | 46 | 3.6 | 251 | 3.8 |
| Apulia | 182 | 4.5 | 114 | 9.4 | 68 | 5.3 | 364 | 5.5 |
| Basilicata | 21 | 0.5 | 7 | 0.6 | 9 | 0.7 | 37 | 0.6 |
| Calabria | 43 | 1.1 | 25 | 2.1 | 10 | 0.8 | 78 | 1.2 |
| Sicily | 199 | 4.9 | 68 | 5.6 | 51 | 4.0 | 318 | 4.8 |
| Sardinia | 29 | 0.7 | 8 | 0.7 | 37 | 2.9 | 74 | 1.1 |

Table 4.23. Shoulder. Primary total replacement: number of hospitals by region and class of volume (main and secondary procedures)

| | Class of volume of total replacements | | | | | |
|-----------------------|---------------------------------------|------------|-----------|-----------|-----------|------------|
| | 1-4 | 5-9 | 10-14 | 15-24 | ≥25 | TOTAL |
| Italy | 221 | 105 | 60 | 51 | 54 | 491 |
| Piedmont | 14 | 8 | 8 | 5 | 5 | 40 |
| Valle d'Aosta | 2 | 0 | 0 | 0 | 0 | 2 |
| Lombardy | 36 | 24 | 7 | 8 | 12 | 87 |
| AP Bolzano | 4 | 3 | 0 | 0 | 0 | 7 |
| AP Trento | 5 | 2 | 1 | 1 | 0 | 9 |
| Veneto | 7 | 6 | 9 | 12 | 7 | 41 |
| Friuli Venezia Giulia | 2 | 3 | 3 | 1 | 1 | 10 |
| Liguria | 8 | 1 | 1 | 1 | 1 | 12 |
| Emilia-Romagna | 17 | 8 | 8 | 5 | 7 | 45 |
| Tuscany | 14 | 8 | 2 | 2 | 7 | 33 |
| Umbria | 3 | 4 | 3 | 2 | 0 | 12 |
| Marche | 11 | 4 | 1 | 1 | 2 | 19 |
| Lazio | 26 | 6 | 5 | 4 | 4 | 45 |
| Abruzzi | 8 | 5 | 1 | 1 | 1 | 16 |
| Molise | 1 | 1 | 0 | 0 | 0 | 2 |
| Campania | 17 | 4 | 2 | 4 | 1 | 28 |
| Apulia | 12 | 8 | 3 | 2 | 3 | 28 |
| Basilicata | 2 | 1 | 0 | 1 | 0 | 4 |
| Calabria | 4 | 3 | 2 | 1 | 0 | 10 |
| Sicily | 21 | 5 | 3 | 0 | 3 | 32 |
| Sardinia | 7 | 1 | 1 | 0 | 0 | 9 |

Table 4.24. Shoulder. Primary partial replacement: number of hospitals by region and class of volume (main and secondary procedures)

| | Class of volume of partial replacements | | | | | |
|-----------------------|---|-----------|-----------|-----------|----------|------------|
| | 1-4 | 5-9 | 10-14 | 15-24 | ≥25 | TOTAL |
| Italy | 247 | 52 | 14 | 16 | 1 | 330 |
| Piedmont | 26 | 1 | 0 | 0 | 0 | 27 |
| Valle d'Aosta | 0 | 0 | 0 | 0 | 0 | 0 |
| Lombardy | 45 | 12 | 3 | 4 | 0 | 64 |
| AP Bolzano | 2 | 1 | 1 | 0 | 0 | 4 |
| AP Trento | 6 | 1 | 0 | 0 | 0 | 7 |
| Veneto | 18 | 5 | 4 | 7 | 0 | 34 |
| Friuli Venezia Giulia | 6 | 2 | 0 | 0 | 0 | 8 |
| Liguria | 3 | 2 | 0 | 0 | 0 | 5 |
| Emilia-Romagna | 22 | 3 | 0 | 2 | 0 | 27 |
| Tuscany | 14 | 3 | 0 | 1 | 1 | 19 |
| Umbria | 3 | 1 | 1 | 0 | 0 | 5 |
| Marche | 6 | 2 | 0 | 0 | 0 | 8 |
| Lazio | 27 | 6 | 1 | 0 | 0 | 34 |
| Abruzzi | 8 | 2 | 2 | 0 | 0 | 12 |
| Molise | 1 | 0 | 0 | 0 | 0 | 1 |
| Campania | 13 | 2 | 0 | 1 | 0 | 16 |
| Apulia | 15 | 3 | 1 | 1 | 0 | 20 |
| Basilicata | 4 | 0 | 0 | 0 | 0 | 4 |
| Calabria | 6 | 0 | 0 | 0 | 0 | 6 |
| Sicily | 16 | 4 | 0 | 0 | 0 | 20 |
| Sardinia | 6 | 2 | 1 | 0 | 0 | 9 |

Table 4.25. Shoulder. Distribution of arthroplasties (main procedure)

| | | N | % |
|----------------|-------------------------------------|--------------|------|
| ICD-9-CM codes | Arthroplasty | 6,323 | |
| 81.80 | Total shoulder replacement | 5,179 | |
| | Total replacement (elective) | 4,006 | 77.4 |
| | Total replacement (emergency) | 1,173 | 22.6 |
| 81.81 | Partial shoulder replacement | 1,144 | |

Table 4.26. Shoulder. Demographic characteristics by type of procedure

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | TOTAL | |
|----------------------|------------------------------|------|-------------------------------|------|---------------------|------|-------|------|
| | N | % | N | % | N | % | N | % |
| Patients | 3,965 | | 1,166 | | 1,113 | | 6,244 | |
| Procedures | 4,006 | | 1,173 | | 1,144 | | 6,323 | |
| Gender | | | | | | | | |
| Males | 1,171 | 29.2 | 170 | 14.5 | 384 | 33.6 | 1,725 | 27.3 |
| Females | 2,835 | 70.8 | 1,003 | 85.5 | 760 | 66.4 | 4,598 | 72.7 |
| Age by gender | | | | | | | | |
| Males | | | | | | | | |
| Mean age | 68.5 | | 72.3 | | 62.9 | | 67.6 | |
| Standard deviation | 9.4 | | 10.4 | | 13.6 | | 10.9 | |
| Median age | 70 | | 74 | | 64 | | 70 | |
| Interquartile range | 64-75 | | 67-79 | | 55-72 | | 62-75 | |
| Females | | | | | | | | |
| Mean age | 72.7 | | 75.0 | | 70.9 | | 72.9 | |
| Standard deviation | 7.4 | | 7.1 | | 10.8 | | 8.1 | |
| Median age | 74 | | 76 | | 73 | | 74 | |
| Interquartile range | 69-77 | | 71-80 | | 65-79 | | 69-78 | |
| Class of age | | | | | | | | |
| Males | | | | | | | | |
| <45 | 31 | 2.7 | 3 | 1.8 | 33 | 8.6 | 67 | 3.9 |
| 45 - 54 | 81 | 6.9 | 7 | 4.1 | 53 | 13.8 | 141 | 8.2 |
| 55 - 64 | 196 | 16.7 | 24 | 14.1 | 107 | 27.9 | 327 | 19.0 |
| 65 - 74 | 551 | 47.1 | 52 | 30.6 | 118 | 30.7 | 721 | 41.8 |
| 75 - 84 | 299 | 25.5 | 70 | 41.2 | 65 | 16.9 | 434 | 25.2 |
| ≥ 85 | 13 | 1.1 | 14 | 8.2 | 8 | 2.1 | 35 | 2.0 |
| Females | | | | | | | | |
| <45 | 17 | 0.6 | 0 | 0.0 | 16 | 2.1 | 33 | 0.7 |
| 45 - 54 | 36 | 1.3 | 11 | 1.1 | 42 | 5.5 | 89 | 1.9 |
| 55 - 64 | 254 | 9.0 | 68 | 6.8 | 114 | 15.0 | 436 | 9.5 |
| 65 - 74 | 1,261 | 44.5 | 357 | 35.6 | 275 | 36.2 | 1,893 | 41.2 |
| 75 - 84 | 1,194 | 42.1 | 499 | 49.8 | 266 | 35.0 | 1,959 | 42.6 |
| ≥ 85 | 73 | 2.6 | 68 | 6.8 | 47 | 6.2 | 188 | 4.1 |

Table 4.27. Shoulder. Arthroplasties by main diagnosis (ICD-9-CM)

| | Males | | Females | | TOTAL | |
|--|--------------|------|--------------|------|--------------|------|
| | N | % | N | % | N | % |
| Total replacement | 1,341 | | 3,838 | | 5,179 | |
| Osteoarthritis | 840 | 62.6 | 2,093 | 54.5 | 2,933 | 56.6 |
| Fracture of humerus | 168 | 12.5 | 986 | 25.7 | 1,154 | 22.3 |
| Other disorders of synovium, tendon, and bursa | 81 | 6.0 | 161 | 4.2 | 242 | 4.7 |
| Other and unspecified arthropathies | 48 | 3.6 | 113 | 2.9 | 161 | 3.1 |
| Other disorders of bone and cartilage | 39 | 2.9 | 82 | 2.1 | 121 | 2.3 |
| Complications peculiar to certain specified procedures | 32 | 2.4 | 72 | 1.9 | 104 | 2.0 |
| Other and unspecified disorders of joint | 25 | 1.9 | 76 | 2.0 | 101 | 2.0 |
| Peripheral enthesopathies and allied syndromes | 36 | 2.7 | 63 | 1.6 | 99 | 1.9 |
| Late effects of musculoskeletal and connective tissue injuries | 20 | 1.5 | 55 | 1.4 | 75 | 1.5 |
| Other derangement of joint | 15 | 1.1 | 38 | 1.0 | 53 | 1.0 |
| Fracture of scapula | 9 | 0.7 | 29 | 0.8 | 38 | 0.7 |
| Dislocation of shoulder | 3 | 0.2 | 29 | 0.8 | 32 | 0.6 |
| Organ or tissue replaced by other means | 5 | 0.4 | 6 | 0.2 | 11 | 0.2 |
| Other orthopedic aftercare | 5 | 0.4 | 5 | 0.1 | 10 | 0.2 |
| Rheumatoid arthritis and other inflammatory polyarthropathies | 3 | 0.2 | 4 | 0.1 | 7 | 0.1 |
| Sprains and strains of shoulder and upper arm | 2 | 0.2 | 4 | 0.1 | 6 | 0.1 |
| Secondary malignant neoplasm of other specified sites | 3 | 0.2 | 2 | 0.1 | 5 | 0.1 |
| Crystal arthropathies | 1 | 0.1 | 3 | 0.1 | 4 | 0.1 |
| Malignant neoplasm of bone and articular cartilage | 1 | 0.1 | 3 | 0.1 | 4 | 0.1 |
| Other diagnoses (N.14, freq <0.1% on total) | 5 | 0.4 | 14 | 0.4 | 19 | 0.4 |
| Partial replacement | 384 | | 760 | | 1,144 | |
| Fracture of humerus | 114 | 29.7 | 387 | 50.9 | 501 | 43.8 |
| Osteoarthritis | 97 | 25.3 | 184 | 24.2 | 281 | 24.6 |
| Other disorders of synovium, tendon, and bursa | 71 | 18.5 | 67 | 8.8 | 138 | 12.1 |
| Other disorders of bone and cartilage | 17 | 4.4 | 26 | 3.4 | 43 | 3.8 |
| Complications peculiar to certain specified procedures | 17 | 4.4 | 16 | 2.1 | 33 | 2.9 |
| Peripheral enthesopathies and allied syndromes | 15 | 3.9 | 13 | 1.7 | 28 | 2.5 |

Follow

Table 4.27. Follow

| | Males | | Females | | TOTAL | |
|---|-------|-----|---------|-----|-------|-----|
| | N | % | N | % | N | % |
| Sprains and strains of shoulder and upper arm | 6 | 1.6 | 10 | 1.3 | 16 | 1.4 |
| Dislocation of shoulder | 14 | 3.7 | 2 | 0.3 | 16 | 1.4 |
| Other and unspecified disorders of joint | 3 | 0.8 | 10 | 1.3 | 13 | 1.1 |
| Malignant neoplasm of bone and articular cartilage | 6 | 1.6 | 7 | 0.9 | 13 | 1.1 |
| Other and unspecified arthropathies | 5 | 1.3 | 3 | 0.4 | 8 | 0.7 |
| Late effects of musculoskeletal and connective tissue injuries | 1 | 0.3 | 6 | 0.8 | 7 | 0.6 |
| Other orthopedic aftercare | 5 | 1.3 | 1 | 0.1 | 6 | 0.5 |
| Other derangement of joint | 3 | 0.8 | 2 | 0.3 | 5 | 0.4 |
| Fracture of neck of femur | 0 | 0.0 | 5 | 0.7 | 5 | 0.4 |
| Organ or tissue replaced by other means | 1 | 0.3 | 4 | 0.5 | 5 | 0.4 |
| Fracture of scapula | 0 | 0.0 | 4 | 0.5 | 4 | 0.4 |
| Secondary malignant neoplasm of other specified sites | 3 | 0.8 | 1 | 0.1 | 4 | 0.4 |
| III-defined fractures of upper limb | 0 | 0.0 | 2 | 0.3 | 2 | 0.2 |
| Osteomyelitis, periostitis, and other infections involving bone | 1 | 0.3 | 1 | 0.1 | 2 | 0.2 |
| Certain adverse effects not elsewhere classified | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Other congenital anomalies of limbs | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Other and unspecified disorders of joint | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Sprains and strains of sacroiliac region | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Epilepsy | 1 | 0.3 | 0 | 0.0 | 1 | 0.1 |
| Fracture of radius and ulna | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Fracture of clavicle | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Dislocation of hip | 1 | 0.3 | 0 | 0.0 | 1 | 0.1 |
| Symptoms involving cardiovascular system | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Toxoplasmosis | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Benign neoplasm of bone and articular cartilage | 0 | 0.0 | 1 | 0.1 | 1 | 0.1 |
| Malignant neoplasm of connective and other soft tissue | 1 | 0.3 | 0 | 0.0 | 1 | 0.1 |
| Malignant neoplasm of trachea, bronchus, and lung | 1 | 0.3 | 0 | 0.0 | 1 | 0.1 |
| Burn of upper limb, except wrist and hand | 1 | 0.3 | 0 | 0.0 | 1 | 0.1 |

Table 4.28. Shoulder. Primary procedures by type of discharge

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | TOTAL | |
|---|------------------------------|------|-------------------------------|------|---------------------|------|-------|------|
| | N | % | N | % | N | % | N | % |
| Type of discharge | 4,006 | | 1,173 | | 1,144 | | 6,323 | |
| Dead | 1 | 0.0 | 3 | 0.3 | 0 | 0.0 | 4 | 0.1 |
| Ordinary discharge | 3,729 | 93.1 | 1,066 | 90.9 | 1,076 | 94.1 | 5,871 | 92.9 |
| Discharge to a residential health care | 8 | 0.2 | 17 | 1.5 | 7 | 0.6 | 32 | 0.5 |
| Discharge to hospital at home | 3 | 0.1 | 4 | 0.3 | 3 | 0.3 | 10 | 0.2 |
| Discharge against medical advice | 3 | 0.1 | 1 | 0.1 | 1 | 0.1 | 5 | 0.1 |
| Transfer to an acute admission unit of a different hospital | 20 | 0.5 | 13 | 1.1 | 11 | 1.0 | 44 | 0.7 |
| Transfer in the same hospital | 171 | 4.3 | 27 | 2.3 | 12 | 1.1 | 210 | 3.3 |
| Transfer to an inpatient rehabilitation hospital | 69 | 1.7 | 33 | 2.8 | 31 | 2.7 | 133 | 2.1 |
| Discharge to a nursing home | 2 | 0.1 | 9 | 0.8 | 3 | 0.3 | 14 | 0.2 |

Table 4.29. Shoulder. Length of stay (days) by procedure and region

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | |
|-----------------------|---------------------------------|------------|----------------------------------|------------|------------------------|------------|
| | Arthroplasty | 4,006 | | 1,173 | | 1,144 |
| | Mean | S.D. | Mean | S.D. | Mean | S.D. |
| Italy | 5.6 | 3.4 | 9.7 | 5.8 | 7.4 | 8.2 |
| Piedmont | 5.9 | 2.9 | 8.3 | 6.7 | 6.1 | 3.6 |
| Valle d'Aosta | 5.0 | 2.6 | 8.0 | 4.6 | - | - |
| Lombardy | 5.5 | 2.9 | 9.6 | 5.0 | 9.0 | 11.2 |
| AP Bolzano | 6.9 | 2.2 | 11.3 | 3.7 | 11.2 | 6.8 |
| AP Trento | 4.5 | 1.7 | 9.5 | 4.3 | 4.9 | 2.4 |
| Veneto | 6.8 | 4.2 | 11.0 | 6.4 | 5.5 | 6.5 |
| Friuli Venezia Giulia | 5.2 | 3.1 | 11.9 | 9.5 | 5.3 | 5.1 |
| Liguria | 3.4 | 4.0 | 11.3 | 7.5 | 7.7 | 3.2 |
| Emilia-Romagna | 5.0 | 2.5 | 9.0 | 5.9 | 7.0 | 4.3 |
| Tuscany | 4.8 | 1.4 | 8.2 | 5.6 | 6.3 | 3.8 |
| Umbria | 4.7 | 2.7 | 8.8 | 2.9 | 5.0 | 4.0 |
| Marche | 3.4 | 2.4 | 7.9 | 7.2 | 4.8 | 2.1 |
| Lazio | 6.5 | 4.9 | 10.7 | 6.2 | 10.0 | 14.5 |
| Abruzzi | 5.6 | 4.2 | 9.8 | 4.4 | 7.3 | 5.6 |
| Molise | 12.3 | 4.3 | 12.8 | 2.8 | 8.0 | 2.8 |
| Campania | 8.0 | 5.1 | 10.9 | 4.9 | 6.2 | 4.6 |
| Apulia | 5.7 | 3.2 | 9.4 | 3.9 | 7.4 | 7.2 |
| Basilicata | 4.5 | 2.0 | 9.8 | 6.2 | 10.3 | 5.6 |
| Calabria | 6.0 | 2.8 | 7.9 | 3.0 | 9.5 | 3.8 |
| Sicily | 5.4 | 3.4 | 10.9 | 5.7 | 8.0 | 3.6 |
| Sardinia | 7.4 | 3.9 | 10.8 | 3.3 | 10.1 | 7.6 |

Table 4.30. Shoulder. Primary procedures by hospitalization burden

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | TOTAL | |
|-------------------------------|---------------------------------|------|----------------------------------|------|------------------------|------|--------------|------|
| | N | % | N | % | N | % | N | % |
| Hospitalization burden | 4,006 | | 1,173 | | 1,144 | | 6,323 | |
| National Health System | 3,843 | 95.9 | 1,147 | 97.8 | 1,112 | 97.2 | 6,102 | 96.5 |
| Other | 163 | 4.1 | 26 | 2.2 | 32 | 2.8 | 221 | 3.5 |

Table 4.31. Shoulder. Primary procedures by hospitalization burden. Details of the *Other* value of Table 4.30

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | TOTAL | |
|--|---------------------------------|------|----------------------------------|------|------------------------|------|------------|------|
| | N | % | N | % | N | % | N | % |
| Hospitalization burden | 163 | | 26 | | 32 | | 221 | |
| Patient's contribution for the accommodation payment | 89 | 54.6 | 8 | 30.8 | 12 | 37.5 | 109 | 49.3 |
| Reimbursement | 4 | 2.5 | 0 | 0.0 | 0 | 0.0 | 4 | 1.8 |
| Solvent | 50 | 30.7 | 11 | 42.3 | 10 | 31.3 | 71 | 32.1 |
| Surgeon's fee charged to patient | 4 | 2.5 | 1 | 3.9 | 2 | 6.3 | 7 | 3.2 |
| Surgeon's fee and accomodation charged to patient | 16 | 9.8 | 5 | 19.2 | 4 | 12.5 | 25 | 11.3 |
| Foreigners charged to NHS | 0 | 0.0 | 1 | 3.9 | 3 | 9.4 | 4 | 1.8 |
| Indigent foreigners charged to NHS | 0 | 0.0 | 0 | 0.0 | 1 | 3.1 | 1 | 0.5 |

Table 5.1. Participation rate and completeness of the regions involved in the RIAP project (2015)

| Regions | Joint | N. hospitals RIAP | N. hospitals HDR | Participation (%) | N. hospital admissions RIAP | N. hospital admissions HDR | Completeness (%) |
|--------------|-------------|----------------------|---------------------|----------------------|-----------------------------------|----------------------------------|---------------------|
| Lombardy | Hip | 112 | 112 | 100.0 | 20,359 | 20,651 | 98.6 |
| | Knee | 112 | 112 | 100.0 | 14,467 | 14,581 | 99.2 |
| AP Bolzano | Hip | 8 | 10 | 80.0 | 1,492 | 1,529 | 97.6 |
| | Knee | 8 | 10 | 80.0 | 917 | 958 | 95.7 |
| AP Trento | Hip | 7 | 8 | 87.5 | 837 | 1,082 | 77.4 |
| | Knee | 5 | 8 | 62.5 | 40 | 620 | 6.5 |
| Tuscany | Hip | 1 | 45 | 2.2 | 383 | 8,840 | 4.3 |
| | Knee | 1 | 45 | 2.2 | 156 | 7,072 | 2.2 |
| Marche | Hip | 12 | 19 | 63.2 | 1,028 | 2,402 | 42.8 |
| | Knee | 12 | 21 | 57.1 | 861 | 1,603 | 53.7 |
| Abruzzi | Hip | 2 | 22 | 9.1 | 90 | 2,242 | 4.0 |
| | Knee | 2 | 22 | 9.1 | 32 | 1,903 | 1.7 |
| Apulia | Hip | 43 | 43 | 100.0 | 4,636 | 4,657 | 99.5 |
| | Knee | 43 | 43 | 100.0 | 3,286 | 3,287 | 100.0 |
| Basilicata | Hip | 2 | 6 | 33.3 | 348 | 657 | 53.0 |
| | Knee | 2 | 6 | 33.3 | 301 | 417 | 72.2 |
| Calabria | Hip | 11 | 20 | 55.0 | 690 | 1,712 | 40.3 |
| | Knee | 10 | 20 | 50.0 | 671 | 1,228 | 54.6 |
| Sicily | Hip | 8 | 72 | 11.1 | 637 | 5,479 | 11.6 |
| | Knee | 8 | 71 | 11.3 | 778 | 4,809 | 16.2 |
| Total | Hip | 206 | 357 | 57.7 | 30,500 | 49,251 | 61.9 |
| | Knee | 203 | 358 | 56.7 | 21,509 | 36,478 | 59.0 |

Participation rate: number of hospitals participating in the RIAP out of number of hospitals included in the Hospital Discharge Records (HDR)
 Completeness: number of admissions recorded in RIAP out of number of admissions recorded in HDR

Table 5.2. Hip. Completeness of the hospitals participating in the RIAP project (2015)

| Regions | Hospitals | Completeness (%) |
|----------------|--|-------------------------|
| Lombardy | Casa di Cura S. Francesco, Bergamo | 100.6 |
| | Casa di Cura Beato Luigi Talamoni, Lecco | 100.0 |
| | Casa di Cura Beato Palazzolo, Bergamo | 100.0 |
| | Casa di Cura Figlie di S. Camillo, Cremona | 100.0 |
| | Casa di Cura G.B. Mangioni, Lecco | 100.0 |
| | Casa di Cura Igea, Milano | 100.0 |
| | Casa di Cura Policlinico, Monza | 100.0 |
| | Casa di Cura S. Maria delle Grazie, Voghera (PV) | 100.0 |
| | Casa di Cura S. Pio X, Milano | 100.0 |
| | Clinica San Carlo, Paderno Dugnano (MI) | 100.0 |
| | IRCCS Policlinico San Donato Milanese (MI) | 100.0 |
| | Istituti Clinici Zucchi Spa, Monza | 100.0 |
| | Istituti Ospitalieri, Cremona | 100.0 |
| | Istituto Clinico S.Ambrogio Spa, Milano | 100.0 |
| | Istituto Clinico S. Siro Spa, Milano | 100.0 |
| | Istituto Clinico Beato Matteo, Vigevano (PV) | 100.0 |
| | Istituto Clinico Città di Brescia, Brescia | 100.0 |
| | Istituto Clinico Città Studi, Milano | 100.0 |
| | Istituto Clinico S. Anna, Brescia | 100.0 |
| | Istituto Clinico S. Rocco Spa, Ome (BS) | 100.0 |
| | Istituto Clinico Villa Aprica Spa, Como | 100.0 |
| | Nuovo Ospedale di Brond-Stradella (PV) | 100.0 |
| | Ospedale Caduti Bollatesi, Bollate (MI) | 100.0 |
| | Ospedale Carlo Ondoli, Angera (VA) | 100.0 |
| | Ospedale Civico di Codogno (LO) | 100.0 |
| | Ospedale Civile di Voghera (PV) | 100.0 |
| | Ospedale Civile G. Fornaroli, Magenta (MI) | 100.0 |
| | Ospedale Civile La Memoria di Gavardo (BS) | 100.0 |
| | Ospedale Costantino Cantù, Abbiategrasso (MI) | 100.0 |
| | Ospedale di Chiavenna (SO) | 100.0 |

Follow

Table 5.2. Follow

| Regions | Hospitals | Completeness (%) |
|-------------------|--|-------------------------|
| Lombardy (follow) | Ospedale di Circolo, Rho (MI) | 100.0 |
| | Ospedale di Cuggiono (MI) | 100.0 |
| | Ospedale di Desenzano (BS) | 100.0 |
| | Ospedale di Manerbio (BS) | 100.0 |
| | Ospedale di Treviglio-Caravaggio (BG) | 100.0 |
| | Ospedale G. Salvini, Garbagnate Milanese (MI) | 100.0 |
| | Ospedale Galmarini, Tradate (VA) | 100.0 |
| | Ospedale Generale Provinciale, Saronno (VA) | 100.0 |
| | Ospedale L. Sacco, Milano | 100.0 |
| | Ospedale M.O. Antonio Locatelli, Piaro (BG) | 100.0 |
| | Ospedale Morelli, Sondalo (SO) | 100.0 |
| | Ospedale Moriggia Pelascini, Gravedona (CO) | 100.0 |
| | Ospedale Pesenti Fenaroli, Alzano Lombardo (BG) | 100.0 |
| | Ospedale S. Antonio Abate, Cantù (CO) | 100.0 |
| | Ospedale S. Gerardo, Monza | 100.0 |
| | Ospedale Sacra Famiglia - Fatebenefratelli, Erba (CO) | 100.0 |
| | Ospedale San Pellegrino, Castiglione delle Stiviere (MN) | 100.0 |
| | Ospedale SS. Capitanio e Gerosa, Lovere (BG) | 100.0 |
| | Ospedale SS. Trinità, Romano di Lombardia (BG) | 100.0 |
| | Ospedale Valcamonica, Edolo (BS) | 100.0 |
| | Ospedale Valcamonica, Esine (BS) | 100.0 |
| | PO Centro Traumatologico Ortopedico, Milano | 100.0 |
| | PO di Carate Brianza (MB) | 100.0 |
| | PO di Chiari (BS) | 100.0 |
| | PO di Gardone Val Trompia (BS) | 100.0 |
| | PO di Iseo (BS) | 100.0 |
| | PO di Montichiari (BS) | 100.0 |
| | PO Spedali Civili di Brescia | 100.0 |
| | Policlinico S. Matteo, Pavia | 100.0 |
| | Policlinico San Marco, Osio Sotto (BG) | 100.0 |
| | | <i>Follow</i> |

Table 5.2. Follow

| Regions | Hospitals | Completeness (%) |
|-------------------|--|-------------------------|
| Lombardy (follow) | Policlinico San Pietro, Ponte S. Pietro (BG) | 100.0 |
| | Polo Chirurgico Capitanio, Milano | 100.0 |
| | Ospedale di Suzzara (MN) | 99.7 |
| | Casa di Cura Mater Domini, Castellanza (VA) | 99.7 |
| | Fondazione Poliambulanza, Brescia | 99.6 |
| | Istituto Ortopedico Galeazzi Spa, Milano | 99.6 |
| | Ospedale di Circolo, Busto Arsizio (VA) | 99.6 |
| | Ospedale di Circolo A. Manzoni, Lecco | 99.5 |
| | Ospedale di Sondrio | 99.4 |
| | Ospedale Civile di Legnano (MI) | 99.4 |
| | Ospedale Papa Giovanni XXIII, Bergamo | 99.4 |
| | Istituto Clinico Humanitas, Rozzano (MI) | 99.3 |
| | Ospedale Civile di Vigevano (PV) | 99.3 |
| | Casa di Cura S. Clemente, Mantova | 99.1 |
| | Ospedale Maggiore, Crema (CR) | 99.1 |
| | PO di Oglio PO, Casalmaggiore (CR) | 99.0 |
| | IRCCS Multimedica, Milano | 99.0 |
| | Ospedale S. Antonio Abate, Gallarate (VA) | 98.9 |
| | Ospedale Niguarda Ca' Granda, Milano | 98.7 |
| | Ospedale Bolognini, Seriate (BG) | 98.6 |
| | Casa di Cura S. Camillo, Cremona | 98.6 |
| | Ospedale di Circolo S.L. Mandic, Merate (LC) | 98.6 |
| | IRCCS S. Raffaele, Milano | 98.5 |
| | Ospedale Luini Confalonieri, Luino (VA) | 98.4 |
| | Ospedale di Vizzolo Predabissi (MI) | 98.2 |
| | Ospedale S. Maria delle Stelle, Melzo (MI) | 98.2 |
| | Istituto di Cura Città di Pavia, Pavia | 98.1 |
| | Ospedale Ubaldo, Cernusco sul Naviglio (MI) | 97.7 |
| | Ospedale Causa Pia Luvini di Cittiglio, Cittiglio (VA) | 97.7 |
| | Ospedale Civile di S. Giovanni Bianco (BG) | 97.6 |

Follow

Table 5.2. Follow

| Regions | Hospitals | Completeness (%) |
|-------------------|---|-------------------------|
| Lombardy (follow) | PO di Desio (MB) | 97.5 |
| | Ospedale Erba-Renaldi-Menaggio (CO) | 97.4 |
| | PO di Giussano (MB) | 96.5 |
| | Ospedale di Circolo e Fondazione Macchi, Varese | 96.3 |
| | PO di Vimercate (MB) | 95.9 |
| | Ospedale Fatebenefratelli e Oftalmico, Milano | 95.8 |
| | Cliniche Gavazzeni Spa, Bergamo | 95.7 |
| | Istituto Ortopedico Gaetano Pini, Milano | 95.7 |
| | Ospedale S. Paolo, Milano | 93.1 |
| | Casa di Cura La Cittadella Sociale, Pieve del Cairo (PV) | 92.6 |
| | Clinica Castelli, Bergamo | 92.6 |
| | Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Milano | 92.6 |
| | Ospedale S. Carlo Borromeo, Milano | 92.1 |
| | Casa di Cura S. Camillo, Brescia | 91.3 |
| | Casa di Cura S. Maria di Castellanza (VA) | 90.9 |
| | Ospedale S. Anna, Como | 90.8 |
| | Casa di Cura S. Giovanni, Milano | 90.6 |
| | Ospedale Carlo Poma, Mantova | 90.1 |
| | Ospedale Civile Destra Secchia, Pieve di Coriano (MN) | 89.6 |
| | Ospedale Maggiore di Lodi | 81.8 |
| | Ospedale Bassini, Cinisello Balsamo (MI) | 80.0 |
| | COF Lanzo Hospital, Ramponio Verna (CO) | 77.0 |
| AP Bolzano | Ospedale di San Candido | 100.0 |
| | Ospedale di Vipiteno | 100.0 |
| | Ospedale di Bolzano | 99.6 |
| | Ospedale di Brunico | 98.8 |
| | Ospedale di Bressanone | 96.6 |
| | Casa di Cura Privata S. Maria | 96.3 |
| | Ospedale di Merano | 95.9 |
| | Ospedale di Silandro | 92.7 |

Follow

Table 5.2. Follow

| Regions | Hospitals | Completeness (%) |
|----------------|---|-------------------------|
| AP Trento | Ospedale di Cavalese | 100.0 |
| | Ospedale San Camillo, Trento | 99.4 |
| | Ospedale di Trento | 95.8 |
| | Ospedale di Cles | 94.0 |
| | Ospedale di Rovereto | 79.0 |
| | Ospedale di Tione | 69.4 |
| | Ospedale di Borgo Valsugana | 61.5 |
| Tuscany | AOU Pisana, Pisa | 62.2 |
| Marche | Ospedale Civile E. Profili, Fabriano (AN) | 95.5 |
| | Ospedale di Fermo, Fermo | 94.9 |
| | Casa di Cura Villa Serena, Jesi (AN) | 94.7 |
| | Casa di Cura Villa Pini Sanatrix Gestione, Civitanova Marche (MC) | 93.7 |
| | Casa di Cura Villa Igea, Ancona | 80.2 |
| | PO ZT4, Senigallia (AN) | 77.4 |
| | Ospedale S. Maria della Misericordia, Urbino | 64.7 |
| | PO Umberto I, Ancona | 47.3 |
| | Ospedale San Salvatore, Pesaro | 28.5 |
| | Ospedali Riuniti di Jesi, Jesi (AN) | 28.1 |
| Abruzzi | Casa di cura Villa San Marco, Ascoli Piceno | 13.0 |
| | Ospedale Generale Di Zona, Civitanova Marche (MC) | 12.7 |
| | PO Clinicizzato SS. Annunziata, Chieti | 34.4 |
| | Ospedale G. Bernabeo, Ortona | 19.0 |
| | PO G. Tatarella, Cerignola (FG) | 100.0 |
| | PO S. Caterina Novella, Galatina (LE) | 100.0 |
| | PO Sacro Cuore di Gesù, Gallipoli (LE) | 100.0 |
| | PO Orientale, Manduria (TA) | 100.0 |
| | PO Valle d'Itria, Martina Franca (TA) | 100.0 |
| | EE Miulli, Acquaviva (BA) | 100.0 |
| Apulia | EE Cardinale G. Panico, Triggiano (BA) | 100.0 |
| | Casa di Cura S. Maria, Bari | 100.0 |

Follow

Table 5.2. Follow

| Regions | Hospitals | Completeness (%) |
|-----------------|---|-------------------------|
| Apulia (follow) | Casa Di Cura Villa Lucia, Conversano (BA) | 100.0 |
| | Casa di Cura Villa Bianca, Lecce | 100.0 |
| | Casa di Cura Bernardini, Taranto | 100.0 |
| | Casa di Cura F. D'Amore, Taranto | 100.0 |
| | Casa di Cura San Camillo, Taranto | 100.0 |
| | Casa di Cura Anthea, Bari | 100.0 |
| | Casa Bianca Hospital, Cassano | 100.0 |
| | CBH Presidio Mater Dei, Bari | 100.0 |
| | Casa di Cura Città di Lecce | 100.0 |
| | PO Umberto I, Altamura (BA) | 100.0 |
| | PO San Paolo, Bari | 100.0 |
| | PO Umberto I, Corato (BA) | 100.0 |
| | PO Don Tonino Bello, Molfetta (BA) | 100.0 |
| | PO San Giacomo, Monopoli (BA) | 100.0 |
| | PO S. Maria degli Angeli, Putignano (BA) | 100.0 |
| | PO Ostuni (BR) | 100.0 |
| | PO Camberlingo, Francavilla-Ceglie (BR) | 100.0 |
| | PO T. Masselli Mascia, San Severo (FG) | 100.0 |
| | PO S. Camillo De Lellis, Manfredonia (FG) | 100.0 |
| | PO S. Giuseppe da Copertino, Copertino-Nardò (LE) | 100.0 |
| | PO Veris dellì Ponti, Scorrano (LE) | 100.0 |
| | PO F. Ferrari, Casarano (LE) | 100.0 |
| | PO Occidentale, Castellaneta (TA) | 100.0 |
| | PO Di Venere, Bari | 100.0 |
| | PO A. Perrino, Brindisi | 100.0 |
| | PO Vito Fazzi, Lecce | 100.0 |
| | PO SS. Annunziata, Taranto | 100.0 |
| | PO L. Bonomo, Andria-Canosa (BT) | 100.0 |
| | PO Monsignor R. Dimiccoli, Barletta | 100.0 |
| | PO Bisceglie, Trani (BT) | 100.0 |

Follow

Table 5.2. Follow

| Regions | Hospitals | Completeness (%) |
|-----------------|---|-------------------------|
| Apulia (follow) | Cliniche Riunite Villa Serena e San Francesco, Foggia | 100.0 |
| | PO San Marco, Grottaglie (TA) | 100.0 |
| | IRCCS Casa Sollievo della Sofferenza, San Giovanni Rotondo (FG) | 100.0 |
| | AOU Policlinico, Bari | 100.0 |
| | AOU Ospedali Riuniti, Foggia | 100.0 |
| Basilicata | AOR San Carlo (PZ) | 71.2 |
| | Ospedale Madonna Delle Grazie (MT) | 33.9 |
| Calabria | AO Mater Domini P.U., Catanzaro | 100.0 |
| | Casa di Cura Villa Serena, Catanzaro | 100.0 |
| | PO Lamezia Terme (CZ) | 100.0 |
| | Casa di Cura Scarnati, Cosenza | 96.0 |
| | Casa di Cura Villa Caminiti, Reggio Calabria | 96.0 |
| | PO Rossano-Corigliano (CS) | 89.0 |
| | Istituto Ortopedico del Mezzogiorno d'Italia, Reggio Calabria | 76.0 |
| | PO S. Maria degli Ungheresi, Polistena (RC) | 62.0 |
| | Casa di Cura Cascini, Cosenza | 49.0 |
| | AO di Cosenza | 32.7 |
| Sicily | AO Pugliese Ciaccio, Catanzaro | 4.0 |
| | IMOI F. Scalabrino Ganzirri, Messina | 100.0 |
| | San Raffaele-G. Giglio, Cefalù (PA) | 86.6 |
| | Ospedale Maria Paternò Arezzo, Ragusa | 86.0 |
| | PO M. Chiello, Piazza Armerina (EN) | 72.1 |
| | PO Umberto I, Enna | 56.2 |
| | PO Gravina e S. Pietro, Caltagirone (CT) | 40.7 |
| | Policlinico Giaccone, Palermo | 13.4 |
| | Casa di Cura S. Barbara Sogesa, Gela (CL) | 8.7 |

Table 5.3. Knee. Completeness of the hospitals participating in the RIAP project (2015)

| Regions | Hospitals | Completeness (%) |
|----------------|--|-------------------------|
| Lombardy | Casa di Cura Beato Palazzolo, Bergamo | 102.5 |
| | Casa di Cura Beato Luigi Talamoni, Lecco | 100.0 |
| | Casa di Cura Igea, Milano | 100.0 |
| | Casa di Cura S. Camillo, Cremona | 100.0 |
| | Casa di Cura S. Maria delle Grazie, Voghera (PV) | 100.0 |
| | Casa di Cura S. Pio X, Milano | 100.0 |
| | Clinica San Carlo, Paderno Dugnano (MI) | 100.0 |
| | Istituto Clinico S. Ambrogio Spa, Milano | 100.0 |
| | Istituto Clinico Beato Matteo, Vigevano (PV) | 100.0 |
| | Istituto Clinico Città Studi, Milano | 100.0 |
| | Nuovo Ospedale di Broni-Stradella (PV) | 100.0 |
| | Ospedale Bassini, Cinisello Balsamo (MI) | 100.0 |
| | Ospedale Caduti Bollatesi, Bollate (MI) | 100.0 |
| | Ospedale Carlo Poma, Mantova | 100.0 |
| | Ospedale Civico di Codogno (LO) | 100.0 |
| | Ospedale Civile di S. Giovanni Bianco (BG) | 100.0 |
| | Ospedale Civile di Vigevano (PV) | 100.0 |
| | Ospedale Civile di Voghera (PV) | 100.0 |
| | Ospedale Civile G. Fornaroli, Magenta (MI) | 100.0 |
| | Ospedale Civile La Memoria di Gavardo (BS) | 100.0 |
| | Ospedale Costantino Cantù, Abbiategrasso (MI) | 100.0 |
| | Ospedale di Chiavenna (SO) | 100.0 |
| | Ospedale di Circolo A. Manzoni, Lecco | 100.0 |
| | Ospedale di Circolo, Busto Arsizio (VA) | 100.0 |
| | Ospedale di Circolo, Rho (MI) | 100.0 |
| | Ospedale di Cuggiono (MI) | 100.0 |
| | Ospedale di Desenzano (BS) | 100.0 |
| | Ospedale di Manerbio (BS) | 100.0 |
| | Ospedale di Sondrio | 100.0 |
| | Ospedale di Treviglio-Caravaggio (BG) | 100.0 |

Follow

Table 5.3. Follow

| Regions | Hospitals | Completeness (%) |
|-------------------|---|-------------------------|
| Lombardy (follow) | Ospedale di Vizzolo Predabissi (MI) | 100.0 |
| | Ospedale Fatebenefratelli e Oftalmico, Milano | 100.0 |
| | Ospedale G. Salvini, Garbagnate Milanese (MI) | 100.0 |
| | Ospedale Galmarini, Tradate (VA) | 100.0 |
| | Ospedale Generale Provinciale, Saronno (VA) | 100.0 |
| | Ospedale L. Sacco, Milano | 100.0 |
| | Ospedale Luini Confalonieri, Luino (VA) | 100.0 |
| | Ospedale M.O. Antonio Locatelli, Pario (BG) | 100.0 |
| | Ospedale Maggiore, Crema (CR) | 100.0 |
| | Ospedale Niguarda Ca' Granda, Milano | 100.0 |
| | Ospedale Pesenti Fenaroli, Alzano Lombardo (BG) | 100.0 |
| | Ospedale S. Antonio Abate, Gallarate (VA) | 100.0 |
| | Ospedale S. Gerardo, Monza | 100.0 |
| | Ospedale S. Maria delle Stelle, Melzo (MI) | 100.0 |
| | Ospedale SS. Capitanio e Gerosa, Lovere (BG) | 100.0 |
| | Ospedale SS. Trinità, Romano di Lombardia (BG) | 100.0 |
| | Ospedale Uboldo, Cernusco sul Naviglio (MI) | 100.0 |
| | Ospedale Valcamonica, Edolo (BS) | 100.0 |
| | Ospedale Valcamonica, Esine (BS) | 100.0 |
| | PO di Carate Brianza (MB) | 100.0 |
| | PO di Chiari (BS) | 100.0 |
| | PO di Desio (MB) | 100.0 |
| | PO di Gardone Val Trompia (BS) | 100.0 |
| | PO di Giussano (MB) | 100.0 |
| | PO di Montichiari (BS) | 100.0 |
| | PO di Oglio PO, Casalmaggiore (CR) | 100.0 |
| | PO Spedali Civili di Brescia | 100.0 |
| | Policlinico S. Matteo, Pavia | 100.0 |
| | Policlinico San Marco, Osio Sotto (BG) | 100.0 |
| | Polo Chirurgico Capitanio, Milano | 100.0 |

Follow

Table 5.3. Follow

| Regions | Hospitals | Completeness (%) |
|-------------------|--|-------------------------|
| Lombardy (follow) | Casa di Cura Policlinico, Monza | 99.9 |
| | Istituto Clinico Humanitas, Rozzano (MI) | 99.8 |
| | Istituti Clinici Zucchi Spa, Monza | 99.8 |
| | Fondazione Poliambulanza, Brescia | 99.7 |
| | Istituto Clinico Villa Aprica Spa, Como | 99.6 |
| | Casa di Cura Mater Domini, Castellanza (VA) | 99.4 |
| | Istituto Clinico S. Rocco Spa, Ome (BS) | 99.4 |
| | Policlinico San Pietro, Ponte S. Pietro (BG) | 99.3 |
| | IRCCS Policlinico San Donato Milanese (MI) | 99.3 |
| | Istituto Ortopedico Galeazzi Spa, Milano | 99.3 |
| | Ospedale San Pellegrino, Castiglione delle Stiviere (MN) | 99.2 |
| | Casa di Cura S. Francesco, Bergamo | 99.1 |
| | Istituto Clinico S. Siro Spa, Milano | 99.0 |
| | Casa di Cura S. Clemente, Mantova | 99.0 |
| | PO Centro Traumatologico Ortopedico, Milano | 99.0 |
| | Ospedale di Suzzara (MN) | 99.0 |
| | Casa di Cura Figlie di S. Camillo, Cremona | 98.9 |
| | COF Lanzo Hospital, Ramponio Verna (CO) | 98.9 |
| | Ospedale Civile di Legnano (MI) | 98.7 |
| | Ospedale Papa Giovanni XXIII, Bergamo | 98.6 |
| | Ospedale Carlo Ondoli, Angera (VA) | 98.6 |
| | Istituto Clinico S. Anna, Brescia | 98.4 |
| | Ospedale di Circolo S.L. Mandic, Merate (LC) | 98.2 |
| | Istituti Ospitalieri, Cremona | 98.1 |
| | Istituto Clinico Città di Brescia, Brescia | 98.1 |
| | PO di Iseo (BS) | 97.0 |
| | Ospedale Morelli, Sondalo (SO) | 96.8 |
| | PO di Vimercate (MB) | 96.7 |
| | Istituto di Cura Città di Pavia, Pavia | 96.6 |
| | Ospedale Sacra Famiglia - Fatebenefratelli, Erba (CO) | 96.5 |

Follow

Table 5.3. Follow

| Regions | Hospitals | Completeness (%) |
|-------------------|---|-------------------------|
| Lombardy (follow) | Ospedale di Circolo e Fondazione Macchi, Varese | 95.8 |
| | Ospedale Civile Destra Secchia, Pieve di Coriano (MN) | 95.7 |
| | Cliniche Gavazzeni Spa, Bergamo | 95.1 |
| | Casa di Cura S. Camillo, Brescia | 94.1 |
| | Ospedale S. Anna, Como | 94.1 |
| | IRCCS Multimedica, Milano | 93.8 |
| | Casa di Cura S. Giovanni, Milano | 93.3 |
| | Clinica Castelli, Bergamo | 92.3 |
| | IRCCS S. Raffaele, Milano | 92.3 |
| | Ospedale S. Antonio Abate, Cantù (CO) | 92.3 |
| | Casa di Cura G.B. Mangioni, Lecco | 91.8 |
| | Ospedale Erba-Renaldi-Menaggio (CO) | 91.7 |
| | Istituto Ortopedico Gaetano Pini, Milano | 91.2 |
| | Ospedale Bolognini, Seriate (BG) | 90.4 |
| | Casa di Cura S. Maria di Castellanza (VA) | 90.3 |
| | Ospedale Maggiore di Lodi | 90.0 |
| | Ospedale S. Paolo, Milano | 89.5 |
| | Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinico, Milano | 87.5 |
| | Ospedale Moriggia Pelascini, Gravedona (CO) | 85.7 |
| | Ospedale Causa Pia Luvini di Cittiglio, Cittiglio (VA) | 83.0 |
| | Casa di Cura La Cittadella Sociale, Pieve del Cairo (PV) | 78.4 |
| | Ospedale S. Carlo Borromeo, Milano | 66.7 |
| AP Bolzano | Casa di Cura Privata S. Maria | 100.0 |
| | Ospedale di Bolzano | 100.0 |
| | Ospedale di Vipiteno | 100.0 |
| | Ospedale di Bressanone | 99.4 |
| | Ospedale di San Candido | 98.6 |
| | Ospedale di Brunico | 97.7 |
| | Ospedale di Silandro | 94.3 |
| | Ospedale di Merano | 93.3 |

Follow

Table 5.3. Follow

| Regions | Hospitals | Completeness (%) |
|----------------|---|-------------------------|
| AP Trento | Ospedale di Trento | 13.5 |
| | Ospedale di Rovereto | 10.3 |
| | Ospedale San Camillo, Trento | 6.8 |
| | Ospedale di Tione | 1.8 |
| Tuscany | Ospedale di Cavalese | 1.5 |
| | A.O.U. Pisana, Pisa | 40.3 |
| | Ospedale Civile E. Profili, Fabriano (AN) | 100.0 |
| | Casa di Cura Villa Serena, Jesi (AN) | 93.8 |
| Marche | Casa di Cura Villa Pini Sanatrix Gestione, Civitanova Marche (MC) | 90.2 |
| | Ospedale di Fermo, Fermo | 87.9 |
| | Ospedale S. Maria della Misericordia, Urbino | 87.3 |
| | Casa di Cura Villa Igea, Ancona | 83.5 |
| | PO ZT4, Senigallia (AN) | 68.3 |
| | Ospedale Generale Di Zona, Civitanova Marche (MC) | 66.7 |
| | Presidio Ospedaliero Umberto I°, Ancona | 66.7 |
| | Ospedale San Salvatore, Pesaro | 60.7 |
| | Ospedali Riuniti di Jesi, Jesi (AN) | 28.8 |
| | Casa di cura Villa San Marco, Ascoli Piceno | 11.8 |
| Abruzzi | PO Clinicizzato SS. Annunziata, Chieti | 41.4 |
| | Ospedale G. Bernabeo, Ortona | 17.4 |
| Apulia | PO G. Tatarella, Cerignola (FG) | 100.0 |
| | PO S. Caterina Novella, Galatina (LE) | 100.0 |
| | PO Sacro Cuore di Gesù, Gallipoli (LE) | 100.0 |
| | PO Orientale, Manduria (TA) | 100.0 |
| | PO Valle d'Itria, Martina Franca (TA) | 100.0 |
| | EE Miulli, Acquaviva (BA) | 100.0 |
| | EE Cardinale G. Panico, Triggiano (BA) | 100.0 |
| | Casa di Cura S. Maria, Bari | 100.0 |
| | Casa Di Cura Villa Lucia, Conversano (BA) | 100.0 |
| | Casa di Cura Villa Bianca, Lecce | 100.0 |

Follow

Table 5.3. Follow

| Regions | Hospitals | Completeness (%) |
|-----------------|---|-------------------------|
| Apulia (follow) | Casa di Cura Bernardini, Taranto | 100.0 |
| | Casa di Cura F. D'Amore, Taranto | 100.0 |
| | Casa di Cura San Camillo, Taranto | 100.0 |
| | Casa di Cura Anthea, Bari | 100.0 |
| | Casa Bianca Hospital, Cassano | 100.0 |
| | CBH Presidio Mater Dei, Bari | 100.0 |
| | Casa di Cura Città di Lecce | 100.0 |
| | PO Umberto I, Altamura (BA) | 100.0 |
| | PO San Paolo, Bari | 100.0 |
| | PO Umberto I, Corato (BA) | 100.0 |
| | PO Don Tonino Bello, Molfetta (BA) | 100.0 |
| | PO San Giacomo, Monopoli (BA) | 100.0 |
| | PO S. Maria degli Angeli, Putignano (BA) | 100.0 |
| | PO Ostuni (BR) | 100.0 |
| | PO Camberlingo, Francavilla-Ceglie (BR) | 100.0 |
| | PO T. Masselli Mascia, San Severo (FG) | 100.0 |
| | PO S. Camillo De Lellis, Manfredonia (FG) | 100.0 |
| | PO S. Giuseppe da Copertino, Copertino-Nardò (LE) | 100.0 |
| | PO Veris delli Ponti, Scorrano (LE) | 100.0 |
| | PO F. Ferrari, Casarano (LE) | 100.0 |
| | PO Occidentale, Castellaneta (TA) | 100.0 |
| | PO Di Venere, Bari | 100.0 |
| | PO A. Perrino, Brindisi | 100.0 |
| | PO Vito Fazzi, Lecce | 100.0 |
| | PO SS. Annunziata, Taranto | 100.0 |
| | PO L. Bonomo, Andria-Canosa (BT) | 100.0 |
| | PO Monsignor R. Dimiccoli, Barletta | 100.0 |
| | PO Bisceglie, Trani (BT) | 100.0 |
| | Cliniche Riunite Villa Serena e San Francesco, Foggia | 100.0 |
| | PO San Marco, Grotttaglie (TA) | 100.0 |

Follow

Table 5.3. Follow

| Regions | Hospitals | Completeness (%) |
|----------------|---|-------------------------|
| Basilicata | IRCCS Casa Sollievo della Sofferenza, San Giovanni Rotondo (FG) | 100.0 |
| | AOU Policlinico, Bari | 100.0 |
| | AOU Ospedali Riuniti, Foggia | 100.0 |
| | Ospedale Madonna Delle Grazie (MT) | 88.5 |
| | AOR San Carlo (PZ) | 56.6 |
| | Casa di Cura Villa Caminiti, Reggio Calabria | 100.0 |
| | Istituto Ortopedico del Mezzogiorno d'Italia, Reggio Calabria | 100.0 |
| | PO Lamezia Terme (CZ) | 100.0 |
| | PO Rossano-Corigliano (CS) | 100.0 |
| Calabria | Casa di Cura Scarnati, Cosenza | 98.0 |
| | Casa di Cura Villa Serena, Catanzaro | 95.0 |
| | AO Mater Domini P.U., Catanzaro | 93.0 |
| | PO S. Maria degli Ungheresi, Polistena (RC) | 40.0 |
| | Casa di Cura Cascini, Cosenza | 37.0 |
| | AO Pugliese Ciaccio, Catanzaro | 8.0 |
| | OMI F. Scalabrin Ganzirri, Messina | 100.0 |
| | PO Gravina e S. Pietro, Caltagirone (CT) | 93.3 |
| | Ospedale Maria Paternò Arezzo, Ragusa | 81.4 |
| Sicily | COT - Cure Ortopediche Traumatologiche, Messina | 68.7 |
| | San Raffaele-G. Giglio, Cefalù (PA) | 66.2 |
| | Casa di Cura S. Barbara Sogesa, Gela (CL) | 55.4 |
| | Ospedale Generale di Zona, Lentini (SR) | 13.2 |
| | PO M. Chiello, Piazza Armerina (EN) | 6.0 |

Table 5.4. Procedures distribution. Comparison between 2014 HDR data and 2015 evaluable RIAP data

| | HDR (2014) | | RIAP (2015 evaluable data) | |
|-----------------------------------|----------------|-------------|----------------------------|-------------|
| | N | % | N | % |
| Total procedures | 166,862 | | 48,135 | |
| Hip | 99,497 | 59.6 | 28,415 | 59.0 |
| Total replacement and resurfacing | 66,115 | 66.4 | 19,911 | 70.1 |
| Total replacement (elective) | 56,561 | 85.5 | 17,854 | 89.7 |
| Total replacement (emergency) | 9,484 | 14.3 | 2,057 | 10.3 |
| Partial replacement | 25,313 | 25.4 | 6,831 | 24.0 |
| Revision | 8,069 | 8.1 | 1,673 | 5.9 |
| Knee | 67,365 | 40.4 | 19,720 | 41.0 |
| Total replacement | 62,886 | 93.4 | 18,793 | 95.3 |
| Revision | 4,479 | 6.6 | 927 | 4.7 |

Table 5.5. Hip. Type of procedure

| | N | % |
|---|---------------|-------------|
| Total procedures | 28,415 | |
| Total replacement | 19,911 | 70.2 |
| Total replacement (elective) | 17,854 | 89.7 |
| Total replacement (emergency) | 2,057 | 10.3 |
| Partial replacement | 6,831 | 24.0 |
| Revision (total or partial) | 1,319 | 4.6 |
| Removal of prosthesis | 348 | 1.2 |
| Conversion from partial to total replacement | 2 | 0.0 |
| Spacer revision | 4 | 0.0 |

Table 5.6. Hip. Procedures by type of provider

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision (*) | | TOTAL | |
|-------------------------|---------------------------------|------|----------------------------------|------|---------------------|------|--------------|------|--------|------|
| | N | % | N | % | N | % | N | % | N | % |
| Provider | 17,854 | | 2,057 | | 6,831 | | 1,673 | | 28,415 | |
| Public | 6,613 | 37.0 | 1,494 | 72.6 | 5,355 | 78.4 | 607 | 36.3 | 14,069 | 49.5 |
| Private, accredited | 11,154 | 62.5 | 558 | 27.1 | 1,473 | 21.6 | 1,056 | 63.1 | 14,241 | 50.1 |
| Private, not accredited | 87 | 0.5 | 5 | 0.2 | 3 | 0.0 | 10 | 0.6 | 105 | 0.4 |

(*) Revision (total or partial), removal of prosthesis, conversion from partial to total replacement, spacer revision

Table 5.7. Hip. Demographic characteristics by type of procedure

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision (*) | | TOTAL | |
|-------------------------------|------------------------------|------|-------------------------------|------|---------------------|------|--------------|------|---------------|------|
| | N | % | N | % | N | % | N | % | N | % |
| Gender | 17,854 | | 2,057 | | 6,831 | | 1,673 | | 28,415 | |
| Males | 8,103 | 45.4 | 601 | 29.2 | 1,926 | 28.2 | 647 | 38.7 | 11,277 | 39.7 |
| Females | 9,751 | 54.6 | 1,456 | 70.8 | 4,905 | 71.8 | 1,026 | 61.3 | 17,138 | 60.3 |
| Age by gender(**) | | | | | | | | | | |
| Males | | | | | | | | | | |
| Mean age | 65.2 | | 70.4 | | 80.6 | | 70.1 | | 68.5 | |
| Standard deviation | 12.1 | | 12.4 | | 12.1 | | 12.4 | | 13.5 | |
| Median age | 67 | | 72 | | 84 | | 72 | | 70 | |
| Interquartile range | 58-74 | | 63-79 | | 77-88 | | 63-79 | | 60-78 | |
| Females | | | | | | | | | | |
| Mean age | 69.3 | | 73.1 | | 84.0 | | 72.3 | | 74.1 | |
| Standard deviation | 11.1 | | 9.4 | | 7.4 | | 11.2 | | 11.9 | |
| Median age | 71 | | 74 | | 85 | | 74 | | 76 | |
| Interquartile range | 63-77 | | 67-79 | | 80-89 | | 65-81 | | 67-83 | |
| Class of age by gender | | | | | | | | | | |
| Males | 8,103 | | 601 | | 1,926 | | 647 | | 11,277 | |
| <45 | 477 | 5.9 | 22 | 3.7 | 34 | 1.8 | 18 | 2.8 | 551 | 4.9 |
| 45 - 54 | 988 | 12.2 | 37 | 6.2 | 79 | 4.1 | 65 | 10.0 | 1,169 | 10.4 |
| 55 - 64 | 1,708 | 21.1 | 99 | 16.5 | 85 | 4.4 | 88 | 13.6 | 1,980 | 17.6 |
| 65 - 74 | 2,557 | 31.6 | 185 | 30.8 | 148 | 7.7 | 173 | 26.7 | 3,063 | 27.2 |
| 75 - 84 | 1,758 | 21.7 | 174 | 29.0 | 656 | 34.1 | 192 | 29.7 | 2,780 | 24.7 |
| ≥ 85 | 127 | 1.6 | 64 | 10.6 | 863 | 44.8 | 55 | 8.5 | 1,109 | 9.8 |
| n.a. | 488 | 6.0 | 20 | 3.3 | 61 | 3.2 | 56 | 8.7 | 625 | 5.5 |
| Females | 9,751 | | 1,456 | | 4,905 | | 1,026 | | 17,138 | |
| <45 | 260 | 2.7 | 8 | 0.5 | 8 | 0.2 | 12 | 1.2 | 288 | 1.7 |
| 45 - 54 | 688 | 7.1 | 40 | 2.7 | 19 | 0.4 | 58 | 5.7 | 805 | 4.7 |
| 55 - 64 | 1,610 | 16.5 | 178 | 12.2 | 54 | 1.1 | 139 | 13.5 | 1,981 | 11.6 |
| 65 - 74 | 3,207 | 32.9 | 532 | 36.5 | 315 | 6.4 | 275 | 26.8 | 4,329 | 25.3 |
| 75 - 84 | 2,999 | 30.8 | 493 | 33.9 | 1,930 | 39.3 | 353 | 34.4 | 5,775 | 33.7 |
| ≥ 85 | 398 | 4.1 | 151 | 10.4 | 2,435 | 49.6 | 109 | 10.6 | 3,093 | 18.0 |
| n.a. | 589 | 6.0 | 54 | 3.7 | 144 | 2.9 | 80 | 7.8 | 867 | 5.1 |

(*) Revision (total or partial), removal of prosthesis, conversion from partial to total replacement, spacer revision

(**) AP Bolzano data have not been included because not available

Table 5.8. Hip. Surgical practice for primary procedures

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Revision (*) | | TOTAL | |
|---|------------------------------|------|-------------------------------|------|---------------------|------|--------------|------|---------------|------|
| | N | % | N | % | N | % | N | % | N | % |
| Operated side | 17,854 | | 2,057 | | 6,831 | | 1,673 | | 28,415 | |
| Right | 9,708 | 54.4 | 1,033 | 50.2 | 3,377 | 49.4 | 872 | 52.1 | 14,990 | 52.8 |
| Left | 7,880 | 44.1 | 1,020 | 49.6 | 3,446 | 50.4 | 793 | 47.4 | 13,139 | 46.2 |
| Bilateral | 266 | 1.5 | 4 | 0.2 | 8 | 0.1 | 8 | 0.5 | 286 | 1.0 |
| Approach | 17,854 | | 2,057 | | 6,831 | | 1,673 | | 28,415 | |
| Anterior | 2,666 | 14.9 | 102 | 5.0 | 270 | 4.0 | 83 | 5.0 | 3,121 | 11.0 |
| Anterolateral | 2,158 | 12.1 | 502 | 24.4 | 1,647 | 24.1 | 233 | 13.9 | 4,540 | 16.0 |
| Lateral | 3,510 | 19.7 | 517 | 25.1 | 2,128 | 31.2 | 370 | 22.1 | 6,525 | 23.0 |
| Posterolateral | 9,373 | 52.5 | 932 | 45.3 | 2,727 | 39.9 | 971 | 58.0 | 14,003 | 49.3 |
| Other | 147 | 0.8 | 4 | 0.2 | 59 | 0.9 | 16 | 1.0 | 226 | 0.8 |
| Fixation | 17,854 | | 2,057 | | 6,831 | | 1,673 | | 28,415 | |
| Cemented (stem + cup) | 758 | 4.2 | 149 | 7.2 | 302 | 4.4 | 75 | 4.5 | 1,284 | 4.5 |
| Cementless (stem + cup) | 15,941 | 89.3 | 1,552 | 75.4 | 2,321 | 34.0 | 1,051 | 62.8 | 20,865 | 73.4 |
| Hybrid (cemented stem and cementless cup) | 387 | 2.2 | 205 | 10.0 | 2,008 | 29.4 | 46 | 2.7 | 2,646 | 9.3 |
| Reverse hybrid (cementless stem and cemented cup) | 125 | 0.7 | 66 | 3.2 | 85 | 1.2 | 85 | 5.1 | 361 | 1.3 |
| Only cemented stem | 5 | 0.0 | 8 | 0.4 | 915 | 13.4 | 31 | 1.9 | 959 | 3.4 |
| Only cementless stem | 39 | 0.2 | 17 | 0.8 | 931 | 13.6 | 109 | 6.5 | 1,096 | 3.9 |
| Only cemented cup | 17 | 0.1 | 0 | 0.0 | 7 | 0.1 | 32 | 1.9 | 56 | 0.2 |
| Only cementless cup | 371 | 2.1 | 25 | 1.2 | 128 | 1.9 | 140 | 8.4 | 664 | 2.3 |
| Incorrectly coded | 211 | 1.2 | 35 | 1.7 | 134 | 2.0 | 104 | 6.2 | 484 | 1.7 |

(*) Revision (total or partial), removal of prosthesis, conversion from partial to total replacement, spacer revision

Note: incorrectly coded data could be included because hip partial replacement does not foresee cup implant

Table 5.9. Hip. Primary replacements by diagnosis and previous surgery

| | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | TOTAL | |
|--|------------------------------|------|-------------------------------|------|---------------------|------|---------------|------|
| | N | % | N | % | N | % | N | % |
| Diagnosis | 17,854 | | 2,057 | | 6,831 | | 26,742 | |
| Primary osteoarthritis | 15,380 | 86.1 | | | 283 | 4.1 | 15,663 | 58.6 |
| Post-traumatic osteoarthritis | 355 | 2.0 | | | 41 | 0.6 | 396 | 1.5 |
| Rheumatoid arthritis | 90 | 0.5 | | | 0 | 0.0 | 90 | 0.3 |
| Neoplasia | 41 | 0.2 | | | 27 | 0.4 | 68 | 0.3 |
| Aseptic necrosis of femoral head | 745 | 4.2 | | | 18 | 0.3 | 763 | 2.9 |
| Congenital dislocation/hip dysplasia | 613 | 3.4 | | | 7 | 0.1 | 620 | 2.3 |
| Perthes disease or epiphysiolyis | 39 | 0.2 | | | 0 | 0.0 | 39 | 0.1 |
| Fractured neck of femur | 0 | 0.0 | 1,994 | 96.9 | 6,306 | 92.3 | 8,300 | 31.0 |
| Septic coxitis | 1 | 0.0 | | | 0 | 0.0 | 1 | 0.0 |
| Pseudoarthrosis neck fracture related | 16 | 0.1 | | | 9 | 0.1 | 25 | 0.1 |
| Other | 219 | 1.2 | | | 71 | 1.0 | 290 | 1.1 |
| Previous surgery inconsistent with primary procedure (a) | 141 | 0.8 | 13 | 0.6 | 40 | 0.6 | 194 | 0.7 |
| Previous surgery incorrectly coded | 214 | 1.2 | 50 | 2.4 | 29 | 0.4 | 293 | 1.1 |
| Previous surgery | 17,854 | | 2,057 | | 6,831 | | 26,742 | |
| None | 15,778 | 88.4 | 1,832 | 89.1 | 6,090 | 89.2 | 23,700 | 88.6 |
| Osteosynthesis | 228 | 1.3 | 33 | 1.6 | 46 | 0.7 | 307 | 1.1 |
| Osteotomy | 87 | 0.5 | 2 | 0.1 | 1 | 0.0 | 90 | 0.3 |
| Arthrodesis | 2 | 0.0 | 0 | 0.0 | 0 | 0.0 | 2 | 0.0 |
| Other | 1,404 | 7.9 | 127 | 6.2 | 625 | 9.1 | 2,156 | 8.1 |
| Previous surgery inconsistent with primary procedure (a) | 141 | 0.8 | 13 | 0.6 | 40 | 0.6 | 194 | 0.7 |
| Previous surgery incorrectly coded | 214 | 1.2 | 50 | 2.4 | 29 | 0.4 | 293 | 1.1 |

(a) Inconsistent previous surgery: total or partial replacement, spacer or prosthesis removal (Girdlestone)

Table 5.10. Hip. Revision procedures by diagnosis and previous surgery

| Diagnosis | Revision (*) | |
|---|--------------|------|
| | N | % |
| Diagnosis | 1,673 | |
| Pain | 88 | 5.3 |
| Osteolysis | 53 | 3.2 |
| Wear | 106 | 6.3 |
| Implant breakage acetabulum | 4 | 0.2 |
| Implant breakage acetabular insert | 4 | 0.2 |
| Implant breakage head | 4 | 0.2 |
| Implant breakage modular neck | 3 | 0.2 |
| Implant breakage stem | 3 | 0.2 |
| Implant breakage spacer | 0 | 0.0 |
| Prosthesis dislocation | 156 | 9.3 |
| Periprosthetic fracture | 164 | 9.8 |
| Infection | 102 | 6.1 |
| Prosthesis removal | 23 | 1.4 |
| Acetabular cup aseptic loosening | 353 | 21.1 |
| Femoral stem aseptic loosening | 169 | 10.1 |
| Total aseptic loosening | 113 | 6.8 |
| Osteoarthritis progression | 0 | 0.0 |
| Other | 80 | 4.8 |
| Previous surgery inconsistent with revision (a) | 202 | 12.1 |
| Diagnosis incorrectly coded | 46 | 2.7 |
| Previous surgery | 1,673 | |
| Total hip replacement | 1,143 | 68.3 |
| Revision of hip replacement | 65 | 3.9 |
| Spacer or prosthesis removal | 127 | 7.6 |
| Partial hip replacement | 94 | 5.6 |
| Other | 41 | 2.5 |
| Previous surgery inconsistent with revision (a) | 202 | 12.1 |
| Previous surgery incorrectly coded | 1 | 0.1 |

(*) Revision (total or partial), removal of prosthesis, conversion from partial to total replacement, spacer revision

(a) Inconsistent previous surgery: none, osteosynthesis, ostectomy, arthrodesis

Table 5.11. Knee. Type of procedure

| | N | % |
|--|---------------|-------------|
| Total procedures | 19,720 | |
| Primary | 18,793 | 95.3 |
| Total | 15,858 | 84.4 |
| Unicondylar | 2,935 | 15.6 |
| Revision | 927 | 4.7 |
| Total revision | 268 | 28.9 |
| Partial revision | 623 | 67.2 |
| Primary patello-femoral implant on pre-existing prosthesis | 14 | 1.5 |
| Removal of prosthesis | 17 | 1.8 |
| Spacer revision | 5 | 0.5 |

Table 5.12. Knee. Procedures by type of provider

| Provider | Total replacement | | Revision (*) | | TOTAL | |
|-------------------------|-------------------|------|--------------|------|---------------|-------------|
| | N | % | N | % | N | % |
| Provider | 18,793 | | 927 | | 19,720 | |
| Public | 5,144 | 27.4 | 236 | 25.5 | 5,380 | 27.3 |
| Private, accredited | 13,600 | 72.4 | 688 | 74.2 | 14,288 | 72.5 |
| Private, not accredited | 49 | 0.3 | 3 | 0.3 | 52 | 0.3 |

(*) Total or partial revision, primary patello-femoral implant on pre-existing prosthesis, removal of prosthesis, spacer revision

Table 5.13. Knee. Demographic characteristics by type of procedure

| | Primary | | Revision (*) | | TOTAL | |
|-------------------------------|---------------|------|--------------|------|---------------|------|
| | N | % | N | % | N | % |
| Gender | 18,793 | | 927 | | 19,720 | |
| Males | 5,840 | 31.1 | 277 | 29.9 | 6,117 | 31.0 |
| Females | 12,953 | 68.9 | 650 | 70.1 | 13,603 | 69.0 |
| Age by gender(**) | | | | | | |
| Males | 5,626 | | 342 | | 5,968 | |
| Mean age | 69.4 | | 68.6 | | 69.4 | |
| Standard deviation | 9.2 | | 10.0 | | 9.3 | |
| Median age | 71 | | 70 | | 71 | |
| Interquartile range | 65-76 | | 64-76 | | 63-77 | |
| Females | 12,661 | | 817 | | 13,478 | |
| Mean age | 70.7 | | 70.5 | | 70.7 | |
| Standard deviation | 8.2 | | 9.0 | | 8.2 | |
| Median age | 72 | | 71 | | 74 | |
| Interquartile range | 66-76 | | 66-77 | | 67-80 | |
| Class of age by gender | | | | | | |
| Males | 5,840 | | 277 | | 6,117 | |
| <45 | 72 | 1.2 | 3 | 1.1 | 75 | 1.2 |
| 45 - 54 | 289 | 4.9 | 19 | 6.9 | 308 | 5.0 |
| 55 - 64 | 1,000 | 17.1 | 48 | 17.3 | 1,048 | 17.1 |
| 65 - 74 | 2,373 | 40.6 | 108 | 39.0 | 2,481 | 40.6 |
| 75 - 84 | 1,673 | 28.6 | 70 | 25.3 | 1,743 | 28.5 |
| ≥ 85 | 84 | 1.4 | 4 | 1.4 | 88 | 1.4 |
| n.a. | 349 | 6.0 | 25 | 9.0 | 374 | 6.1 |
| Females | 12,953 | | 650 | | 13,603 | |
| <45 | 54 | 0.4 | 6 | 0.9 | 60 | 0.4 |
| 45 - 54 | 422 | 3.3 | 27 | 4.2 | 449 | 3.3 |
| 55 - 64 | 1,986 | 15.3 | 102 | 15.7 | 2,088 | 15.3 |
| 65 - 74 | 5,497 | 42.4 | 260 | 40.0 | 5,757 | 42.3 |
| 75 - 84 | 4,301 | 33.2 | 211 | 32.5 | 4,512 | 33.2 |
| ≥ 85 | 183 | 1.4 | 11 | 1.7 | 194 | 1.4 |
| n.a. | 510 | 3.9 | 33 | 5.1 | 543 | 4.0 |

(*) Total or partial revision, primary patello-femoral implant on pre-existing prosthesis, removal of prosthesis, spacer revision

(**) AP Bolzano data have not been included because not available

Table 5.14. Knee. Surgical practice for primary procedures

| | Primary | | Revision (*) | | TOTAL | |
|---|---------------|------|--------------|------|---------------|------|
| | N | % | N | % | N | % |
| Operated side | 18,793 | | 927 | | 19,720 | |
| Right | 10,002 | 53.2 | 472 | 50.9 | 10,474 | 53.1 |
| Left | 8,617 | 45.9 | 451 | 48.7 | 9,068 | 46.0 |
| Bilateral | 174 | 0.9 | 4 | 0.4 | 178 | 0.9 |
| Approach | 18,793 | | 927 | | 19,720 | |
| Medial parapatellar | 16,387 | 87.2 | 826 | 89.1 | 17,213 | 87.3 |
| Lateral parapatellar | 409 | 2.2 | 16 | 1.7 | 425 | 2.2 |
| Mid-vastus | 754 | 4.0 | 33 | 3.6 | 787 | 4.0 |
| Minimally invasive Mid-Vastus | 701 | 3.7 | 25 | 2.7 | 726 | 3.7 |
| Quad-sparing | 69 | 0.4 | 1 | 0.1 | 70 | 0.4 |
| Sub-Vastus | 181 | 1.0 | 9 | 1.0 | 190 | 1.0 |
| Minimally invasive Sub-Vastus | 39 | 0.2 | 3 | 0.3 | 42 | 0.2 |
| V Quadriceps | 1 | 0.0 | 1 | 0.1 | 2 | 0.0 |
| Tibial tuberosity osteotomy | 25 | 0.1 | 7 | 0.8 | 32 | 0.2 |
| Other | 227 | 1.2 | 6 | 0.6 | 233 | 1.2 |
| Fixation | 18,793 | | 927 | | 19,720 | |
| Cemented | 13,132 | 69.9 | 439 | 47.4 | 13,571 | 68.8 |
| Cementless | 3,619 | 19.3 | 216 | 23.3 | 3,835 | 19.4 |
| Hybrid (cementless femoral component and cemented tibial component) | 773 | 4.1 | 60 | 6.5 | 833 | 4.2 |
| Only patella | 7 | 0.0 | 33 | 3.6 | 40 | 0.2 |
| Other | 475 | 2.5 | 24 | 2.6 | 499 | 2.5 |
| Inconsistent with the type of procedure(**) | 787 | 4.2 | 155 | 16.7 | 942 | 4.8 |

(*) Total or partial revision, primary patello-femoral implant on pre-existing prosthesis, removal of prosthesis, spacer revision

(**) Primary: only cemented femoral component, only cementless femoral component, only cemented tibial component, only cementless tibial component, no fixed component. Revision: no fixed component

Table 5.15. Knee. Primary replacements by diagnosis and previous surgery

| | Primary | |
|--|---------------|------|
| | N | % |
| Diagnosis | 18,793 | |
| Primary osteoarthritis | 17,247 | 91.8 |
| Post-traumatic osteoarthritis | 272 | 1.4 |
| Rheumatoid arthritis | 125 | 0.7 |
| Neoplasia | 15 | 0.1 |
| Osteonecrosis | 199 | 1.1 |
| Other | 473 | 2.5 |
| Previous surgery inconsistent with primary procedure (a) | 462 | 2.5 |
| Previous surgery | 18,793 | |
| None | 16,272 | 86.6 |
| Osteotomy | 3 | 0.0 |
| Arthroscopy | 134 | 0.7 |
| Arthrodesis | 479 | 2.5 |
| Other | 1,443 | 7.7 |
| Previous surgery inconsistent with primary procedure (a) | 462 | 2.5 |

(a) Inconsistent previous surgery: primary total, primary unicondylar, prosthesis reimplantation, spacer

Table 5.16. Knee. Revision procedures by diagnosis and previous surgery

| | Revision (*) | |
|---|--------------|------|
| | N | % |
| Diagnosis | 927 | |
| Aseptic loosening (more components) | 211 | 22.8 |
| Femoral component aseptic loosening | 42 | 4.5 |
| Tibial component aseptic loosening | 91 | 9.8 |
| Patellar component aseptic loosening | 1 | 0.1 |
| Wear | 31 | 3.3 |
| Dislocation | 20 | 2.2 |
| Instability | 37 | 4.0 |
| Periprosthetic fracture | 14 | 1.5 |
| Implant breakag prosthesis | 9 | 1.0 |
| Implant breakag spacer | 0 | 0.0 |
| Infection | 143 | 15.4 |
| Rigidity | 22 | 2.4 |
| Osteoarthritis progression | 19 | 2.0 |
| Pain | 205 | 22.1 |
| Other | 15 | 1.6 |
| Previous surgery inconsistent with revision (a) | 55 | 5.9 |
| Previous surgery incorrectly coded | 12 | 1.3 |
| Previous surgery | 1,021 | |
| Total replacement | 565 | 60.9 |
| Primary unicondylar | 119 | 12.8 |
| Removal of prosthesis | 59 | 6.4 |
| Spacer revision | 56 | 6.0 |
| Other | 61 | 6.6 |
| Previous surgery inconsistent with revision (a) | 55 | 5.9 |
| Previous surgery incorrectly coded | 12 | 1.3 |

(*) Total or partial revision, primary patello-femoral implant on pre-existing prosthesis, removal of prosthesis, spacer revision

(a) Inconsistent previous surgery: none, osteotomy

Table 5.17. Hip. Primary total replacement: type of implanted medical device

| | Total replacement (elective) | | Total replacement (emergency) | | TOTAL | |
|----------------------------------|---------------------------------|------|----------------------------------|-------|---------------|------|
| | N | % | N | % | N | % |
| Cup | 11,525 | | 1,086 | | 12,611 | |
| Cemented for primary procedure | 198 | 1.7 | 60 | 5.5 | 258 | 2.0 |
| metal | 196 | 99.0 | 60 | 100.0 | 256 | 99.2 |
| polyethylene | 1 | 0.5 | 0 | 0.0 | 1 | 0.4 |
| other | 1 | 0.5 | 0 | 0.0 | 1 | 0.4 |
| Cementless for primary procedure | 9,068 | 78.7 | 767 | 70.6 | 9,835 | 78.0 |
| metal | 8,318 | 91.7 | 694 | 90.5 | 9,012 | 91.6 |
| other | 750 | 8.3 | 73 | 9.5 | 823 | 8.4 |
| Cementless for revision | 2,259 | 19.6 | 259 | 23.8 | 2,518 | 20.0 |
| Insert | 11,525 | | 1,086 | | 12,611 | |
| Polyethylene | 8,220 | 71.3 | 925 | 85.2 | 9,145 | 3.3 |
| Ceramic | 3,290 | 28.5 | 157 | 14.5 | 3,447 | 96.7 |
| Metal | 15 | 0.1 | 4 | 0.4 | 19 | 0.6 |
| Head | 11,525 | | 1,086 | | 12,611 | |
| For partial replacement | 1,301 | 11.3 | 55 | 5.1 | 1,356 | 10.8 |
| ceramic | 1,151 | 88.5 | 42 | 76.4 | 1,193 | 88.0 |
| metal | 150 | 11.5 | 13 | 23.6 | 163 | 12.0 |
| For total replacement | 10,224 | 88.7 | 1,031 | 94.9 | 11,255 | 89.2 |
| ceramic | 9,514 | 93.1 | 875 | 84.9 | 10,389 | 92.3 |
| metal | 710 | 6.9 | 156 | 15.1 | 866 | 7.7 |
| Stem | 11,525 | | 1,086 | | 12,611 | |
| Cemented for primary procedure | 539 | 4.7 | 136 | 12.5 | 675 | 5.4 |
| non-modular straight | 496 | 92.0 | 126 | 92.6 | 622 | 92.1 |
| non-modular anatomical | 29 | 5.4 | 4 | 2.9 | 33 | 4.9 |
| modular | 9 | 1.7 | 6 | 4.4 | 15 | 2.2 |
| conservative | 5 | 0.9 | 0 | 0.0 | 5 | 0.7 |
| Cementless for primary procedure | 10,731 | 93.1 | 935 | 86.1 | 11,666 | 92.5 |
| non-modular straight | 7,296 | 68.0 | 699 | 74.8 | 7,995 | 68.5 |
| non-modular anatomical | 494 | 4.6 | 53 | 5.7 | 547 | 4.7 |
| modular | 867 | 8.1 | 151 | 16.1 | 1,018 | 8.7 |
| conservative | 2,074 | 19.3 | 32 | 3.4 | 2,106 | 18.1 |
| Cemented for revision | 0 | 0.0 | 1 | 0.1 | 1 | 0.0 |
| Cementless for revision | 254 | 2.2 | 13 | 1.2 | 267 | 2.1 |
| Large resections | 1 | 0.0 | 1 | 0.1 | 2 | 0.0 |

Table 5.18. Hip. Primary total replacement: bearing surface

| | Total replacement (elective) | | Total replacement (emergency) | | TOTAL | |
|--------------------------------------|---------------------------------|------|----------------------------------|------|---------------|------|
| | N | % | N | % | N | % |
| Bearing surface (head/insert) | 11,525 | | 1,086 | | 12,611 | |
| Ceramic-Polyethylene | 7,383 | 64.1 | 758 | 69.8 | 8,141 | 64.6 |
| Ceramic-Ceramic | 3,273 | 28.4 | 157 | 14.5 | 3,430 | 27.2 |
| Metal-Polyethylene | 837 | 7.3 | 167 | 15.4 | 1,004 | 8.0 |
| Metal-Ceramic | 17 | 0.1 | 0 | 0.0 | 17 | 0.1 |
| Ceramic-Metal | 9 | 0.1 | 2 | 0.2 | 11 | 0.1 |
| Metal-Metal | 6 | 0.1 | 2 | 0.2 | 8 | 0.1 |

Table 5.19. Knee. Primary total replacement: type of implanted medical device

| | Primary | |
|------------------------------|---------------|------|
| | N | % |
| Femoral component | 15,269 | |
| Bicondylar | 13,597 | 89.0 |
| cemented | 11,679 | 85.9 |
| cementless | 1,552 | 11.4 |
| cementable | 366 | 2.7 |
| Unicondylar | 1,480 | 9.7 |
| Revision | 192 | 1.3 |
| Tibial insert | 15,269 | |
| Bicondylar | 13,434 | 88.0 |
| mobile | 4,043 | 30.1 |
| fixed | 9,391 | 69.9 |
| Unicondylar | 1,444 | 9.5 |
| Revision | 391 | 2.6 |
| mobile | 93 | 23.8 |
| fixed | 298 | 76.2 |
| Tibial component | 15,269 | |
| Cemented bicondylar | 11,670 | 76.4 |
| mobile | 3,055 | 26.2 |
| fixed | 8,615 | 73.8 |
| Cementless bicondylar | 1,014 | 6.6 |
| mobile | 823 | 81.2 |
| fixed | 191 | 18.8 |
| Cementable mobile bicondylar | 302 | 2.0 |
| mobile | 266 | 88.1 |
| fixed | 36 | 11.9 |
| Unicondylar | 1,490 | 12.9 |
| Revision | 793 | 5.2 |
| mobile | 75 | 9.5 |
| fixed | 718 | 90.5 |

Table 5.20. Knee. Primary total replacement: implant of patella

| | N | % |
|----------------|---------------|------|
| Implant | 15,269 | |
| No patella | 13,679 | 89.6 |
| With patella | 1,590 | 10.4 |

Figure 4.1. Temporal trend of elective primary joint replacements in Italy




Figure 4.2. Hip. Incidence rates per 100,000 population by type of procedure




Figure 4.3. Knee. Incidence rates per 100,000 population by type of procedure




Figure 4.4. Shoulder. Incidence rates per 100,000 population by type of procedure




Figure 4.5. Hip. Primary total replacement: percentage of hospitals by region and volume of activity




Figure 4.6. Hip. Revision: percentage of hospitals by region and volume of activity




Figure 4.7. Hip. Elective primary total replacement: inter-regional mobility index (%), a) attraction index, b) escape index




Figure 4.8. Knee. Primary total replacement: percentage of hospitals by region and volume of activity




Figure 4.9. Knee. Revision: percentage of hospitals by region and volume of activity




Figure 4.10. Knee. Primary total replacement: inter-regional mobility index (%), a) attraction index, b) escape index




Figure 4.11. Shoulder. Primary total replacement: percentage of hospitals by region and volume of activity




Figure 4.12. Shoulder. Primary partial replacement: percentage of hospitals by region and volume of activity




Figure 4.13. Shoulder. Elective primary total replacement: inter-regional mobility index (%),
a) attraction index, b) escape index

