

Italian Arthroplasty Registry

Addendum to the Annual Report 2019

English version of Tables and Figures

©Istituto Superiore di Sanità First edition: October 2020

Excerpt from: Registro Italiano ArtroProtesi. Report Annuale 2019.

Il Pensiero Scientifico Editore

Via San Giovanni Valdarno 8, 00138 Roma

Telefono (+39) 06 86281 – Fax (+39) 06 86282250

pensiero@pensiero.it

www.pensiero.it – www.vapensiero.info

www.facebook.com/PensieroScientifico

Layout: Doppiosegno s.n.c. Roma

Editorial coordination: Martina Teodoli

Authors

Iuliia Urakcheeva Alessia Biondi and Marina Torre

Italian National Institute of Health Scientific secretariat of the Presidency RIPI working group Rome

Acknowledgements

Special thanks to the RIAP working group: Eugenio Carrani, Stefania Ceccarelli, Enrico Ciminello, Attanasio Cornacchia, Michela Franzò, Fabio Galati, Paola Laricchiuta, Mascia Masciocchi and Emanuela Saquella; and to the orthopaedic surgeons experts of the RIAP Steering committee: Filippo Boniforti, Stefano Lepore, Silvano Piffer, Emilio Romanini, Stefano Tornago and Gustavo Zanoli.

Please cite as follows:

Urakcheeva I, Biondi A and Torre M. Italian Arthroplasty Registry. Annual Report 2019 – Addendum. Roma: Il Pensiero Scientifico Editore, 2020.

Tables of contents

| RIAP: the m | nost important things to know | 8 |
|-------------|---|----|
| RIAP in 201 | 8-2019: progress and key achievements | 10 |
| | | |
| Inde | x of Tables and Figures | |
| Table 2.1. | Number of participating hospitals and coverage and number of collected procedure | es |
| | and completeness, by participating institution and by joint (year 2018) | 19 |
| Table 2.2. | RIAP completeness (years 2017 and 2018). Comparison with HDD data 2017 | |
| | and estimate 2018, by joint | 21 |
| Table 2.3. | Hip. Number of procedures analysed, by procedure type | 22 |
| Table 2.4. | Hip. Number of procedures by type of hospital, by procedure type | 23 |
| Table 2.5. | Hip. Number of procedures by patient gender and age group, by procedure type | 24 |
| Table 2.6. | Hip. Number of procedures by operated side, surgical approach and fixation, | |
| | by procedure type | 25 |
| Table 2.7. | Hip. Number of primary procedures by indication for surgery and previous procedure, by procedure type | 26 |
| Table 2.8. | Hip. Number of revision procedures by indication for surgery and previous procedure | 27 |
| Table 2.9. | Hip. Number of procedures analysed, by device | 28 |
| Table 2.10. | Hip. Number of procedures by type of fixation, by procedure type | 29 |
| Table 2.11. | Hip. Number of total replacement procedures by type of bearing, | |
| | by procedure type | 30 |
| Table 2.12. | Hip. Number of revision procedures by type of bearing | 31 |
| Table 2.13. | Hip. Number of total replacement procedures by stem type, by procedure type | 32 |
| Table 2.14. | Knee. Number of procedures analysed, by procedure type | 33 |
| Table 2.15. | Knee. Number of procedures by type of hospital, by procedure type | 34 |
| Table 2.16. | Knee. Number of procedures by patient gender and age group, | |
| | by procedure type | 35 |
| | | |

| Table 2.17. | Knee. Number of procedures by operated side and surgical approach, by procedure type | 36 |
|-------------|--|----|
| Table 2.18. | Knee. Number of primary procedures, by indication for surgery and previous procedure | 37 |
| Table 2.19. | Knee. Number of revision procedures, by indication for surgery and previous procedure | 38 |
| Table 2.20. | Knee. Number of procedures analysed for the implanted device, by procedure type | 39 |
| Table 2.21. | Knee. Number of procedures by type of fixation, by procedure type | 40 |
| Table 2.22. | Knee. Number of primary procedures by type of tibial tray | 42 |
| Table 2.23. | Shoulder. Number of procedures analysed, by procedure type | 43 |
| Table 2.24. | Shoulder. Number of procedures by prosthesis type implanted in the total replacement | 44 |
| Table 2.25. | Shoulder. Number of procedures by type of hospital, by procedure type | 45 |
| Table 2.26. | Shoulder. Number of procedures by patient gender and age group, by procedure type | 46 |
| Table 2.27. | Shoulder. Number of procedures by operated side and surgical approach, by procedure type | 47 |
| Table 2.28. | Shoulder. Number of primary procedures by indication for surgery and previous procedure | 48 |
| Table 2.29. | Shoulder. Number of revision procedures by indication for surgery and previous procedure | 49 |
| Table 2.30. | Shoulder. Number of procedures by type of fixation, by procedure type | 50 |
| Figure 1.1. | RIPI organisation | 16 |
| Figure 1.2. | Participation in the RIAP | 17 |
| Figure 1.3. | Flow diagram of the RIAP data collection model | 18 |
| Figure 2.1. | Flowchart of the RIAP data quality control process: procedures | 51 |
| Figure 2.2. | Flowchart of the RIAP data quality control process: implanted medical devices | 52 |
| Figure 2.3. | Hip. Types of bearing. Total replacement (elective procedures) | 53 |
| Figure 2.4. | Hip. Types of bearing. Total replacement (emergency) | 54 |
| | | |

Appendix 2A. Joint replacements in Italy. Hospital Discharge Data analysis 2016-2017

| Table 1. | Joint replacements in Italy, per year of discharge. 2001-2017 and estimate 2018 | 56 |
|-----------|---|----|
| Table 2. | Hip. Number of procedures by region of admission and procedure type. 2016-2017 | 58 |
| Table 3. | Hip. Primary total replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017 | 60 |
| Table 4. | Hip. Revision. Number of hospitals by region of admission and class of volume. 2016 and 2017 | 62 |
| Table 5. | Hip. Percent distribution of hospital discharges by patient gender and age group, and by procedure type. 2016 and 2017 | 63 |
| Table 6. | Hip. Percent distribution of hospital discharges by mode of discharge and by procedure type. 2016 and 2017 | 64 |
| Table 7. | Knee. Number of procedures by region of admission and procedure type. 2016-2017 | 66 |
| Table 8. | Knee. Primary total replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017 | 68 |
| Table 9. | Knee. Revision. Number of hospitals by region of admission and class of volume. 2016 and 2017 | 70 |
| Table 10. | Knee. Percent distribution of hospital discharges by patient gender and age group, and by procedure type. 2016 and 2017 | 71 |
| Table 11. | Knee. Percent distribution of hospital discharges by mode of discharge and by procedure type. 2016 and 2017 | 72 |
| Table 12. | Shoulder. Number of procedures by region of admission and procedure type. 2016-2017 | 74 |
| Table 13. | Shoulder. Total replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017 | 76 |
| Table 14. | Shoulder. Partial replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017 | 78 |
| Table 15. | Shoulder. Percent distribution of hospital discharges by patient gender and age group, and by procedure type. 2016 and 2017 | 80 |
| | | |

| lable 16. | and by procedure type. 2016 and 2017 | 81 |
|-----------|---|----|
| Table 17. | Ankle. Total replacement. Number of procedures by region of admission and procedure type. 2016-2017 | 83 |
| Table 18. | Ankle. Total replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017 | 84 |
| Table 19. | Ankle. Total replacement. Percent distribution of hospital discharges by patient gender and age group, and by procedure type. 2016 and 2017 | 86 |
| Table 20. | Ankle. Total replacement. Percent distribution of hospital discharges by mode of discharge, by procedure type. 2016 and 2017 | 87 |
| Figure 1. | Hip. Elective total replacement. Inter-regional mobility (attraction and escape indices) in 2016 (a) and in 2017 (b) | 65 |
| Figure 2. | Knee. Total replacement. Inter-regional mobility (attraction and escape indices) in 2016 (a) and in 2017 (b) | 73 |
| Figure 3. | Shoulder. Elective total replacement. Inter-regional mobility (attraction and escape indices) in 2016 (a) and in 2017 (b) | 82 |

This summary provides a brief overview of the main findings described in the 2019 RIAP Annual Report, as by the end of 2019.

It contains all the tables and figures, while the entire Report is available only in Italian (you can find it here). The technical Appendix includes an analysis of the most recent national Hospital Discharge Data available (years 2016 and 2017).

RIAP: the most important things to know

What is RIAP?

The Italian Arthroplasty Registry, RIAP, was started in 2006 within the framework of a collaboration between the Italian Ministry of Health, DG for Medical Devices and Pharmaceutical Services (DGDMF) and the Italian National Institute of Health (Istituto Superiore di Sanità, ISS), to set up a data flow to monitor the joint prostheses performance.

What are the goals?

Main aims of the RIAP are to monitor the longterm effectiveness of hip, knee, shoulder and ankle prostheses (measured as implant survival), and support Regions and hospitals when recall of patients is needed because of problems reported on specific implants.

What information is collected?

The information collected includes Hospital Discharge Data (HDD) integrated with an additional Minimum Data Set (MDS) - specific for each joint - about the procedure, the operated body side and the data needed to identify the implanted device. In 2018, 70,584 procedures were collected (+4.7% more than a year before), representing about 33,3% of the national volume.

Personal data are treated by RIAP in compliance with the current European legislation on privacy (EU Regulation 2016/679). Clinical, health and demographic data are treated applying criteria ensuring the highest confidentiality, in compliance with the security regulations for digital and paper-based archives.

How is RIAP organised?

RIAP is a federation of regional registries coordinated by the ISS. Through a web interface, the surgeons collect the MDS, which is successively linked to the HDD by the regional coordinating center. The latter is responsible for the transmission of the linked data to the ISS. Currently, participation of Regions and structures is on a voluntary basis.

A distinctive feature of the RIAP model is its transferability into areas beyond arthroplasty, especially where the replacement interventions are mainly carried out in structures that can provide the Hospital Discharge Data (Schede di dimissione ospedaliera, SDO).

Why a registry, and why a national registry?

Joint prosthesis is a recognised solution for the treatment of disabling joint diseases. In Italy, like in many countries, arthroplasties are constantly growing. A national registry allows assessing the outcomes of primary and revision procedures, based on patient's specificity, and intervention and implanted device characteristics. If needed, there must be the possibility to recall all the implanted patients, including the patients that were operated in a different region from the one they live in. This is the reason why the Registry must have national coverage.

How is the implanted device identified and characterised?

The RIAP-DM Dictionary, a database containing data supplied by manufacturers useful to identify each implanted device, is available for the operators.

To identify each implanted device, RIAP made the RIAP-DM Dictionary available to operators. It is a database containing data supplied by manufacturers, including single-device specific data. The information contained in the RIAP-DM Dictionary is checked and matched against the information from the National Database of medical devices of the Ministry of Health.

RIAP is now developing a collaboration with the National Joint Registry for England, Wales, Northern Ireland and the Isle of Man, to utilise its component database and classification, recently upgraded in cooperation with the Endoprothesen Register Deutschland (EPRD) to include in the RIAP-DM Dictionary the specs of the implanted device.

Here you can consult the Report summaries (Addendum) of years 2014-2018.

RIAP in 2018-2019: progress and key achievements

Highlights

- Starting from February 2019, the RIAP is part of the Italian Implantable Prostheses Registry (RIPI) and represents the model for the other registries considered by the RIPI.
- Capitalising the experience gained with the RIAP, we have been supporting the Italian Ministry of Health in the development of a pilot platform for the National Registry of Breast Implants. We also worked on drafting the regulation prescribed by the law that will render mandatory for the Italian Regions to feed the registries.
- We implemented a new module in our IT platform RaDaR to allow the collection of additional data regarding ankle arthroplasty. We also kept updating the RIAP-DM Dictionary, our database of implantable devices, increasing the number of product codes (+1000) and contributors. The Quality Control procedures have been further improved.
- In 2019 we exchanged knowledge

- with the English National Joint Registry (NJR). This collaboration has allowed us to support the Ministry of Health in updating the national Classification of Medical Devices (CND) and in its transition to the European Medical Device Nomenclature.
- Three new topics resulting from the RIAP analytical work in 2019 are reproduced in the Report: first survival analysis in AP of Bolzano, start of retrospective data gathering in AP Trento, and analysis on national HDD 2016-2017.

2019 has been a year of ongoing activities, but also of new achievements and improved procedures that allowed us to provide surgeons and patients with better information to support decision-making on prostheses. Our team kept growing and was enforced by PhD students and researchers.

Hereby is a summary of the main activities and achievements of 2019:

 The drafting of the regulation required by the decree (DPCM - Decreto del Presidente del Consiglio) has begun, the last piece of the regulatory framework for the registries which, with the law n.145/2018, made it mandatory for the Regions to feed them.

- A special commitment has been dedicated to the drafting of the launch of the Italian Implantable Prosthesis Registry (Registro Italiano delle Protesi Impiantabili. RIPI) in order to make it operational in accordance with the law. The RIPI has been launched in February 2019 (see ripi.iss.it). It is a coordinating structure of registries of implantable devices divided by subject: Italian Spinal Implants Registry (RIDIS), Italian Implantable Cardioverter-defibrillator and Pacemaker Registry (RIDEP), Italian Heart Valves Registry (RIVAC) and the RIAP which is its most developed component and a model (Figure 1.1).
- Thanks to the feedback provided by the participants, the quality of data gathered in the regions has been growing: 96.1% of cases passed QC on interventions this year, vs 91% a year before.
- The process of the IT infrastructure reengineering started in order to include and automatise the quality control processes and provide faster feedback to the participants

In February 2019, we made the new module for Ankle arthroplasty data collection avail-

able to registered operators on the IT platform RaDaR

The RIAP-DM Dictionary has been updated, adding 1000 product codes by the end of the year (over 66,000) relating to 101 manufacturers, sent by 39 companies.

- We took part in the works of the International Society of Arthroplasty registries (ISAR). We have developed communications with the Network of Orthopedic Registries of Europe (NORE) and with the English National Joint Registry. It is our intention to sign a collaboration agreement with the NJR by 2020. The aim is to access, consult and possibly feed the Component Library NJR-EPRD (the database of devices shared between the English NJR register and the German EPRD registry).
- We further optimised the RIAP communication assets - the annual report and the website, and launched the website of the RIPI, in Italian and English.
- Through the application of the new function of patients pseudonymisation the Autonomous Province of Bolzano provided the RIAP with its complete 10 years historical data. At the same time, the Autonomous Province of Trento has

launched a retrospective recovery process of missing information (mainly on hip procedures) from 2010 to 2018 using our web application RaDaR2. This result is a first step to produce implant survival analysis, made possible thanks to the proactive approach of the regional RIAP participants.

To provide the reader with figures about the national number of arthroplasties performed in Italy, this edition of the annual Report contains detailed analysis of the 2016 and 2017 HDD data, the last available to the Institute (see Appendix 2A). HDD data represent an important reference to perform statistical and epidemiological analysis in public health and to calculate the actual RIAP completeness at both national and regional levels.

To do this, HDD was browsed using the ICD9-CM procedure codes of interest for RIAP (Hip, Knee, Shoulder and Ankle primary and revision arthroplasties) listed in Table 1. All the procedures performed during a hospitalisation were included. Hip and shoulder arthroplasties were further classified on the basis of the associated diagnosis; emergency procedures are those performed to treat a fracture (ICD9-CM codes 820.XX and 812.XX). Inter-regional mobility for every joint was measured using the

attraction and escape indices computed considering only the ICD9-CM code registered in the so-called "principal procedure". Appendix 2A shows the temporal trend of the number of procedures over 17 years of observation (2001-2017) and, for each joint and for the years 2016 and 2017, the volumes of activity broken down by region, the characteristics of the operated patients (sex, age), the modality of discharge and the interregional mobility (the latter not for ankle).

RELATED ACTIVITIES AND COLLABORATIONS

In 2019, the RIAP team actively collaborated with the Italian Ministry of Health. Capitalising on the Registry's experience, we provided support to the Ministry in the development of the pilot platform of the National Registry of Breast Implants.

Moreover, the RIAP team collaborated with the ministerial Healthcare Technical Committee for updating the Italian CND classification, taken as the basic nomenclature of reference in the pan-european medical devices database EU-DAMED.

In June 2019, we presented a first study on harmonisation of CND and NJR classifications, to create a shared international database of the orthopaedic prostheses. Both CND and NJR taxonomies were matched and compared for hip joint components. A combination of attributes in the NJR taxonomy was found for each CND terminal level. This is the first step towards the organisation of an international component database since their comparability enables the integration of devices from different databases

Key findings from the annual report 2019

In 2019, RIAP collected and admitted to quality control the data on **70,584** procedures performed in 2018 in eight regions (Lombardy, Tuscany, Marche, Apulia, Basilicata, Calabria, Sicily, and Campania), two Autonomous Provinces (Bolzano, Trento) and two hospital structures ("Policlinico Città di Alessandria" of Alessandria and "Santa Maria della Misericordia" of Udine) (Table 2.1, Figure 1.2).

After the quality control, 96.1% and 92.6% of the received records, respectively, were admitted to the interventions analysis and to the devices analysis. Both percentage figures are higher than in 2018. (Figure 2.1, Figure 2.2).

The overall 2018 data completeness of RIAP was 64.7% (Table 2.1). Compared to 2017, a slight reduction in completeness was recorded for hip (- 0.1%) and knee (- 0.8%) and a bigger drop for shoulder (- 40.7%) resulting in

general reduction in completeness of - 0.9%. As in previous years, in the regions where no provision has been adopted that effectively makes data recording mandatory, we observe low values of data completeness and coverage. For the shoulder, another possible cause of downtrend is the start of data collection in Marche and Sicily where only some structures have given their contribution, and therefore, the number of procedures recorded was significantly lower than that of the interventions carried out globally in the region.

Overall, in 2018 the estimated number of replacements performed in Italy was around 212,205 of which RIAP data represented 33.3% (estimate) vs 33.2% of 2017 (Table 2.2).

Hip: Of 38,734 interventions collected, 37,709 passed the QC for interventions analysis. Total hip replacements represent 72.5% of analysed cases, partial hip replacements - 21.9%, and revisions - 5.6%. 36,353 passed the QC for devices analysis (fixation type, materials of bearing surfaces, type of stem). (Tables 2.3 - 2.13, Figures 2.3, 2.4).

Knee: Of 29,918 interventions collected, 29,198 passed the QC for interventions analysis. 83.6% of primary interventions recorded were total, while 16.4% were unicondylar

knee replacements. 28,497 (94% of interventions received) passed the QC for devices analysis (fixation type, tibial tray type). (Tables 2.14 - 2.22).

Shoulder: Of 1038 interventions collected, 939 passed the QC for interventions analysis. 56.7% of the interventions were total replacements, 3.5% were partial replacements and 2% were revisions. 39.8% of primary interventions were not specified (those are 366 interventions from the Register of Orthopaedic Prosthetic Implantology of the Apulia Region). 481 interventions (45.8% of interventions received) passed the QC for devices analysis. (See Tables 2.23 - 2.30).

Challenges

The first goal for the upcoming years is to make RIAP a nationwide and 100% complete registry. This means to achieve the enrollment of all the 21 Italian Regions and full collection of required data. A challenging task, especially considering that in the Italian National Health Service, the Regions are autonomous in regards to health services planning and organisation. Achieving this goal highly depends on the strategies adopted at local level. In the RIAP network, the data completeness and quality varies across Regions even if the law (L145/2018) made it mandatory to feed the

national Registry. Highest levels of completeness are registered where regional data collection was made compulsory. Therefore, it is of high importance that rules rendering data collection indispensable are introduced at regional level, i.e. by linking the reimbursement of interventions costs to the registration of the performed procedure. In the near future, the Regulation required by law will be finalised, defining the detailed procedures of feeding the RIAP and other national Registries.

As for the quality improvement of regional data, it remains crucial that the regional coordination centres will continue to communicate the results of the QC to each structure, in order to monitor compilation errors and improve the quality of collected data. The RIAP team will continue to work on the automatised QC to provide the regional data operators with immediate feedback.

Future development and plans for 2020-2022

The activities of the RIAP will now continue within **the operational framework of the RIPI**, of which the RIAP development constitutes an important research line. In 2019, we have taken the first concrete steps to structure RIPI, organising a first working group for the Italian Spinal Implants Registry (RIDIS) and

collaborating with the Italian Association of Arrhythmology and Cardiostimulation (AIAC) to design the Italian Implantable Cardioverterdefibrillator and Pacemaker Registry (RIDEP).

In 2020-2021, the consolidated experience of the RIAP will be extended to design the registries of spinal devices, pacemakers and defibrillators, and heart valves. Support will be further provided to the Ministry of Health to launch the National Registry of Breast Implants. The goal is to use the RIAP as a benchmark for a national system of registries that, soon, might be integrated into the current flows of the National Health Service. Re-engineering the IT infrastructure and designing the RIPI platform could become the first step of this process.

With the entry into force of the new European regulation EU 2017/745, the classification and traceability of medical devices become critically important. The implantable devices should be included in the European database on medical devices (EUDAMED) by 26 May, 2021. In this situation, it is our aim to further collaborate

with the Italian Ministry of Health to **contribute to updating the national classification CND for joint prostheses**. With the same aim, in order to keep the RIAP-DM Dictionary constantly updated and to start the design of the dictionaries of the other registries, we will intensify the collaboration with the industry through Confindustria-DM.

We will keep strengthening the international collaborations, contributing to the work of the International Society of Arthroplasty Registry (ISAR), the Network of Orthopaedic Registries in Europe (NORE) and participating in international technical meetings and calls of European projects (Horizon). In 2020 we should formalise the collaboration with the NJR for access to the NJR / EPRD database

We will continue **raising stakeholder aware- ness** about the importance of recording the replacement surgeries. Moreover, we plan to increase our focus on the needs of our target audiences in 2020-2021, by developing a scientific research regarding the communication strategy of the RIPI.

Figure 1.1. RIPI organisation

Italian Arthroplasty Registry Italian Spinal Implants Registry Italian Implantable CardioverterDefibrillator and Pacemaker Registry Registry Italian Heart Valves Registry Registry

Figure 1.2. Participation in the RIAP




Figure 1.3. Flow diagram of the RIAP data collection model




Table 2.1. Number of participating hospitals and *coverage* and number of collected procedures and *completeness*, by participating institution and by joint (year 2018)

| Participating authority | Joint | Participating authority | Coverage (*) | Collected procedures | Completeness (**) |
|-------------------------|----------|-------------------------|---|----------------------|----------------------|
| Regions | | N | % | N | % |
| Lombardy | Hip | 105 | 100.0 | 23,903 | 97.7 |
| | Knee | 107 | 100.0 | 19,115 | 97.5 |
| AP Bolzano | Hip | 12 | 100.0 | 1,341 | 96.0 |
| | Knee | 12 | 100.0 | 902 | 95.0 |
| | Shoulder | 4 | 44.4 | 23 | 31.1 |
| AP Trento | Hip | 8 | 100.0 | 1,358 | 97.6 |
| | Knee | 8 | 100.0 | 772 | 97.0 |
| Tuscany | Hip | 1 | 2.3 | 476 | 5.0 |
| | Knee | 1 | 2.2 | 257 | 3.0 |
| Marche | Hip | 11 | 61.1 | 767 | 31.1 |
| | Knee | 10 | 55.6 | 895 | 45.0 |
| | Shoulder | 6 | 55.6 895 33.3 23 84.4 4,901 87.3 3,199 | 8.9 | |
| Campania | Hip | 65 | 84.4 | 4,901 | 75.7 |
| | Knee | 62 | 87.3 | 3,199 | 78.4 |
| | Shoulder | 34 | 75.6 | 515 | 79.8 |
| Apulia | Hip | 45 | 100.0 | 4,782 | 99.8 |
| | Knee | 42 | 100.0 | 3,669 | 99.9 |
| | Shoulder | 34 | 100.0 | 471 | 99.8 |
| Basilicata | Hip | 1 | 16.7 | 126 | 21.1 |
| | Knee | 1 | 16.7 | 74 | 26.6 |
| Calabria | Hip | 16 | 76.2 | 885 | 46.1 |
| | Knee | 14 | 70.0 | 468 | 27.7 |
| Sicily | Hip | 4 | 5.5 | 195 | 3.3 |
| | Knee | 4 | 5.8 | 567 | 10.4 |
| | Shoulder | 1 | 1.9 | 6 | 1.1 |
| Subtotal (regions) | Hip | 268 | 65.5 | 38,734 | 65.6 |
| | Knee | 261 | 65.6 | 29,918 | 63.6 |
| | Shoulder | 79 | 49.7 | 1,038 | 51.7 |

(continued)

Table 2.1. (continued)

| Participating authority | Joint | Participating authority | Coverage (*) | Collected procedures | Completeness (**) |
|--|----------|-------------------------|--------------|----------------------|----------------------|
| Single hospitals | | N | % | N | % |
| Città di Alessandria Hospital (Alessandria) | Hip | 1 | - | 321 | 77.9 |
| | Knee | 1 | - | 261 | 62.9 |
| Santa Maria della Misericordia Hospital (Udine) | Hip | 1 | - | 161 | 100.0 |
| | Knee | 1 | - | 139 | 100.0 |
| | Shoulder | 1 | - | 12 | 100.0 |
| Subtotal (single hospitals) | Hip | 2 | - | 482 | 84.1 |
| | Knee | 2 | - | 400 | 72.2 |
| | Shoulder | 1 | - | 12 | 100.0 |
| Total number of procedures admitted to quality control | | N | % | N | % |
| | Hip | 270 | - | 39,216 | 65.8 |
| otal number of procedures admitted | Knee | 263 | - | 30,318 | 63.7 |
| | Shoulder | 80 | - | 1,050 | 52.0 |
| | ALL | 273 | | 70,584 | 64.7 |

^(*) Coverage: ratio between hospitals participating in RIAP and hospitals performing arthroplasties, based on data from HDD

^(**) Completeness: ratio between procedures collected by RIAP and procedures performed by all hospitals in the Region, based on data from HDD

Table 2.2. RIAP completeness (years 2017 and 2018). Comparison with HDD data 2017 and estimate 2018, by joint

| | | 2017 | | | | | |
|----------|---------|--------|--------------|-------------------|--------|----------------------------|--|
| | HDD | RIAP | Completeness | HDD (estimate) | RIAP | Completeness (estimate) | |
| | N | N | % | N | N | % | |
| Joint | 202,747 | 67,366 | 33.2 | 212,205 | 70,584 | 33.3 | |
| Hip | 112,375 | 38,460 | 34.2 | 115,308 | 39,216 | 34.0 | |
| Knee | 81,271 | 28,023 | 34.5 | 86,735 | 30,318 | 35.0 | |
| Shoulder | 9,101 | 883 | 9.7 | 10,162 | 1,050 | 10.3 | |

Table 2.3. Hip. Number of procedures analysed, by procedure type

| | N | % |
|----------------------------|--------|------|
| Procedure type | 37,709 | |
| Total replacement | 27,329 | 72.5 |
| - elective | 24,049 | 88.0 |
| - emergency | 3,280 | 12.0 |
| Partial replacement | 8,269 | 21.9 |
| Revision | 2,111 | 5.6 |
| Partial revision (*) | 1,459 | 69.1 |
| Total revision | 230 | 10.9 |
| Removal of prosthesis (**) | 422 | 20.0 |

^(*) Includes: conversion from partial to total prosthesis

^(**) Includes: removal of prosthesis, removal with spacer implant, spacer revision

Table 2.4. Hip. Number of procedures by type of hospital, by procedure type

| | Tot | al repla | cement | | Partial | | Revision (*) | | TOTAL | |
|-----------------------------------|----------|----------|-----------|------|-------------|------|--------------|------|--------|------|
| | elective | | emergency | | replacement | | | | | |
| | N | % | N | % | N | % | N | % | N | % |
| Type of hospital | 24,049 | | 3,280 | | 8,269 | | 2,111 | | 37,709 | |
| Public hospitals | 10,064 | 41.8 | 2,779 | 84.7 | 6,884 | 83.3 | 1,091 | 51.7 | 20,818 | 55.2 |
| Private hospitals, accredited | 12,744 | 53.0 | 477 | 14.5 | 1,274 | 15.4 | 946 | 44.8 | 15,441 | 40.9 |
| Private hospitals, not accredited | 1,241 | 5.2 | 24 | 0.7 | 111 | 1.3 | 74 | 3.5 | 1,450 | 3.8 |

^(*) Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision

Table 2.5. Hip. Number of procedures by patient gender and age group, by procedure type

| | | | acement | | | placement | | | TOTAL | |
|---------------------|--------|------|---------|------|-------|-----------|-------|------|--------|------|
| | electi | ive | emerge | ency | | | | | | |
| | N | % | N | % | N | % | N | % | N | % |
| Gender | 24,049 | | 3,280 | | 8,269 | | 2,111 | | 37,709 | |
| Male | 11,198 | 46.6 | 915 | 27.9 | 2,162 | 26.1 | 836 | 39.6 | 15,111 | 40.1 |
| Female | 12,851 | 53.4 | 2,365 | 72.1 | 6,107 | 73.9 | 1,275 | 60.4 | 22,598 | 59.9 |
| Age group by gender | | | | | | | | | | |
| Male | 11,198 | | 915 | | 2,162 | | 836 | | 15,111 | |
| Mean age | 65 | | 71 | | 83 | | 69 | | 68 | |
| Standard deviation | 12 | | 12 | | 8 | | 13 | | 13 | |
| <45 | 634 | 5.7 | 16 | 1.7 | 5 | 0.2 | 35 | 4.2 | 690 | 4.6 |
| 45 - 54 | 1,693 | 15.1 | 103 | 11.3 | 12 | 0.6 | 92 | 11.0 | 1,900 | 12.6 |
| 55 - 64 | 2,770 | 24.7 | 128 | 14.0 | 50 | 2.3 | 155 | 18.5 | 3,103 | 20.5 |
| 65 - 74 | 3,541 | 31.6 | 252 | 27.5 | 175 | 8.1 | 252 | 30.1 | 4,220 | 27.9 |
| 75 - 84 | 2,285 | 20.4 | 289 | 31.6 | 858 | 39.7 | 232 | 27.8 | 3,664 | 24.2 |
| ≥ 85 | 275 | 2.5 | 127 | 13.9 | 1,062 | 49.1 | 70 | 8.4 | 1,534 | 10.2 |
| Female | 12,851 | | 2,365 | | 6,107 | | 1,275 | | 22,598 | |
| Mean age | 70 | | 74 | | 84 | | 74 | | 74 | |
| Standard deviation | 11 | | 10 | | 7 | | 11 | | 12 | |
| <45 | 309 | 2.4 | 14 | 0.6 | 4 | 0.1 | 14 | 1.1 | 341 | 1.5 |
| 45 - 54 | 935 | 7.3 | 81 | 3.4 | 16 | 0.3 | 59 | 4.6 | 1,091 | 4.8 |
| 55 - 64 | 2,382 | 18.5 | 286 | 12.1 | 56 | 0.9 | 163 | 12.8 | 2,887 | 12.8 |
| 65 - 74 | 4,403 | 34.3 | 810 | 34.2 | 390 | 6.4 | 361 | 28.3 | 5,964 | 26.4 |
| 75 - 84 | 4,161 | 32.4 | 834 | 35.3 | 2,445 | 40.0 | 481 | 37.7 | 7,921 | 35.1 |
| ≥ 85 | 661 | 5.1 | 340 | 14.4 | 3,196 | 52.3 | 197 | 15.5 | 4,394 | 19.4 |

 $^{(\}star) \, \text{Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision}$

Table 2.6. Hip. Number of procedures by operated side, surgical approach and fixation, by procedure type

| | T | otal rep | lacement | | Parti | | Revisio | on (*) | TOTA | ۱L |
|-------------------|--------|----------|----------|------|-------------|------|---------|--------|--------|------|
| | electi | ve | emerge | ency | replacement | | | | | |
| | N | % | N | % | N | % | N | % | N | % |
| Operated side | 24,049 | | 3,280 | | 8,269 | | 2,111 | | 37,709 | |
| Right | 12,864 | 53.5 | 1,671 | 50.9 | 4,179 | 50.5 | 1,060 | 50.2 | 19,774 | 52.4 |
| Left | 10,440 | 43.4 | 1,599 | 48.8 | 4,061 | 49.1 | 1,038 | 49.2 | 17,138 | 45.4 |
| Bilateral | 745 | 3.1 | 10 | 0.3 | 29 | 0.4 | 13 | 0.6 | 797 | 2.1 |
| Surgical approach | 24,049 | | 3,280 | | 8,269 | | 2,111 | | 37,709 | |
| Anterior | 4,661 | 19.4 | 210 | 6.4 | 448 | 5.4 | 122 | 5.8 | 5,441 | 14.4 |
| Anterolateral | 2,413 | 10.0 | 622 | 19.0 | 1,736 | 21.0 | 259 | 12.3 | 5,030 | 13.3 |
| Lateral | 4,037 | 16.8 | 898 | 27.4 | 2,757 | 33.3 | 513 | 24.3 | 8,205 | 21.8 |
| Posterolateral | 12,479 | 51.9 | 1,531 | 46.7 | 3,302 | 39.9 | 1,198 | 56.8 | 18,510 | 49.1 |
| Other | 459 | 1.9 | 19 | 0.6 | 26 | 0.3 | 19 | 0.9 | 523 | 1.4 |

^(*) Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision

Table 2.7. Hip. Number of primary procedures by indication for surgery and previous procedure, by procedure type

| | To | Total replacement | | | | | TOTA | ۱L |
|---------------------------------------|--------|-------------------|-------|-------|----------|------|--------|------|
| | electi | ve | emerg | jency | replacer | nent | | |
| | N | % | N | % | N | % | N | % |
| Indication for surgery | 24,049 | | 3,280 | | 8,269 | | 35,598 | |
| Primary osteoarthritis | 21,615 | 89.9 | 0 | 0.0 | 134 | 1.6 | 21,749 | 61.1 |
| Post-traumatic osteoarthritis | 343 | 1.4 | 0 | 0.0 | 33 | 0.4 | 376 | 1.1 |
| Rheumatoid arthritis | 58 | 0.2 | 0 | 0.0 | 0 | 0.0 | 58 | 0.2 |
| Neoplasia | 36 | 0.1 | 0 | 0.0 | 30 | 0.4 | 66 | 0.2 |
| Aseptic necroisis of femoral head | 952 | 4.0 | 0 | 0.0 | 15 | 0.2 | 967 | 2.7 |
| Congenital dislocation/hip dysplasia | 590 | 2.5 | 0 | 0.0 | 4 | 0.0 | 594 | 1.7 |
| Perthes disease or epiphysiolysis | 62 | 0.3 | 0 | 0.0 | 9 | 0.1 | 71 | 0.2 |
| Fractured neck of femur | 0 | 0.0 | 3,280 | 100.0 | 7,968 | 96.4 | 11,248 | 31.6 |
| Septic coxitis | 5 | 0.0 | 0 | 0.0 | 0 | 0.0 | 5 | 0.0 |
| Pseudoarthrosis neck fracture related | 28 | 0.1 | 0 | 0.0 | 12 | 0.1 | 40 | 0.1 |
| Other | 360 | 1.5 | 0 | 0.0 | 64 | 0.8 | 424 | 1.2 |
| Previous procedure | 24,049 | | 3,280 | | 8,269 | | 35,598 | |
| None | 21,990 | 91.4 | 3,098 | 94.5 | 7,613 | 92.1 | 32,701 | 91.9 |
| Osteosynthesis | 303 | 1.3 | 60 | 1.8 | 57 | 0.7 | 420 | 1.2 |
| Osteotomy | 91 | 0.4 | 0 | 0.0 | 4 | 0.0 | 95 | 0.3 |
| Arthrodesis | 4 | 0.0 | 0 | 0.0 | 2 | 0.0 | 6 | 0.0 |
| Other | 1,661 | 6.9 | 122 | 3.7 | 593 | 7.2 | 2,376 | 6.7 |

Table 2.8. Hip. Number of revision procedures by indication for surgery and previous procedure

| | Revision (*) | |
|--|--------------|------|
| | N | % |
| Indication for surgery | 2,111 | |
| Pain | 118 | 5.6 |
| Lysis | 57 | 2.7 |
| Wear | 174 | 8.2 |
| Implant fracture | 73 | 3.5 |
| Prosthesis dislocation | 293 | 13.9 |
| Periprosthetic fracture | 260 | 12.3 |
| Infection | 185 | 8.8 |
| Previous prosthesis removal | 34 | 1.6 |
| Aseptic loosening, cup | 382 | 18.1 |
| Aseptic loosening, stem | 228 | 10.8 |
| Aseptic loosening, total | 148 | 7.0 |
| Disease progression | 2 | 0.1 |
| Other | 157 | 7.4 |
| Previous procedure | 2,111 | |
| Total hip replacement | 1,607 | 76.1 |
| Revision of hip replacement | 80 | 3.8 |
| Spacer implantation or prosthesis removal (**) | 182 | 8.6 |
| Partial hip replacement | 172 | 8.1 |
| Other | 70 | 3.3 |

^(*) Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision

^(**) Includes: removal of prosthesis, removal with spacer implant, spacer revision

Table 2.9. Hip. Number of procedures analysed, by device

| | N | % |
|----------------------------|--------|------|
| Procedure type | 36,353 | |
| Total replacement | 26,905 | 74.0 |
| - elective | 23,726 | 88.2 |
| - emergency | 3,179 | 11.8 |
| Partial replacement | 7,366 | 20.3 |
| Revision | 2,082 | 5.7 |
| Partial revision (*) | 1,456 | 69.9 |
| Total revision | 205 | 9.8 |
| Removal of prosthesis (**) | 421 | 20.2 |

^(*) Includes: conversion from partial to total prosthesis

^(**) Includes: removal of prosthesis, removal with spacer implant, spacer revision

Table 2.10. Hip. Number of procedures by type of fixation, by procedure type

| | T | otal rep | olacement | | | Partial | | Revision (*) | | TOTAL | |
|---|--------|----------|-----------|------|-------------|---------|-------|--------------|--------|-------|--|
| | electi | ve | emerge | ncy | replacement | | | | | | |
| | N | % | N | % | N | % | N | % | N | % | |
| Fixation | 23,726 | | 3,179 | | 7,366 | | 2,082 | | 36,353 | | |
| Cemented (stem + cup) | 1,120 | 4.7 | 196 | 6.2 | 0 | 0.0 | 92 | 4.4 | 1,408 | 3.9 | |
| Reverse hybrid (uncemented stem and cemented cup) | 301 | 1.3 | 141 | 4.4 | 0 | 0.0 | 123 | 5.9 | 565 | 1.6 | |
| Only cemented cup | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 71 | 3.4 | 71 | 0.2 | |
| Hybrid (cemented stem and uncementled cup) | 1,113 | 4.7 | 202 | 6.4 | 0 | 0.0 | 68 | 3.3 | 1,383 | 3.8 | |
| Uncemented (stem + cup) | 21,192 | 89.3 | 2,640 | 83.0 | 0 | 0.0 | 1,254 | 60.2 | 25,086 | 69.0 | |
| Only uncemented cup | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 169 | 8.1 | 169 | 0.5 | |
| Only cemented stem | 0 | 0.0 | 0 | 0.0 | 3,091 | 42.0 | 38 | 1.8 | 3,129 | 8.6 | |
| Only uncemented stem | 0 | 0.0 | 0 | 0.0 | 4,275 | 58.0 | 182 | 8.7 | 4,457 | 12.3 | |
| Fixation declared "not applicable" for cup and stem even if the component is present | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 85 | 4.1 | 85 | 0.2 | |

^(*) Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision

Table 2.11. Hip. Number of total replacement procedures by type of bearing, by procedure type

| | | Total rep | TOTAL | | | |
|--|--------|-----------|-------|-------|--------|------|
| | elec | tive | emer | gency | | |
| | N | % | N | % | N | % |
| Bearing type (head/insert) | 23,726 | | 3,179 | | 26,905 | |
| Ceramics-Ceramics | 3,467 | 14.6 | 204 | 6.4 | 3,671 | 13.6 |
| Ceramics-Metal | 118 | 0.5 | 29 | 0.9 | 147 | 0.5 |
| Ceramics-Polyethylene | 14,616 | 61.6 | 1,623 | 51.1 | 16,239 | 60.4 |
| Metal-Ceramics | 39 | 0.2 | 7 | 0.2 | 46 | 0.2 |
| Metal-Metal | 108 | 0.5 | 22 | 0.7 | 130 | 0.5 |
| Metal-Polyethylene | 2,176 | 9.2 | 588 | 18.5 | 2,764 | 10.3 |
| Procedures that do not report the implantation of a head and an insert | 3,202 | 13.5 | 706 | 22.2 | 3,908 | 14.5 |

Table 2.12. Hip. Number of revision procedures by type of bearing

| | Revis | ion (*) |
|--|-------|---------|
| | N | % |
| Type of bearing (head/insert) | 2,082 | |
| Ceramics-Ceramics | 42 | 2.0 |
| Ceramics-Metal | 31 | 1.5 |
| Ceramics-Polyethylene | 625 | 30.0 |
| Metal-Ceramics | 1 | 0.0 |
| Metal-Metal | 45 | 2.2 |
| Metal-Polyethylene | 327 | 15.7 |
| Procedures that do not report the implantation of a head and an insert | 1,011 | 48.6 |

^(*) Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision

Table 2.13. Hip. Number of total replacement procedures by stem type, by procedure type

| | | Total rep | | TOTAL | | | |
|--|--------|-----------|-------|-------|--------|------|--|
| | elec | tive | emer | gency | | | |
| | N | % | N | % | N | % | |
| Stem type | 23,726 | | 3,179 | | 26,905 | | |
| Uncemented | 19,232 | 81.1 | 2,370 | 74.6 | 21,602 | 80.3 | |
| Modular | 1,122 | 5.8 | 307 | 13.0 | 1,429 | 6.6 | |
| Non-modular | 18,110 | 94.2 | 2,063 | 87.0 | 20,173 | 93.4 | |
| Straight | 13,563 | 74.9 | 1,851 | 89.7 | 15,414 | 76.4 | |
| Anatomical | 2,076 | 11.5 | 125 | 6.1 | 2,201 | 10.9 | |
| Conservative | 2,470 | 13.6 | 87 | 4.2 | 2,557 | 12.7 | |
| Cemented | 1,375 | 5.8 | 412 | 13.0 | 1,787 | 6.6 | |
| Modular | 52 | 3.8 | 12 | 2.9 | 64 | 3.6 | |
| Non-modular | 1,323 | 96.2 | 400 | 97.1 | 1,723 | 96.4 | |
| Straight | 1,205 | 91.1 | 376 | 94.0 | 1,581 | 91.8 | |
| Anatomical | 101 | 7.6 | 23 | 5.8 | 124 | 7.2 | |
| Conservative | 17 | 1.3 | 1 | 0.3 | 18 | 1.0 | |
| Other stem type / Stem type not reported | 3,119 | 13.1 | 397 | 12.5 | 3,516 | 13.1 | |

Table 2.14. Knee. Number of procedures analysed, by procedure type

| | N | % |
|--|--------|------|
| Procedure type | 29,198 | |
| Primary | 27,588 | 94.5 |
| - total | 23,064 | 83.6 |
| - unicompartmental | 4,524 | 16.4 |
| Revision | 1,610 | 5.5 |
| Partial revision | 358 | 22.2 |
| Total revision | 1,184 | 73.5 |
| Removal of prosthesis | 47 | 2.9 |
| Primary patella implant on existing prosthesis | 21 | 1.3 |

Table 2.15. Knee. Number of procedures by type of hospital, by procedure type

| | Primary | | | | Revision (*) | | TOTAL | |
|-----------------------------------|---------|------|------------------|------|--------------|------|--------|------|
| | total | | unicompartmental | | | | | |
| | N | % | N | % | N | % | N | % |
| Type of hospital | 23,064 | | 4,524 | | 1,610 | | 29,198 | |
| Public hospitals | 9,081 | 39.4 | 922 | 20.4 | 553 | 34.3 | 10,556 | 36.2 |
| Private hospitals, accredited | 12,257 | 53.1 | 3,277 | 72.4 | 987 | 61.3 | 16,521 | 56.6 |
| Private hospitals, not accredited | 1,726 | 7.5 | 325 | 7.2 | 70 | 4.3 | 2,121 | 7.3 |

^(*) Total or partial revision, removal of prosthesis, primary patella implant on existing prosthesis, spacer revision

Table 2.16. Knee. Number of procedures by patient gender and age group, by procedure type

| | | Pr | imary | | Revision (*) | | TOTAL | |
|---------------------|--------|------|----------|----------|--------------|------|--------|------|
| | to | tal | unicompa | rtmental | | | | |
| | N | % | N | % | N | % | N | % |
| Gender | 23,064 | | 4,524 | | 1,610 | | 29,198 | |
| Male | 7,380 | 32.0 | 1,692 | 37.4 | 516 | 32.0 | 9,588 | 32.8 |
| Female | 15,684 | 68.0 | 2,832 | 62.6 | 1,094 | 68.0 | 19,610 | 67.2 |
| Age group by gender | | | | | | | | |
| Male | 7,380 | | 1,692 | | 516 | | 9,588 | |
| Mean age | 70 | | 67 | | 67 | | 69 | |
| Standard deviation | 9 | | 10 | | 10 | | 9 | |
| <45 | 67 | 0.9 | 23 | 1.4 | 7 | 1.4 | 97 | 1.0 |
| 45 - 54 | 376 | 5.1 | 148 | 8.7 | 27 | 5.2 | 551 | 5.7 |
| 55 - 64 | 1,381 | 18.7 | 424 | 25.1 | 93 | 18.0 | 1,898 | 19.8 |
| 65 - 74 | 3,132 | 42.4 | 730 | 43.1 | 203 | 39.3 | 4,065 | 42.4 |
| 75 - 84 | 2,319 | 31.4 | 339 | 20.0 | 174 | 33.7 | 2,832 | 29.5 |
| ≥85 | 105 | 1.4 | 28 | 1.7 | 12 | 2.3 | 145 | 1.5 |
| Female | 15,684 | | 2,832 | | 1,094 | | 19,610 | |
| Mean age | 71 | | 68 | | 68 | | 71 | |
| Standard deviation | 8 | | 9 | | 9 | | 8 | |
| <45 | 42 | 0.3 | 23 | 0.8 | 1 | 0.1 | 66 | 0.3 |
| 45 - 54 | 510 | 3.3 | 195 | 6.9 | 42 | 3.8 | 747 | 3.8 |
| 55 - 64 | 2,481 | 15.8 | 632 | 22.3 | 166 | 15.2 | 3,279 | 16.7 |
| 65 - 74 | 6,940 | 44.2 | 1,219 | 43.0 | 477 | 43.6 | 8,636 | 44.0 |
| 75 - 84 | 5,398 | 34.4 | 714 | 25.2 | 368 | 33.6 | 6,480 | 33.0 |
| ≥85 | 313 | 2.0 | 49 | 1.7 | 40 | 3.7 | 402 | 2.0 |

^(*) Total or partial revision, removal of prosthesis, primary patella implant on existing prosthesis, spacer revision

Table 2.17. Knee. Number of procedures by operated side and surgical approach, by procedure type

| | | Primary | | | | Revision (*) | | TOTAL | |
|-------------------------------|--------|----------------------|-------|------|-------|--------------|--------|-------|--|
| | tot | tal unicompartmental | | | | | | | |
| | N | % | N | % | N | % | N | % | |
| Operated side | 23,064 | | 4,524 | | 1,610 | | 29,198 | | |
| Right | 12,178 | 52.8 | 2,203 | 48.7 | 830 | 51.6 | 15,211 | 52.1 | |
| Left | 10,565 | 45.8 | 1,977 | 43.7 | 768 | 47.7 | 13,310 | 45.6 | |
| Bilateral | 321 | 1.4 | 344 | 7.6 | 12 | 0.7 | 677 | 2.3 | |
| Surgical approach | 23,064 | | 4,524 | | 1,610 | | 29,198 | | |
| Medial parapatellar | 20,274 | 87.9 | 3,123 | 69.0 | 1,388 | 86.2 | 24,785 | 84.9 | |
| Lateral parapatellar | 477 | 2.1 | 249 | 5.5 | 25 | 1.6 | 751 | 2.6 | |
| Mid-vastus | 1,244 | 5.4 | 356 | 7.9 | 102 | 6.3 | 1,702 | 5.8 | |
| Minimally invasive mid-vastus | 365 | 1.6 | 537 | 11.9 | 37 | 2.3 | 939 | 3.2 | |
| Quad-sparing | 59 | 0.3 | 124 | 2.7 | 3 | 0.2 | 186 | 0.6 | |
| Sub-vastus | 258 | 1.1 | 42 | 0.9 | 10 | 0.6 | 310 | 1.1 | |
| Minimally invasive sub-vastus | 86 | 0.4 | 54 | 1.2 | 5 | 0.3 | 145 | 0.5 | |
| V Quadriceps | 1 | 0.0 | 0 | 0.0 | 4 | 0.2 | 5 | 0.0 | |
| Tibial tuberosity osteotomy | 8 | 0.0 | 2 | 0.0 | 5 | 0.3 | 15 | 0.1 | |
| Other | 292 | 1.3 | 37 | 0.8 | 31 | 1.9 | 360 | 1.2 | |

 $^{(^\}star) \, \text{Total or partial revision, removal of prosthesis, primary patella implant on existing prosthesis, spacer revision}$

Table 2.18. Knee. Number of primary procedures, by indication for surgery and previous procedure

| | | Prin | nary | | TOTAL | | |
|-------------------------------|--------|------|----------|----------|--------|------|--|
| | to | tal | unicompa | rtmental | | | |
| | N | % | N | % | N | % | |
| Indication for surgery | 23,064 | | 4,524 | | 27,588 | | |
| Primary osteoarthritis | 21,764 | 94.4 | 4,184 | 92.5 | 25,948 | 94.1 | |
| Post-traumatic osteoarthritis | 368 | 1.6 | 50 | 1.1 | 418 | 1.5 | |
| Rheumatoid arthritis | 122 | 0.5 | 3 | 0.1 | 125 | 0.5 | |
| Neoplasia | 15 | 0.1 | 2 | 0.0 | 17 | 0.1 | |
| Osteonecrosis | 107 | 0.5 | 170 | 3.8 | 277 | 1.0 | |
| Other | 688 | 3.0 | 115 | 2.5 | 803 | 2.9 | |
| Previous procedure | 23,064 | | 4,524 | | 27,588 | | |
| None | 20,549 | 89.1 | 4,191 | 92.6 | 24,740 | 89.7 | |
| Arthrodesis | 8 | 0.0 | 1 | 0.0 | 9 | 0.0 | |
| Osteotomy | 152 | 0.7 | 12 | 0.3 | 164 | 0.6 | |
| Arthroscopy | 503 | 2.2 | 176 | 3.9 | 679 | 2.5 | |
| Osteosynthesis | 142 | 0.6 | 11 | 0.2 | 153 | 0.6 | |
| Other | 1,710 | 7.4 | 133 | 2.9 | 1,843 | 6.7 | |

Table 2.19. Knee. Number of revision procedures, by indication for surgery and previous procedure

| | Revision (* |) |
|---|-------------|------|
| | N | % |
| Indication for surgery | 1,610 | |
| Aseptic loosening of several components | 418 | 26.0 |
| Aseptic loosening of femur | 88 | 5.5 |
| Aseptic loosening of tibia | 160 | 9.9 |
| Aseptic loosening of patella | 4 | 0.2 |
| Wear | 38 | 2.4 |
| Dislocation | 39 | 2.4 |
| Instability | 56 | 3.5 |
| Periprosthetic fracture | 19 | 1.2 |
| Implant fracture | 23 | 1.4 |
| Fractured spacer | 2 | 0.1 |
| Infection | 249 | 15.5 |
| Stiffness | 25 | 1.6 |
| Disease progression | 37 | 2.3 |
| Pain | 269 | 16.7 |
| Other | 183 | 11.4 |
| Previous procedure | 1,610 | |
| Primary total | 992 | 61.6 |
| Primary unicompartmental | 267 | 16.6 |
| Revision of knee replacement | 116 | 7.2 |
| Spacer | 143 | 8.9 |
| Other | 92 | 5.7 |

^(*) Total or partial revision, removal of prosthesis, primary patella implant on existing prosthesis, spacer revision

Table 2.20. Knee. Number of procedures analysed for the implanted device, by procedure type

| | N | % |
|--------------------|--------|------|
| Procedure type | 28,497 | |
| Primary | 26,917 | 94.5 |
| - total | 22,695 | 84.3 |
| - unicompartmental | 4,222 | 15.7 |
| Revision | 1,580 | 5.5 |

Table 2.21. Knee. Number of procedures by type of fixation, by procedure type

| | | F | Primary | | Revisio | on (*) | TOTA | ٨L |
|--|--------|------|----------|-----------|---------|--------|--------|------|
| | tota | al | unicompa | artmental | | | | |
| | N | % | N | % | N | % | N | % |
| Fixation | 22,695 | | 4,222 | | 1,580 | | 28,497 | |
| Patella not implanted | 19,930 | 87.8 | 4,145 | 98.2 | 601 | 38.0 | 24,676 | 86.6 |
| Cemented (femoral and tibial components) | 13,342 | 66.9 | 2,724 | 65.7 | 380 | 63.2 | 16,446 | 66.6 |
| Cemented femoral component and uncemented tibial component | 678 | 3.4 | 284 | 6.9 | 49 | 8.2 | 1,011 | 4.1 |
| Only cemented femoral component | 0 | 0.0 | 0 | 0.0 | 17 | 2.8 | 17 | 0.1 |
| Uncemented femoral component and cemented tibial component | 1,043 | 5.2 | 373 | 9.0 | 54 | 9.0 | 1,470 | 6.0 |
| Uncemented | 4,867 | 24.4 | 764 | 18.4 | 12 | 2.0 | 5,643 | 22.9 |
| Only uncemented femoral component | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Only cemented tibial component | 0 | 0.0 | 0 | 0.0 | 29 | 4.8 | 29 | 0.1 |
| Only uncemented tibial component | 0 | 0.0 | 0 | 0.0 | 9 | 1.5 | 9 | 0.0 |
| Fixaction declared "not applicable" for both femoral and tibial components | 0 | 0.0 | 0 | 0.0 | 51 | 8.5 | 51 | 0.2 |
| Patella implanted (cemented) | 2,187 | 9.6 | 63 | 1.5 | 296 | 18.7 | 2,546 | 8.9 |
| Cemented (femoral and tibial components) | 2,073 | 94.8 | 36 | 57.1 | 186 | 62.8 | 2,295 | 90.1 |
| Cemented femoral component and uncemented tibial component | 8 | 0.4 | 22 | 34.9 | 5 | 1.7 | 35 | 1.4 |
| Only cemented femoral component | 0 | 0.0 | 0 | 0.0 | 1 | 0.3 | 1 | 0.0 |
| Uncemented femoral component and cemented tibial component | 67 | 3.1 | 1 | 1.6 | 13 | 4.4 | 81 | 3.2 |
| Uncemented | 39 | 1.8 | 4 | 6.3 | 63 | 21.3 | 106 | 4.2 |
| Only uncemented femoral component | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Only cemented tibial component | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Only uncemented tibial component | | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Only patella | 0 | 0.0 | 0 | 0.0 | 28 | 9.5 | 28 | 1.1 |

(continued)

Table 2.21. (continued)

| | | F | Primary | | Revisio | on (*) | TOTAL | |
|--|------|------|----------|-----------|---------|--------|-------|------|
| | tota | ıl | unicompa | artmental | | | | |
| | N | % | N | % | N | % | N | % |
| Patella implanted (uncemented) | 578 | 2.5 | 14 | 0.3 | 683 | 43.2 | 1,275 | 4.5 |
| Cemented (femoral and tibial components) | 156 | 27.0 | 11 | 78.6 | 275 | 40.3 | 442 | 34.7 |
| Cemented femoral component and uncemented tibial component | 43 | 7.4 | 1 | 7.1 | 38 | 5.6 | 82 | 6.4 |
| Only cemented femoral component | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Uncemented femoral component and cemented tibial component | 48 | 8.3 | 0 | 0.0 | 70 | 10.2 | 118 | 9.3 |
| Uncemented | 331 | 57.3 | 2 | 14.3 | 300 | 43.9 | 633 | 49.6 |
| Only uncemented femoral component | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Only cemented tibial component | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Only uncemented tibial component | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Only patella | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Procedures not admitted to the analysis of the fixation mode | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |

^(*) Total or partial revision, removal of prosthesis, primary patella implant on existing prosthesis, spacer revision

42

Table 2.22. Knee. Number of primary procedures by type of tibial tray

| 1 31 331 | N | % |
|--------------------------|--------|------|
| | 14 | 70 |
| Type of tibial tray | 22,695 | |
| Mobile bearing | 5,452 | 24.0 |
| Cemented | 3,664 | 67.2 |
| Uncemented | 1,557 | 28.6 |
| Cementable | 231 | 4.2 |
| Fixed | 11,597 | 51.1 |
| Cemented | 11,107 | 95.8 |
| Uncemented | 458 | 3.9 |
| Cementable | 32 | 0.3 |
| Tibial tray not reported | 5,646 | 24.9 |

Table 2.23. Shoulder. Number of procedures analysed, by procedure type

| | N | % |
|---------------------|-----|------|
| Procedure type | 939 | |
| Primary | 920 | 98.0 |
| Total | 522 | 56.7 |
| - elective | 410 | 78.5 |
| - emergency | 112 | 21.5 |
| Partial replacement | 32 | 3.5 |
| Not specified | 366 | 39.8 |
| Revision | 19 | 2.0 |

Table 2.24. Shoulder. Number of procedures by prosthesis type implanted in the total replacement

| | N | % |
|---|-----|------|
| Type of prosthesis implanted in the total replacement | 522 | |
| Elective | 410 | 78.5 |
| - anatomical | 14 | 3.4 |
| - coating | 0 | 0.0 |
| - reverse | 314 | 76.6 |
| - interposition | 82 | 20.0 |
| Emergency | 112 | 21.5 |
| - anatomical | 7 | 6.3 |
| - coating | 0 | 0.0 |
| - reverse | 105 | 93.8 |
| - interposition | 0 | 0.0 |

Table 2.25. Shoulder. Number of procedures by type of hospital, by procedure type

| | | | | Pr | imary | | | | Revision (*) | | TOTAL | |
|-----------------------------------|----------|---------|-----------|-----|-------------|-----|-----------|------|--------------|-----|-------|------|
| | То | tal rep | laceme | nt | Partial | | Not | | | | | |
| | elective | | emergency | | replacement | | specified | | | | | |
| | N | % | N | % | N | % | N | % | N | % | N | % |
| Type of hospital | 410 | | 112 | | 32 | | 366 | | 19 | | 939 | |
| Public hospitals | 388 | 94.6 | 112 | 100 | 32 | 100 | 142 | 38.8 | 19 | 100 | 693 | 73.8 |
| Private hospitals, accredited | 15 | 3.7 | 0 | 0.0 | 0 | 0.0 | 129 | 35.2 | 0 | 0.0 | 144 | 15.3 |
| Private hospitals, not accredited | 7 | 1.7 | 0 | 0.0 | 0 | 0.0 | 95 | 26.0 | 0 | 0.0 | 102 | 10.9 |

 $^{(\}begin{tabular}{l} (\begin{tabular}{l} (\be$

Table 2.26. Shoulder. Number of procedures by patient gender and age group, by procedure type

| | | | | Prima | ry | | | | Revisi | on (*) | TOTAL | | |
|---------------------|-------|-----------|----------|-------|-------------|------|-----------|------|--------|--------|-------|------|--|
| | To | otal repl | lacement | | Par | | No | | | | | | |
| | elect | ive | emerg | ency | replacement | | specified | | | | | | |
| | N | % | N | % | N | % | N | % | N | % | N | % | |
| Gender | 410 | | 112 | | 32 | | 366 | | 19 | | 939 | | |
| Male | 126 | 30.7 | 21 | 18.8 | 15 | 46.9 | 91 | 24.9 | 7 | 36.8 | 260 | 27.7 | |
| Female | 284 | 69.3 | 91 | 81.3 | 17 | 53.1 | 275 | 75.1 | 12 | 63.2 | 679 | 72.3 | |
| Age group by gender | | | | | | | | | | | | | |
| Male | 126 | | 21 | | 15 | | 91 | | 7 | | 260 | | |
| Mean age | 67 | | 72 | | 53 | | 70 | | 63 | | 66 | | |
| Standard deviation | 10 | | 8 | | 10 | | 9 | | 12 | | 10 | | |
| <45 | 2 | 1.6 | 0 | 0.0 | 4 | 26.7 | 1 | 1.1 | 1 | 14.3 | 8 | 3.1 | |
| 45 - 54 | 19 | 15.1 | 0 | 0.0 | 4 | 26.7 | 6 | 6.6 | 1 | 14.3 | 30 | 11.5 | |
| 55 - 64 | 33 | 26.2 | 3 | 14.3 | 5 | 33.3 | 18 | 19.8 | 2 | 28.6 | 61 | 23.5 | |
| 65 - 74 | 47 | 37.3 | 10 | 47.6 | 1 | 6.7 | 42 | 46.2 | 3 | 42.9 | 103 | 39.6 | |
| 75 - 84 | 25 | 19.8 | 6 | 28.6 | 1 | 6.7 | 22 | 24.2 | 0 | 0.0 | 54 | 20.8 | |
| ≥85 | 0 | 0.0 | 2 | 9.5 | 0 | 0.0 | 2 | 2.2 | 0 | 0.0 | 4 | 1.5 | |
| Female | 284 | | 91 | | 17 | | 275 | | 12 | | 679 | | |
| Mean age | 71 | | 74 | | 67 | | 74 | | 68 | | 72 | | |
| Standard deviation | 7 | | 8 | | 9 | | 7 | | 6 | | 7 | | |
| <45 | 2 | 0.7 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 2 | 0.3 | |
| 45 - 54 | 6 | 2.1 | 1 | 1.1 | 1 | 5.9 | 1 | 0.4 | 0 | 0.0 | 9 | 1.3 | |
| 55 - 64 | 47 | 16.5 | 8 | 8.8 | 6 | 35.3 | 17 | 6.2 | 4 | 33.3 | 82 | 12.1 | |
| 65 - 74 | 148 | 52.1 | 40 | 44.0 | 7 | 41.2 | 120 | 43.6 | 6 | 50.0 | 321 | 47.3 | |
| 75 - 84 | 80 | 28.2 | 33 | 36.3 | 2 | 11.8 | 124 | 45.1 | 2 | 16.7 | 241 | 35.5 | |
| ≥85 | 1 | 0.4 | 9 | 9.9 | 1 | 5.9 | 13 | 4.7 | 0 | 0.0 | 24 | 3.5 | |

 $^{(\}begin{tabular}{l} (\begin{tabular}{l} (\be$

Table 2.27. Shoulder. Number of procedures by operated side and surgical approach, by procedure type

| | | | | Prin | nary | | | | Revisi | on (*) | TOTAL | | | |
|----------------------|------|-----------|--------------|-------|--------|-----------------------|-----|-----------|--------|-----------|-------|------|--|--|
| | T | otal repl | acemen | t | | Partial No | | | | | | | | |
| | elec | tive | ve emergency | | replac | replacement specified | | specified | | specified | | | | |
| | N | % | N | % | N | % | N | % | N | % | N | % | | |
| Operated side | 410 | | 112 | | 32 | | 366 | | 19 | | 939 | | | |
| Right | 263 | 64.1 | 55 | 49.1 | 18 | 56.3 | 238 | 65.0 | 13 | 68.4 | 587 | 62.5 | | |
| Left | 147 | 35.9 | 57 | 50.9 | 14 | 43.8 | 128 | 35.0 | 6 | 31.6 | 352 | 37.5 | | |
| Bilateral | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | | |
| Surgical approach | 410 | | 112 | | 32 | | 366 | | 19 | | 939 | | | |
| Deltopectoral | 322 | 78.5 | 112 | 100.0 | 26 | 81.3 | 344 | 94.0 | 16 | 84.2 | 820 | 87.3 | | |
| Transdeltoidea | 54 | 13.2 | 0 | 0.0 | 5 | 15.6 | 14 | 3.8 | 0 | 0.0 | 73 | 7.8 | | |
| Other | 34 | 8.3 | 0 | 0.0 | 1 | 3.1 | 8 | 2.2 | 3 | 15.8 | 46 | 4.9 | | |

^(*) Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision

Table 2.28. Shoulder. Number of primary procedures by indication for surgery and previous procedure

| procedure | | | | | | | | | | |
|---------------------------|--------|-----------|---------|-----------|-----|-------------|-----|-------|------|------|
| | | | | Prima | ry | | | | TOTA | L |
| | To | tal repla | acement | | Par | | No | | | |
| | electi | ve | emerg | emergency | | replacement | | ified | | |
| | N | % | N | % | N | % | N | % | N | % |
| Cause | 410 | | 112 | | 32 | | 366 | | 920 | |
| Eccentric osteoarthritis | 228 | 55.6 | 0 | 0.0 | 1 | 3.1 | 132 | 36.1 | 361 | 39.2 |
| Concentric osteoarthritis | 57 | 13.9 | 0 | 0.0 | 13 | 40.6 | 22 | 6.0 | 92 | 10.0 |
| Rheumatoid arthritis | 4 | 1.0 | 0 | 0.0 | 1 | 3.1 | 4 | 1.1 | 9 | 1.0 |
| Neoplasia | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Osteonecrosis | 2 | 0.5 | 0 | 0.0 | 1 | 3.1 | 7 | 1.9 | 10 | 1.1 |
| Fracture | 0 | 0.0 | 112 | 100.0 | 15 | 46.9 | 157 | 42.9 | 284 | 30.9 |
| Previous fracture | 17 | 4.1 | 0 | 0.0 | 1 | 3.1 | 0 | 0.0 | 18 | 2.0 |
| Other | 102 | 24.9 | 0 | 0.0 | 0 | 0.0 | 44 | 12.0 | 146 | 15.9 |
| Previous | 410 | | 112 | | 32 | | 366 | | 920 | |
| None | 375 | 91.5 | 111 | 99.1 | 31 | 96.9 | 365 | 99.7 | 882 | 95.9 |
| Osteosynthesis | 7 | 1.7 | 0 | 0.0 | 1 | 3.1 | 1 | 0.3 | 9 | 1.0 |
| Arthrotomy | 3 | 0.7 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 3 | 0.3 |
| Arthrodesis | 1 | 0.2 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 0.1 |
| Arthroscopy | 18 | 4.4 | 1 | 0.9 | 0 | 0.0 | 0 | 0.0 | 19 | 2.1 |
| Other | 6 | 1.5 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 6 | 0.7 |

Table 2.29. Shoulder. Number of revision procedures by indication for surgery and previous procedure

| | Revisior | ı (*) |
|-------------------------------|----------|-------|
| | N | % |
| Indication for surgery | 19 | |
| Instability | 3 | 15.8 |
| Lysis | 0 | 0.0 |
| Glenoid erosion | 1 | 5.3 |
| Prosthesis breakage | 0 | 0.0 |
| Dislocation | 3 | 15.8 |
| Periprosthetic fracture | 0 | 0.0 |
| Infection | 3 | 15.8 |
| Prosthesis removal outcomes | 0 | 0.0 |
| Aseptic mobilisation | 5 | 26.3 |
| Disease progression | 0 | 0.0 |
| Pain | 1 | 5.3 |
| Other | 3 | 15.8 |
| Previous procedure | 19 | |
| Primary | 17 | 89.5 |
| Removal | 1 | 5.3 |
| Shoulder replacement revision | 1 | 5.3 |
| Other | 0 | 0.0 |

^(*) Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision

Table 2.30. Shoulder. Number of procedures by type of fixation, by procedure type

| | | | | | | | | | Revisi | on (*) | T01 | AL |
|---|-------------------|----------|-----|-----------|---------------|------------------|---|-------|--------|--------|-----|------|
| | Total replacement | | | | tial ement | Not specified | | | | | | |
| | | elective | | emergency | | | | | | | | |
| | N | % | N | % | N | % | N | % | N | % | N | % |
| Fixation | 325 | | 110 | | 24 | | 3 | | 19 | | 481 | |
| Cemented (glenoid + stem) | 5 | 1.5 | 16 | 14.5 | 0 | 0.0 | 0 | 0.0 | 1 | 5.3 | 22 | 4.6 |
| Reverse hybrid (cemented glenoid and uncemented stem) | 3 | 0.9 | 2 | 1.8 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 5 | 1.0 |
| Only cemented glenoid | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Hybrid (uncemented glenoid and cemented stem) | 20 | 6.2 | 28 | 25.5 | 0 | 0.0 | 0 | 0.0 | 3 | 15.8 | 51 | 10.6 |
| Uncemented (glenoid + stem) | 297 | 91.4 | 64 | 58.2 | 0 | 0.0 | 3 | 100.0 | 9 | 47.4 | 373 | 77.5 |
| Only uncemented glenoid | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 3 | 15.8 | 3 | 0.6 |
| Only cemented stem | 0 | 0.0 | 0 | 0.0 | 7 | 29.2 | 0 | 0.0 | 1 | 5.3 | 8 | 1.7 |
| Only stem uncemented | 0 | 0.0 | 0 | 0.0 | 17 | 70.8 | 0 | 0.0 | 2 | 10.5 | 19 | 4.0 |
| Fixation declared "not applicable" for glenoid and stem | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |

^(*) Total or partial revision, removal of prosthesis, conversion from partial to total prosthesis, spacer revision

Figure 2.1. Flowchart of the RIAP data quality control process: procedures




Figure 2.2. Flowchart of the RIAP data quality control process: implanted medical devices





Figure 2.3. Hip. Types of bearing. Total replacement (elective procedures)



Note: the first component indicates the material of the head, the second the material of the insert.

CoP

MoP

CoC

Other (CoM, MoM, MoC)

Chart Legend:
CoP = Ceramics/Polyethylene, CoC = Ceramic/Ceramic,
MoP = Metal/Polyethylene, MoM = Metal/Metal,

Figure 2.4. Hip. Types of bearing. Total replacement (emergency)

66%

Note: the first component indicates the material of the head, the second the material of the insert.

CoM = Ceramic/Metal, MoC = Metal/Ceramic

APPENDIX

Table 1. Joint replacements in Italy, per year of discharge. 2001-2017 and estimate 2018

| ICD-9-CM Code | Procedure | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 |
|------------------|--|---------|---------|---------|---------|---------|---------|---------|---------|
| | Hip | 74,408 | 78,497 | 80,999 | 85,307 | 87,499 | 90,062 | 91,077 | 92,217 |
| 81.51 | Total hip replacement | 46,850 | 49,820 | 52,541 | 55,875 | 57,112 | 59,315 | 60,425 | 60,840 |
| | Total hip replacement (elective) | 40,067 | 42,284 | 44,505 | 47,047 | 47,908 | 49,657 | 50,684 | 51,110 |
| 81.52 | Partial hip replacement | 21,394 | 22,015 | 21,753 | 22,474 | 23,227 | 23,286 | 23,119 | 23,896 |
| 00.85(*) | Hip resurfacing | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| (**) | Revision of hip replacement | 6,164 | 6,662 | 6,705 | 6,958 | 7,160 | 7,461 | 7,533 | 7,481 |
| | Knee | 28,693 | 33,417 | 38,655 | 44,324 | 47,643 | 52,322 | 57,054 | 59,956 |
| 81.54 | Total knee replacement | 27,401 | 31,740 | 36,714 | 42,081 | 45,116 | 49,560 | 53,930 | 56,525 |
| (***) | Total knee replacement | 1,292 | 1,677 | 1,941 | 2,243 | 2,527 | 2,762 | 3,124 | 3,431 |
| | Shoulder | 1,559 | 1,700 | 1,866 | 2,296 | 2,517 | 2,888 | 3,255 | 3,412 |
| 81.80 | Total shoulder replacement | 709 | 820 | 948 | 1,263 | 1,462 | 1,695 | 2,048 | 2,190 |
| | Total shoulder replacement (elective) | 419 | 525 | 644 | 889 | 1,085 | 1,336 | 1,629 | 1,788 |
| 81.81 | Partial shoulder replacement | 850 | 880 | 918 | 1,033 | 1,055 | 1,193 | 1,207 | 1,222 |
| | Ankle | 95 | 115 | 147 | 175 | 179 | 257 | 268 | 284 |
| 81.56 | Total ankle replacement | 95 | 115 | 147 | 175 | 179 | 257 | 268 | 284 |
| | Other joints | 736 | 836 | 870 | 1,154 | 1,668 | 1,713 | 1,570 | 1,315 |
| 81.57 | Replacement of joint of foot and toe | 316 | 391 | 414 | 467 | 604 | 629 | 692 | 656 |
| 81.59 | Revision of joint replacement of lower extremity, not elsewhere classified | 214 | 181 | 173 | 338 | 672 | 588 | 365 | 140 |
| 81.73 | Total wrist replacement | 40 | 42 | 44 | 49 | 60 | 81 | 66 | 69 |
| 81.84 | Total elbow replacement | 90 | 145 | 162 | 203 | 251 | 317 | 314 | 311 |
| 81.97 | Revision of joint replacement of upper extremity | 76 | 77 | 77 | 97 | 81 | 98 | 133 | 139 |
| | Total | 105,491 | 114,565 | 122,537 | 133,256 | 139,506 | 147,242 | 153,224 | 157,184 |

^(°) Average annual increase

^(*) New code introduced on 1st January 2009

^(**) Codes: 00.70, 00.71, 00.72, 00.73 (introduced on 1st January 2009) and 81.53 (Hip replacement revision not specified) (***) Code: 00.80, 00.81, 00.82, 00.83, 00.84 (introduced on 1st January 2009) and 81.55 (Knee replacement revision not otherwise specified)

| 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | % (°) | Estimate 2018 |
|---------|---------|---------|---------|---------|---------|---------|---------|---------|-------|---------------|
| 93,241 | 95,348 | 96,125 | 98,585 | 100,844 | 102,652 | 105,803 | 108,906 | 112,375 | 2.6 | 115,308 |
| 61,601 | 61,775 | 62,664 | 64,503 | 66,257 | 68,190 | 71,178 | 74,660 | 77,787 | 3.2 | |
| 51,769 | 52,187 | 53,157 | 54,852 | 56,598 | 58,491 | 60,661 | 64,102 | 66,917 | 3.3 | |
| 23,393 | 24,847 | 25,091 | 25,346 | 25,979 | 26,141 | 26,222 | 25,879 | 26,101 | 1.3 | |
| 293 | 445 | 162 | 96 | 99 | 45 | 107 | 147 | 65 | -17.2 | |
| 7,954 | 8,281 | 8,208 | 8,640 | 8,509 | 8,276 | 8,296 | 8,220 | 8,422 | 2.0 | |
| 61,079 | 63,255 | 63,749 | 66,007 | 67,634 | 70,313 | 73,191 | 78,779 | 81,271 | 6.7 | 86,735 |
| 57,004 | 59,081 | 59,472 | 61,541 | 62,910 | 65,614 | 68,091 | 73,394 | 75,668 | 6.6 | |
| 4,075 | 4,174 | 4,277 | 4,466 | 4,724 | 4,699 | 5,100 | 5,385 | 5,603 | 9.6 | |
| 3,783 | 4,326 | 4,684 | 5,143 | 5,795 | 6,511 | 7,145 | 8,053 | 9,101 | 11.7 | 10,162 |
| 2,537 | 2,990 | 3,478 | 3,830 | 4,441 | 5,309 | 5,970 | 6,892 | 7,862 | 16.2 | |
| 2,093 | 2,382 | 2,816 | 3,042 | 3,480 | 4,085 | 4,474 | 5,226 | 5,922 | 18.0 | |
| 1,246 | 1,336 | 1,206 | 1,313 | 1,354 | 1,202 | 1,175 | 1,161 | 1,239 | 2.4 | |
| 256 | 255 | 298 | 313 | 330 | 387 | 482 | 546 | 600 | 12.2 | 673 |
| 256 | 255 | 298 | 313 | 330 | 387 | 482 | 546 | 600 | 12.2 | |
| 1,332 | 1,302 | 1,349 | 1,291 | 1,300 | 1,358 | 1,479 | 1,444 | 1,489 | 4.5 | 1,556 |
| 521 | 557 | 543 | 443 | 440 | 453 | 530 | 481 | 468 | 2.5 | |
| 187 | 128 | 107 | 83 | 96 | 103 | 102 | 74 | 77 | -6.2 | |
| 59 | 50 | 68 | 74 | 65 | 49 | 51 | 52 | 37 | -0.5 | |
| 402 | 402 | 434 | 447 | 473 | 491 | 523 | 549 | 608 | 12.7 | |
| 163 | 165 | 197 | 244 | 226 | 262 | 273 | 288 | 299 | 8.9 | |
| 159,691 | 164,486 | 166,205 | 171,339 | 175,903 | 181,221 | 188,100 | 197,728 | 204,836 | 4.2 | 214,434 |

Table 2. Hip. Number of procedures by region of admission and procedure type. 2016-2017

| Region of admission | Tota | l replacen | nent (elec | tive) | Total replacement (emergency) | | | | |
|-----------------------|--------|------------|------------|-------|-------------------------------|------|--------|------|--|
| | 20 | 16 | 20 | 17 | 20 | 16 | 20 | 17 | |
| | N | % | N | % | N | % | N | % | |
| Piedmont | 5,538 | 8.6 | 6,240 | 9.3 | 870 | 8.2 | 1,032 | 9.5 | |
| Aosta Valley | 146 | 0.2 | 158 | 0.2 | 16 | 0.2 | 15 | 0.1 | |
| Lombardy | 14,914 | 23.2 | 15,950 | 23.8 | 1,308 | 12.4 | 1,340 | 12.3 | |
| AP Bolzano | 1,009 | 1.6 | 999 | 1.5 | 114 | 1.1 | 87 | 0.8 | |
| AP Trento | 770 | 1.2 | 794 | 1.2 | 80 | 0.8 | 86 | 0.8 | |
| Veneto | 6,850 | 10.7 | 6,801 | 10.2 | 844 | 8.0 | 781 | 7.2 | |
| Friuli Venezia Giulia | 1,815 | 2.8 | 1,827 | 2.7 | 167 | 1.6 | 183 | 1.7 | |
| Liguria | 1,405 | 2.2 | 1,232 | 1.8 | 728 | 6.9 | 648 | 6.0 | |
| Emilia-Romagna | 7,244 | 11.3 | 7,611 | 11.4 | 753 | 7.1 | 839 | 7.7 | |
| Tuscany | 5,302 | 8.3 | 5,740 | 8.6 | 899 | 8.5 | 842 | 7.7 | |
| Umbria | 827 | 1.3 | 838 | 1.3 | 133 | 1.3 | 157 | 1.4 | |
| Marche | 1,273 | 2.0 | 1,265 | 1.9 | 373 | 3.5 | 448 | 4.1 | |
| Lazio | 5,113 | 8.0 | 5,369 | 8.0 | 1,113 | 10.5 | 1,199 | 11.0 | |
| Abruzzi | 1,424 | 2.2 | 1,429 | 2.1 | 291 | 2.8 | 296 | 2.7 | |
| Molise | 169 | 0.3 | 179 | 0.3 | 19 | 0.2 | 26 | 0.2 | |
| Campania | 3,151 | 4.9 | 3,167 | 4.7 | 891 | 8.4 | 908 | 8.3 | |
| Apulia | 2,490 | 3.9 | 2,428 | 3.6 | 593 | 5.6 | 531 | 4.9 | |
| Basilicata | 280 | 0.4 | 250 | 0.4 | 80 | 0.8 | 88 | 0.8 | |
| Calabria | 879 | 1.4 | 933 | 1.4 | 271 | 2.6 | 268 | 2.5 | |
| Sicily | 2,728 | 4.2 | 2,872 | 4.3 | 869 | 8.2 | 914 | 8.4 | |
| Sardinia | 915 | 1.4 | 894 | 1.3 | 153 | 1.4 | 188 | 1.7 | |
| Italy | 64,242 | 100 | 66,976 | 100 | 10,565 | 100 | 10,876 | 100 | |
| % of national volume | 59.0 | | 59.6 | | 9.7 | | 9.7 | | |

| | Partial rep | lacement | | | Revi | sion | | Total | | | | |
|--------|-------------|----------|------|-------|------|-------|------|---------|------|---------|------|--|
| 20 | 16 | 20 | 17 | 20 | 16 | 20 | 17 | 20 | 16 | 20 | 17 | |
| N | % | N | % | N | % | N | % | N | % | N | % | |
| 1,925 | 7.4 | 1,805 | 6.9 | 751 | 9.1 | 766 | 9.1 | 9,084 | 8.3 | 9,843 | 8.8 | |
| 88 | 0.3 | 73 | 0.3 | 28 | 0.3 | 22 | 0.3 | 278 | 0.3 | 268 | 0.2 | |
| 4,978 | 19.2 | 5,131 | 19.7 | 1,916 | 23.3 | 2,059 | 24.4 | 23,116 | 21.2 | 24,480 | 21.8 | |
| 174 | 0.7 | 216 | 0.8 | 143 | 1.7 | 129 | 1.5 | 1,440 | 1.3 | 1,431 | 1.3 | |
| 272 | 1.1 | 326 | 1.2 | 83 | 1.0 | 96 | 1.1 | 1,205 | 1.1 | 1,302 | 1.2 | |
| 2,195 | 8.5 | 2,291 | 8.8 | 648 | 7.9 | 723 | 8.6 | 10,537 | 9.7 | 10,596 | 9.4 | |
| 839 | 3.2 | 827 | 3.2 | 233 | 2.8 | 181 | 2.1 | 3,054 | 2.8 | 3,018 | 2.7 | |
| 638 | 2.5 | 688 | 2.6 | 269 | 3.3 | 235 | 2.8 | 3,040 | 2.8 | 2,803 | 2.5 | |
| 2,544 | 9.8 | 2,351 | 9.0 | 1,026 | 12.5 | 946 | 11.2 | 11,567 | 10.6 | 11,747 | 10.5 | |
| 2,095 | 8.1 | 2,081 | 8.0 | 806 | 9.8 | 744 | 8.8 | 9,102 | 8.4 | 9,407 | 8.4 | |
| 550 | 2.1 | 610 | 2.3 | 129 | 1.6 | 105 | 1.2 | 1,639 | 1.5 | 1,710 | 1.5 | |
| 582 | 2.2 | 605 | 2.3 | 137 | 1.7 | 177 | 2.1 | 2,365 | 2.2 | 2,495 | 2.2 | |
| 2,145 | 8.3 | 2,033 | 7.8 | 560 | 6.8 | 622 | 7.4 | 8,931 | 8.2 | 9,223 | 8.2 | |
| 642 | 2.5 | 607 | 2.3 | 140 | 1.7 | 163 | 1.9 | 2,497 | 2.3 | 2,495 | 2.2 | |
| 155 | 0.6 | 207 | 0.8 | 24 | 0.3 | 15 | 0.2 | 367 | 0.3 | 427 | 0.4 | |
| 1,569 | 6.1 | 1,660 | 6.4 | 437 | 5.3 | 481 | 5.7 | 6,048 | 5.6 | 6,216 | 5.5 | |
| 1,447 | 5.6 | 1,549 | 5.9 | 299 | 3.6 | 318 | 3.8 | 4,829 | 4.4 | 4,826 | 4.3 | |
| 253 | 1.0 | 268 | 1.0 | 39 | 0.5 | 28 | 0.3 | 652 | 0.6 | 634 | 0.6 | |
| 686 | 2.7 | 581 | 2.2 | 98 | 1.2 | 126 | 1.5 | 1,934 | 1.8 | 1,908 | 1.7 | |
| 1,591 | 6.1 | 1,600 | 6.1 | 369 | 4.5 | 397 | 4.7 | 5,557 | 5.1 | 5,783 | 5.1 | |
| 511 | 2.0 | 592 | 2.3 | 85 | 1.0 | 89 | 1.1 | 1,664 | 1.5 | 1,763 | 1.6 | |
| 25,879 | 100 | 26,101 | 100 | 8,220 | 100 | 8,422 | 100 | 108,906 | 100 | 112,375 | 100 | |
| 23.8 | | 23.2 | | 7.5 | | 7.5 | | 100.0 | | 100.0 | | |

Table 3. Hip. Primary total replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017

| Region of admission | Class of volun | Class of volume | | | | | | | | | | | |
|-----------------------|----------------|-----------------|------|------|------|------|--|--|--|--|--|--|--|
| | 1- | 50 | 51- | 100 | 101- | 200 | | | | | | | |
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | | | | | | | |
| | N | N | N | N | N | N | | | | | | | |
| Piedmont | 17 | 15 | 13 | 10 | 12 | 13 | | | | | | | |
| Aosta Valley | 0 | 0 | 2 | 2 | 0 | 0 | | | | | | | |
| Lombardy | 37 | 33 | 31 | 31 | 24 | 25 | | | | | | | |
| AP Bolzano | 2 | 3 | 4 | 4 | 3 | 3 | | | | | | | |
| AP Trento | 2 | 3 | 2 | 1 | 3 | 3 | | | | | | | |
| Veneto | 10 | 9 | 14 | 9 | 14 | 20 | | | | | | | |
| Friuli Venezia Giulia | 6 | 2 | 8 | 4 | 5 | 6 | | | | | | | |
| Liguria | 3 | 5 | 3 | 2 | 6 | 7 | | | | | | | |
| Emilia-Romagna | 17 | 12 | 23 | 22 | 21 | 21 | | | | | | | |
| Tuscany | 18 | 16 | 16 | 17 | 6 | 6 | | | | | | | |
| Umbria | 5 | 4 | 6 | 5 | 3 | 4 | | | | | | | |
| Marche | 7 | 8 | 4 | 2 | 5 | 6 | | | | | | | |
| Lazio | 48 | 44 | 15 | 20 | 13 | 13 | | | | | | | |
| Abruzzi | 9 | 7 | 6 | 8 | 3 | 3 | | | | | | | |
| Molise | 3 | 3 | 2 | 2 | 0 | 0 | | | | | | | |
| Campania | 47 | 45 | 13 | 13 | 8 | 10 | | | | | | | |
| Apulia | 20 | 21 | 11 | 11 | 9 | 9 | | | | | | | |
| Basilicata | 4 | 4 | 2 | 1 | 1 | 1 | | | | | | | |
| Calabria | 15 | 14 | 1 | 3 | 3 | 2 | | | | | | | |
| Sicily | 45 | 46 | 21 | 16 | 5 | 7 | | | | | | | |
| Sardinia | 15 | 14 | 5 | 6 | 2 | 2 | | | | | | | |
| Italy | 330 | 308 | 202 | 189 | 146 | 161 | | | | | | | |
| % of national volume | 43.0 | 41.1 | 26.3 | 25.2 | 19.0 | 21.5 | | | | | | | |

| 201- | 200 | >3 | 00 | | To | tal. | |
|------|------|------|------|-----|------|------|------|
| 2016 | 2017 | 2016 | | 20 | | 20 | 47 |
| | | | 2017 | | % | | % |
| N 7 | N | N | N | N | | N | |
| | 5 | 4 | 8 | 53 | 6.9 | 51 | 6.8 |
| 0 | 0 | 0 | 0 | 2 | 0.3 | 2 | 0.3 |
| 6 | 7 | 14 | 13 | 112 | 14.6 | 109 | 14.6 |
| 1 | 1 | 0 | 0 | 10 | 1.3 | 11 | 1.5 |
| 1 | 1 | 0 | 0 | 8 | 1.0 | 8 | 1.1 |
| 6 | 4 | 6 | 6 | 50 | 6.5 | 48 | 6.4 |
| 2 | 2 | 0 | 1 | 21 | 2.7 | 15 | 2.0 |
| 2 | 1 | 1 | 1 | 15 | 2.0 | 16 | 2.1 |
| 5 | 7 | 3 | 2 | 69 | 9.0 | 64 | 8.5 |
| 6 | 5 | 5 | 7 | 51 | 6.6 | 51 | 6.8 |
| 0 | 0 | 0 | 0 | 14 | 1.8 | 13 | 1.7 |
| 2 | 2 | 0 | 0 | 18 | 2.3 | 18 | 2.4 |
| 4 | 4 | 3 | 3 | 83 | 10.8 | 84 | 11.2 |
| 2 | 1 | 0 | 1 | 20 | 2.6 | 20 | 2.7 |
| 0 | 0 | 0 | 0 | 5 | 0.7 | 5 | 0.7 |
| 4 | 3 | 0 | 0 | 72 | 9.4 | 71 | 9.5 |
| 2 | 2 | 0 | 0 | 42 | 5.5 | 43 | 5.7 |
| 0 | 0 | 0 | 0 | 7 | 0.9 | 6 | 0.8 |
| 1 | 2 | 0 | 0 | 20 | 2.6 | 21 | 2.8 |
| 2 | 2 | 0 | 0 | 73 | 9.5 | 71 | 9.5 |
| 0 | 0 | 0 | 0 | 22 | 2.9 | 22 | 2.9 |
| 53 | 49 | 36 | 42 | 767 | 100 | 749 | 100 |
| 6.9 | 6.5 | 4.7 | 5.6 | 100 | | 100 | |

Table 4. Hip. Revision. Number of hospitals by region of admission and class of volume. 2016 and 2017

| Region of admission | | | | | | Class of | volume | | | | | |
|-----------------------|------|------|------|------|------|----------|--------|------|-----|------|-----|------|
| | 1-1 | 10 | 11- | 25 | 26- | 50 | >5 | 60 | | Tot | al | |
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 20 | 16 | 201 | 17 |
| | N | N | N | N | N | N | N | N | N | % | N | % |
| Piedmont | 19 | 20 | 23 | 21 | 4 | 6 | 1 | 1 | 47 | 6.9 | 48 | 7.3 |
| Aosta Valley | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | 2 | 0.3 | 2 | 0.3 |
| Lombardy | 54 | 49 | 36 | 32 | 8 | 12 | 7 | 8 | 105 | 15.5 | 101 | 15.3 |
| AP Bolzano | 3 | 4 | 3 | 3 | 2 | 2 | 0 | 0 | 8 | 1.2 | 9 | 1.4 |
| AP Trento | 4 | 5 | 0 | 1 | 2 | 1 | 0 | 0 | 6 | 0.9 | 7 | 1.1 |
| Veneto | 25 | 24 | 17 | 16 | 7 | 7 | 0 | 0 | 49 | 7.2 | 47 | 7.1 |
| Friuli Venezia Giulia | 12 | 10 | 8 | 3 | 1 | 0 | 0 | 1 | 21 | 3.1 | 14 | 2.1 |
| Liguria | 7 | 8 | 3 | 5 | 1 | 2 | 2 | 1 | 13 | 1.9 | 16 | 2.4 |
| Emilia-Romagna | 37 | 30 | 16 | 21 | 8 | 4 | 2 | 2 | 63 | 9.3 | 57 | 8.6 |
| Tuscany | 24 | 23 | 11 | 9 | 8 | 10 | 3 | 3 | 46 | 6.8 | 45 | 6.8 |
| Umbria | 8 | 8 | 3 | 2 | 1 | 1 | 0 | 0 | 12 | 1.8 | 11 | 1.7 |
| Marche | 11 | 9 | 6 | 6 | 0 | 1 | 0 | 0 | 17 | 2.5 | 16 | 2.4 |
| Lazio | 47 | 50 | 12 | 13 | 4 | 5 | 1 | 1 | 64 | 9.4 | 69 | 10.5 |
| Abruzzi | 13 | 12 | 3 | 6 | 1 | 0 | 0 | 0 | 17 | 2.5 | 18 | 2.7 |
| Molise | 3 | 3 | 1 | 0 | 0 | 0 | 0 | 0 | 4 | 0.6 | 3 | 0.5 |
| Campania | 51 | 47 | 6 | 6 | 1 | 2 | 1 | 2 | 59 | 8.7 | 57 | 8.6 |
| Apulia | 30 | 25 | 9 | 8 | 1 | 2 | 0 | 0 | 40 | 5.9 | 35 | 5.3 |
| Basilicata | 3 | 5 | 2 | 1 | 0 | 0 | 0 | 0 | 5 | 0.7 | 6 | 0.9 |
| Calabria | 14 | 14 | 1 | 2 | 1 | 1 | 0 | 0 | 16 | 2.4 | 17 | 2.6 |
| Sicily | 57 | 54 | 9 | 7 | 1 | 2 | 0 | 0 | 67 | 9.9 | 63 | 9.6 |
| Sardinia | 15 | 17 | 2 | 1 | 0 | 0 | 0 | 0 | 17 | 2.5 | 18 | 2.7 |
| Italy | 438 | 418 | 172 | 164 | 51 | 58 | 17 | 19 | 678 | 100 | 659 | 100 |
| % of national volume | 64.6 | 63.4 | 25.4 | 24.9 | 7.5 | 8.8 | 2.5 | 2.9 | 100 | | 100 | |


Table 5. Hip. Percent distribution of hospital discharges by patient gender and age group, and by procedure type. 2016 and 2017

| procedure type. 2010 | Tot replace (elect | ment | Tot replace (emerg | ement | Par replac | | Revi | sion | Tot | al |
|----------------------|--------------------------|------|--------------------------|-------|---------------|------|------|------|------|------|
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 |
| | % | % | % | % | % | % | % | % | % | % |
| Gender | | | | | | | | | | |
| Male | 46.0 | 46.3 | 26.1 | 27.2 | 27.0 | 27.8 | 39.5 | 40.2 | 39.1 | 39.7 |
| Female | 54.0 | 53.7 | 73.9 | 72.8 | 73.0 | 72.2 | 60.5 | 59.8 | 60.9 | 60.3 |
| Age (male) | | | | | | | | | | |
| Mean age | 65.4 | 65.4 | 71.9 | 71.9 | 83.5 | 83.2 | 69.1 | 69.5 | 69.1 | 69.1 |
| Standard deviation | 11.9 | 11.8 | 12.1 | 12.3 | 8.4 | 9.4 | 12.6 | 12.5 | 13.2 | 13.3 |
| Age group | | | | | | | | | | |
| 0-44 | 5.3 | 5.1 | 2.1 | 2.3 | 0.5 | 0.9 | 4.3 | 4.2 | 4.2 | 4.2 |
| 45-54 | 13.4 | 13.3 | 7.6 | 7.7 | 0.5 | 0.9 | 10.0 | 8.9 | 10.6 | 10.6 |
| 55-64 | 23.2 | 23.8 | 14.8 | 15.7 | 1.7 | 2.0 | 16.7 | 17.4 | 18.6 | 19.2 |
| 65-74 | 33.3 | 33.2 | 29.1 | 27.4 | 7.3 | 7.8 | 29.9 | 28.9 | 28.5 | 28.3 |
| 75-84 | 22.8 | 22.3 | 32.1 | 32.3 | 38.7 | 37.6 | 31.3 | 31.9 | 26.7 | 26.2 |
| 85+ | 2.0 | 2.2 | 14.3 | 14.6 | 51.3 | 50.9 | 7.9 | 8.8 | 11.4 | 11.5 |
| Age (female) | | | | | | | | | | |
| Mean age | 69.6 | 69.6 | 74.3 | 74.0 | 84.3 | 84.4 | 73.4 | 73.7 | 74.6 | 74.5 |
| Standard deviation | 10.8 | 10.8 | 9.7 | 9.7 | 7.1 | 7.4 | 11.3 | 11.1 | 11.7 | 11.7 |
| Age group | | | | | | | | | | |
| 0-44 | 2.3 | 2.4 | 0.3 | 0.4 | 0.1 | 0.2 | 2.1 | 1.6 | 1.4 | 1.5 |
| 45-54 | 7.3 | 7.2 | 2.8 | 2.6 | 0.3 | 0.4 | 4.7 | 4.9 | 4.6 | 4.6 |
| 55-64 | 17.6 | 18.0 | 11.9 | 12.6 | 1.1 | 1.0 | 11.3 | 11.5 | 11.7 | 12.1 |
| 65-74 | 35.4 | 35.0 | 33.5 | 33.6 | 6.0 | 5.8 | 27.4 | 26.8 | 26.2 | 26.1 |
| 75-84 | 33.1 | 33.0 | 37.2 | 37.3 | 39.4 | 39.2 | 41.1 | 40.8 | 35.9 | 35.8 |
| 85+ | 4.3 | 4.4 | 14.3 | 13.5 | 53.1 | 53.4 | 13.3 | 14.4 | 20.1 | 19.9 |

Table 6. Hip. Percent distribution of hospital discharges by mode of discharge and by procedure type. 2016 and 2017

| Mode of discharge | Total replace- ment (elective) | | ment | Total replace- ment (emer- gency) | | Partial replacement | | Revision | | Total | |
|---|-----------------------------------|------|------|---|------|------------------------|------|----------|------|-------|--|
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | |
| | % | % | % | % | % | % | % | % | % | % | |
| Deceased | 0.2 | 0.1 | 1.2 | 1.2 | 3.0 | 2.8 | 1.2 | 1.0 | 1.0 | 0.9 | |
| Ordinary discharge | 54.0 | 50.3 | 53.9 | 51.2 | 50.1 | 47.4 | 54.6 | 50.7 | 53.1 | 49.7 | |
| Discharge to a residential health care | 1.0 | 1.5 | 6.0 | 6.7 | 9.6 | 9.4 | 3.3 | 3.9 | 3.7 | 4.0 | |
| Discharge to hospital at home | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.1 | |
| Discharge against medical advice | 0.1 | 0.1 | 0.4 | 0.3 | 0.4 | 0.4 | 0.2 | 0.3 | 0.2 | 0.2 | |
| Transfer to an acute admission unit of a different hospital | 2.2 | 2.1 | 4.4 | 4.9 | 4.5 | 4.3 | 3.1 | 2.8 | 3.0 | 2.9 | |
| Transfer in the same hospital | 23.1 | 27.5 | 9.3 | 10.4 | 7.3 | 10.5 | 18.4 | 21.8 | 17.7 | 21.4 | |
| Transfer to an inpatient rehabilitation hospital | 15.0 | 17.6 | 18.5 | 22.9 | 18.5 | 22.3 | 14.8 | 18.2 | 16.1 | 19.3 | |
| Discharge to a nursing home | 4.4 | 0.7 | 6.1 | 2.2 | 6.1 | 2.6 | 4.3 | 1.2 | 5.0 | 1.3 | |

Figure 1. Hip. Elective total replacement. Inter-regional mobility (attraction and escape indices) in 2016 (a) and in 2017 (b)



NOTE: For regions with low number of procedures (i.e Umbria, Molise and Basilicata) inter-regional mobility indices might be biased

Table 7. Knee. Number of procedures by region of admission and procedure type. 2016-2017

| Region of admission | Total replacement | | | | | | | | |
|-----------------------|-------------------|------|--------|------|--|--|--|--|--|
| | 20 | 16 | 20 | 17 | | | | | |
| | N | % | N | % | | | | | |
| Piedmont | 5,030 | 6.9 | 5,969 | 7.9 | | | | | |
| Aosta Valley | 167 | 0.2 | 136 | 0.2 | | | | | |
| Lombardy | 15,111 | 20.6 | 15,880 | 21.0 | | | | | |
| AP Bolzano | 881 | 1.2 | 903 | 1.2 | | | | | |
| AP Trento | 682 | 0.9 | 709 | 0.9 | | | | | |
| Veneto | 7,830 | 10.7 | 7,531 | 10.0 | | | | | |
| Friuli Venezia Giulia | 1,707 | 2.3 | 1,761 | 2.3 | | | | | |
| Liguria | 1,315 | 1.8 | 1,198 | 1.6 | | | | | |
| Emilia-Romagna | 7,752 | 10.6 | 7,695 | 10.2 | | | | | |
| Tuscany | 7,149 | 9.7 | 7,541 | 10.0 | | | | | |
| Umbria | 1,300 | 1.8 | 1,415 | 1.9 | | | | | |
| Marche | 1,681 | 2.3 | 1,815 | 2.4 | | | | | |
| Lazio | 5,651 | 7.7 | 6,185 | 8.2 | | | | | |
| Abruzzi | 1,988 | 2.7 | 1,915 | 2.5 | | | | | |
| Molise | 227 | 0.3 | 200 | 0.3 | | | | | |
| Campania | 3,281 | 4.5 | 3,537 | 4.7 | | | | | |
| Apulia | 3,426 | 4.7 | 3,370 | 4.5 | | | | | |
| Basilicata | 387 | 0.5 | 264 | 0.3 | | | | | |
| Calabria | 1,350 | 1.8 | 1,439 | 1.9 | | | | | |
| Sicily | 4,959 | 6.8 | 4,795 | 6.3 | | | | | |
| Sardinia | 1,520 | 2.1 | 1,410 | 1.9 | | | | | |
| Italy | 73,394 | 100 | 75,668 | 100 | | | | | |
| % of national volume | 93.2 | | 93.1 | | | | | | |

| | Revi | sion | | Total | | | | | |
|-------|------|-------|------|--------|------|--------|------|--|--|
| 20 | 16 | 20 | 2017 | | 2016 | | 17 | | |
| N | % | N | % | N | % | N | % | | |
| 397 | 7.4 | 433 | 7.7 | 5,427 | 6.9 | 6,402 | 7.9 | | |
| 7 | 0.1 | 9 | 0.2 | 174 | 0.2 | 145 | 0.2 | | |
| 1,299 | 24.1 | 1,407 | 25.1 | 16,410 | 20.8 | 17,287 | 21.3 | | |
| 75 | 1.4 | 98 | 1.7 | 956 | 1.2 | 1,001 | 1.2 | | |
| 43 | 0.8 | 33 | 0.6 | 725 | 0.9 | 742 | 0.9 | | |
| 507 | 9.4 | 447 | 8.0 | 8,337 | 10.6 | 7,978 | 9.8 | | |
| 112 | 2.1 | 110 | 2.0 | 1,819 | 2.3 | 1,871 | 2.3 | | |
| 147 | 2.7 | 134 | 2.4 | 1,462 | 1.9 | 1,332 | 1.6 | | |
| 729 | 13.5 | 702 | 12.5 | 8,481 | 10.8 | 8,397 | 10.3 | | |
| 625 | 11.6 | 648 | 11.6 | 7,774 | 9.9 | 8,189 | 10.1 | | |
| 81 | 1.5 | 86 | 1.5 | 1,381 | 1.8 | 1,501 | 1.8 | | |
| 75 | 1.4 | 115 | 2.1 | 1,756 | 2.2 | 1,930 | 2.4 | | |
| 376 | 7.0 | 393 | 7.0 | 6,027 | 7.7 | 6,578 | 8.1 | | |
| 69 | 1.3 | 89 | 1.6 | 2,057 | 2.6 | 2,004 | 2.5 | | |
| 10 | 0.2 | 5 | 0.1 | 237 | 0.3 | 205 | 0.3 | | |
| 175 | 3.2 | 192 | 3.4 | 3,456 | 4.4 | 3,729 | 4.6 | | |
| 158 | 2.9 | 185 | 3.3 | 3,584 | 4.5 | 3,555 | 4.4 | | |
| 35 | 0.6 | 18 | 0.3 | 422 | 0.5 | 282 | 0.3 | | |
| 56 | 1.0 | 73 | 1.3 | 1,406 | 1.8 | 1,512 | 1.9 | | |
| 353 | 6.6 | 380 | 6.8 | 5,312 | 6.7 | 5,175 | 6.4 | | |
| 56 | 1.0 | 46 | 0.8 | 1,576 | 2.0 | 1,456 | 1.8 | | |
| 5,385 | 100 | 5,603 | 100 | 78,779 | 100 | 81,271 | 100 | | |
| 6.8 | | 6.9 | | 100 | | 100 | | | |

Table 8. Knee. Primary total replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017

| Region of admission | Class of volume | | | | | | | | | | |
|-----------------------|-----------------|------|------|------|---------|------|--|--|--|--|--|
| | 1- | 50 | 51- | 100 | 101-200 | | | | | | |
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | | | | | |
| | N | N | N | N | N | N | | | | | |
| Piedmont | 22 | 20 | 11 | 10 | 9 | 8 | | | | | |
| Aosta Valley | 1 | 1 | 0 | 1 | 1 | 0 | | | | | |
| Lombardy | 47 | 41 | 25 | 34 | 23 | 13 | | | | | |
| AP Bolzano | 4 | 5 | 2 | 2 | 4 | 4 | | | | | |
| AP Trento | 1 | 2 | 5 | 4 | 2 | 2 | | | | | |
| Veneto | 19 | 19 | 11 | 16 | 10 | 6 | | | | | |
| Friuli Venezia Giulia | 9 | 4 | 5 | 3 | 6 | 7 | | | | | |
| Liguria | 7 | 9 | 3 | 3 | 3 | 5 | | | | | |
| Emilia-Romagna | 25 | 21 | 16 | 16 | 16 | 15 | | | | | |
| Tuscany | 18 | 19 | 12 | 9 | 9 | 9 | | | | | |
| Umbria | 2 | 2 | 7 | 7 | 4 | 4 | | | | | |
| Marche | 8 | 9 | 6 | 5 | 1 | 1 | | | | | |
| Lazio | 51 | 47 | 12 | 14 | 12 | 12 | | | | | |
| Abruzzi | 13 | 10 | 3 | 5 | 2 | 2 | | | | | |
| Molise | 3 | 4 | 0 | 0 | 1 | 1 | | | | | |
| Campania | 50 | 39 | 12 | 18 | 5 | 6 | | | | | |
| Apulia | 23 | 22 | 9 | 9 | 6 | 6 | | | | | |
| Basilicata | 4 | 4 | 1 | 1 | 2 | 1 | | | | | |
| Calabria | 14 | 14 | 1 | 3 | 1 | 1 | | | | | |
| Sicily | 39 | 40 | 16 | 16 | 5 | 5 | | | | | |
| Sardinia | 13 | 15 | 2 | 1 | 3 | 3 | | | | | |
| Italy | 373 | 347 | 159 | 177 | 125 | 111 | | | | | |
| % of national volume | 49.6 | 47.1 | 21.1 | 24.0 | 16.6 | 15.1 | | | | | |

| 201- | 200 | >3 | 00 | Total | | | | | |
|-----------|-----------|-----------|-----------|-------------|------|------|------|--|--|
| | 2016 2017 | | | | | 2017 | | | |
| 2016 N | 2017 N | 2016 N | 2017 N | 2016 N % | | N | % | | |
| 4 | 3 | 4 | 7 | N 50 | 6.6 | 48 | 6.5 | | |
| 0 | 0 | 0 | | 2 | 0.3 | 2 | 0.3 | | |
| | | · | 0 | | | | | | |
| 6 | 10 | 12 | 14 | 113 | 15.0 | 112 | 15.2 | | |
| 0 | 0 | 0 | 0 | 10 | 1.3 | 11 | 1.5 | | |
| 0 | 0 | 0 | 0 | 8 | 1.1 | 8 | 1.1 | | |
| 3 | 1 | 7 | 8 | 50 | 6.6 | 50 | 6.8 | | |
| 0 | 0 | 1 | 1 | 21 | 2.8 | 15 | 2.0 | | |
| 1 | 0 | 1 | 0 | 15 | 2.0 | 17 | 2.3 | | |
| 6 | 6 | 4 | 5 | 67 | 8.9 | 63 | 8.5 | | |
| 4 | 4 | 7 | 8 | 50 | 6.6 | 49 | 6.6 | | |
| 1 | 1 | 0 | 0 | 14 | 1.9 | 14 | 1.9 | | |
| 2 | 2 | 1 | 1 | 18 | 2.4 | 18 | 2.4 | | |
| 0 | 0 | 5 | 5 | 80 | 10.6 | 78 | 10.6 | | |
| 1 | 1 | 2 | 2 | 21 | 2.8 | 20 | 2.7 | | |
| 0 | 0 | 0 | 0 | 4 | 0.5 | 5 | 0.7 | | |
| 4 | 4 | 0 | 0 | 71 | 9.4 | 67 | 9.1 | | |
| 4 | 4 | 1 | 1 | 43 | 5.7 | 42 | 5.7 | | |
| 0 | 0 | 0 | 0 | 7 | 0.9 | 6 | 0.8 | | |
| 1 | 1 | 2 | 2 | 19 | 2.5 | 21 | 2.8 | | |
| 8 | 8 | 1 | 1 | 69 | 9.2 | 70 | 9.5 | | |
| 1 | 1 | 1 | 1 | 20 | 2.7 | 21 | 2.8 | | |
| 46 | 46 | 49 | 56 | 752 | 100 | 737 | 100 | | |
| 6.1 | 6.2 | 6.5 | 7.6 | 100 | | 100 | | | |

Table 9. Knee. Revision. Number of hospitals by region of admission and class of volume. 2016 and 2017

| Region of admission | Class of volume | | | | | | | | | | | |
|-----------------------|-----------------|------|------|-------|------|-------|------|------|-----|-------|-----|------|
| | 1-1 | 1-10 | | 11-25 | | 26-50 | | >50 | | Total | | |
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 20 | 16 | 201 | 17 |
| | N | N | N | N | N | N | N | N | N | % | N | % |
| Piedmont | 30 | 30 | 8 | 7 | 2 | 5 | 1 | 0 | 41 | 7.2 | 42 | 7.3 |
| Aosta Valley | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0.4 | 1 | 0.2 |
| Lombardy | 63 | 66 | 18 | 12 | 9 | 10 | 4 | 5 | 94 | 16.6 | 93 | 16.1 |
| AP Bolzano | 5 | 2 | 4 | 5 | 0 | 0 | 0 | 0 | 9 | 1.6 | 7 | 1.2 |
| AP Trento | 6 | 8 | 1 | 0 | 0 | 0 | 0 | 0 | 7 | 1.2 | 8 | 1.4 |
| Veneto | 24 | 28 | 13 | 8 | 0 | 3 | 3 | 1 | 40 | 7.1 | 40 | 6.9 |
| Friuli Venezia Giulia | 16 | 10 | 3 | 3 | 0 | 1 | 0 | 0 | 19 | 3.4 | 14 | 2.4 |
| Liguria | 5 | 7 | 5 | 5 | 0 | 1 | 1 | 0 | 11 | 1.9 | 13 | 2.3 |
| Emilia-Romagna | 40 | 39 | 12 | 12 | 2 | 3 | 3 | 3 | 57 | 10.1 | 57 | 9.9 |
| Tuscany | 29 | 29 | 9 | 6 | 4 | 6 | 3 | 2 | 45 | 7.9 | 43 | 7.5 |
| Umbria | 9 | 7 | 2 | 4 | 0 | 0 | 0 | 0 | 11 | 1.9 | 11 | 1.9 |
| Marche | 14 | 12 | 1 | 1 | 0 | 2 | 0 | 0 | 15 | 2.6 | 15 | 2.6 |
| Lazio | 40 | 46 | 8 | 4 | 4 | 4 | 0 | 0 | 52 | 9.2 | 54 | 9.4 |
| Abruzzi | 8 | 13 | 3 | 3 | 0 | 0 | 0 | 0 | 11 | 1.9 | 16 | 2.8 |
| Molise | 2 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0.4 | 3 | 0.5 |
| Campania | 41 | 38 | 2 | 5 | 1 | 0 | 0 | 0 | 44 | 7.8 | 43 | 7.5 |
| Apulia | 27 | 30 | 3 | 6 | 0 | 0 | 0 | 0 | 30 | 5.3 | 36 | 6.2 |
| Basilicata | 3 | 4 | 1 | 0 | 0 | 0 | 0 | 0 | 4 | 0.7 | 4 | 0.7 |
| Calabria | 9 | 10 | 3 | 2 | 0 | 0 | 0 | 0 | 12 | 2.1 | 12 | 2.1 |
| Sicily | 36 | 41 | 9 | 10 | 2 | 2 | 0 | 0 | 47 | 8.3 | 53 | 9.2 |
| Sardinia | 14 | 11 | 0 | 1 | 0 | 0 | 0 | 0 | 14 | 2.5 | 12 | 2.1 |
| Italy | 423 | 435 | 105 | 94 | 24 | 37 | 15 | 11 | 567 | 100 | 577 | 100 |
| % of national volume | 74.6 | 75.4 | 18.5 | 16.3 | 4.2 | 6.4 | 2.6 | 1.9 | 100 | | 100 | |


Table 10. Knee. Percent distribution of hospital discharges by patient gender and age group, and by procedure type. 2016 and 2017

| | Total rep | lacement | Revi | sion | Total | | |
|--------------------|-----------|----------|------|------|-------|------|--|
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | |
| | % | % | % | % | % | % | |
| Gender | | | | | | | |
| Male | 32.4 | 32.9 | 32.1 | 33.1 | 32.4 | 32.9 | |
| Female | 67.6 | 67.1 | 67.9 | 66.9 | 67.6 | 67.1 | |
| Age (male) | | | | | | | |
| Mean age | 69.5 | 69.4 | 67.7 | 68.6 | 69.4 | 69.4 | |
| Standard deviation | 9.4 | 9.3 | 12.3 | 11.4 | 9.6 | 9.4 | |
| Age group | | | | | | | |
| 0-44 | 1.6 | 1.2 | 4.5 | 3.3 | 1.8 | 1.3 | |
| 45-54 | 5.1 | 5.3 | 8.8 | 6.9 | 5.3 | 5.4 | |
| 55-64 | 18.1 | 19.0 | 16.9 | 19.1 | 18.0 | 19.0 | |
| 65-74 | 42.1 | 42.4 | 37.8 | 37.1 | 41.8 | 42.0 | |
| 75-84 | 31.3 | 30.3 | 28.9 | 30.1 | 31.1 | 30.3 | |
| 85+ | 1.8 | 1.8 | 3.2 | 3.5 | 1.9 | 1.9 | |
| Age (female) | | | | | | | |
| Mean age | 70.8 | 70.8 | 70.7 | 70.6 | 70.8 | 70.8 | |
| Standard deviation | 8.2 | 8.2 | 9.2 | 9.3 | 8.3 | 8.2 | |
| Age group | | | | | | | |
| 0-44 | 0.5 | 0.4 | 1.2 | 1.3 | 0.6 | 0.5 | |
| 45-54 | 3.2 | 3.2 | 3.5 | 3.9 | 3.2 | 3.3 | |
| 55-64 | 15.9 | 16.2 | 16.6 | 17.3 | 15.9 | 16.3 | |
| 65-74 | 43.9 | 44.6 | 40.5 | 39.8 | 43.7 | 44.3 | |
| 75-84 | 34.6 | 33.6 | 35.3 | 34.3 | 34.7 | 33.7 | |
| 85+ | 1.8 | 1.9 | 2.9 | 3.3 | 1.9 | 2.0 | |

Table 11. Knee. Percent distribution of hospital discharges by mode of discharge and by procedure type. 2016 and 2017

| Mode of discharge | | lacement tive) | Revi | sion | Total | | |
|---|------|-------------------|------|------|-------|------|--|
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | |
| | % | % | % | % | % | % | |
| Deceased | 0.1 | 0.0 | 0.1 | 0.2 | 0.1 | 0.0 | |
| Ordinary discharge | 53.3 | 48.5 | 56.5 | 51.8 | 53.5 | 48.7 | |
| Discharge to a residential health care | 0.7 | 1.0 | 1.1 | 1.1 | 0.7 | 1.0 | |
| Discharge to hospital at home | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | |
| Discharge against medical advice | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | |
| Transfer to an acute admission unit of a different hospital | 1.9 | 2.0 | 2.1 | 1.9 | 1.9 | 2.0 | |
| Transfer in the same hospital | 24.7 | 31.1 | 24.7 | 30.2 | 24.7 | 31.1 | |
| Transfer to an inpatient rehabilitation hospital | 15.9 | 16.7 | 12.6 | 14.1 | 15.7 | 16.5 | |
| Discharge to a nursing home | 3.3 | 0.4 | 2.7 | 0.5 | 3.2 | 0.4 | |

Figure 2. Knee. Total replacement. Inter-regional mobility (attraction and escape indices) in 2016 (a) and in 2017 (b)



NOTE: For regions with low number of procedures (i.e Umbria, Molise and Basilicata) inter-regional mobility indices might be biased

Table 12. Shoulder. Number of procedures by region of admission and procedure type. 2016-2017

| Region of admission | То | tal replacen | nent (electiv | re) | Total replacement (emergency) | | | |
|-----------------------|-------|--------------|---------------|------|-------------------------------|------|-------|------|
| | 2016 | | 20 | 17 | 20 | 16 | 20 | 17 |
| | N | % | N | % | N | % | N | % |
| Piedmont | 492 | 9.4 | 489 | 8.3 | 125 | 7.5 | 149 | 7.7 |
| Aosta Valley | 4 | 0.1 | 11 | 0.2 | 2 | 0.1 | 5 | 0.3 |
| Lombardy | 915 | 17.5 | 1,174 | 19.8 | 335 | 20.1 | 398 | 20.5 |
| AP Bolzano | 32 | 0.6 | 48 | 0.8 | 2 | 0.1 | 9 | 0.5 |
| AP Trento | 43 | 0.8 | 41 | 0.7 | 25 | 1.5 | 30 | 1.5 |
| Veneto | 494 | 9.5 | 514 | 8.7 | 199 | 11.9 | 208 | 10.7 |
| Friuli Venezia Giulia | 145 | 2.8 | 108 | 1.8 | 29 | 1.7 | 26 | 1.3 |
| Liguria | 75 | 1.4 | 39 | 0.7 | 27 | 1.6 | 21 | 1.1 |
| Emilia-Romagna | 732 | 14.0 | 828 | 14.0 | 140 | 8.4 | 161 | 8.3 |
| Tuscany | 549 | 10.5 | 635 | 10.7 | 94 | 5.6 | 85 | 4.4 |
| Umbria | 83 | 1.6 | 106 | 1.8 | 27 | 1.6 | 62 | 3.2 |
| Marche | 98 | 1.9 | 113 | 1.9 | 75 | 4.5 | 76 | 3.9 |
| Lazio | 566 | 10.8 | 622 | 10.5 | 166 | 10.0 | 212 | 10.9 |
| Abruzzi | 128 | 2.4 | 160 | 2.7 | 44 | 2.6 | 61 | 3.1 |
| Molise | 3 | 0.1 | 5 | 0.1 | 5 | 0.3 | 6 | 0.3 |
| Campania | 222 | 4.2 | 306 | 5.2 | 45 | 2.7 | 85 | 4.4 |
| Apulia | 230 | 4.4 | 243 | 4.1 | 132 | 7.9 | 117 | 6.0 |
| Basilicata | 44 | 0.8 | 14 | 0.2 | 5 | 0.3 | 6 | 0.3 |
| Calabria | 54 | 1.0 | 68 | 1.1 | 20 | 1.2 | 29 | 1.5 |
| Sicily | 289 | 5.5 | 362 | 6.1 | 151 | 9.1 | 174 | 9.0 |
| Sardinia | 27 | 0.5 | 35 | 0.6 | 19 | 1.1 | 21 | 1.1 |
| Italy | 5,225 | 100 | 5,921 | 100 | 1,667 | 100 | 1,941 | 100 |
| % of national volume | 64.9 | | 65.1 | | 20.7 | | 21.3 | |

| | Partial rep | lacement | | Total | | | | |
|-------|-------------|----------|------|-------|------|-------|------|--|
| 20 | 16 | 20 | 17 | 20 | 16 | 20 | 17 | |
| N | % | N | % | N | % | N | % | |
| 34 | 2.9 | 40 | 3.2 | 651 | 8.1 | 678 | 7.4 | |
| 0 | 0.0 | 0 | 0.0 | 6 | 0.1 | 16 | 0.2 | |
| 234 | 20.2 | 186 | 15.0 | 1,484 | 18.4 | 1,758 | 19.3 | |
| 13 | 1.1 | 9 | 0.7 | 47 | 0.6 | 66 | 0.7 | |
| 25 | 2.2 | 17 | 1.4 | 93 | 1.2 | 88 | 1.0 | |
| 268 | 23.1 | 343 | 27.7 | 961 | 11.9 | 1,065 | 11.7 | |
| 30 | 2.6 | 36 | 2.9 | 204 | 2.5 | 170 | 1.9 | |
| 18 | 1.6 | 10 | 0.8 | 120 | 1.5 | 70 | 0.8 | |
| 72 | 6.2 | 82 | 6.6 | 944 | 11.7 | 1,071 | 11.8 | |
| 81 | 7.0 | 74 | 6.0 | 724 | 9.0 | 794 | 8.7 | |
| 52 | 4.5 | 76 | 6.1 | 162 | 2.0 | 244 | 2.7 | |
| 18 | 1.6 | 10 | 0.8 | 191 | 2.4 | 199 | 2.2 | |
| 104 | 9.0 | 123 | 9.9 | 836 | 10.4 | 957 | 10.5 | |
| 15 | 1.3 | 23 | 1.9 | 187 | 2.3 | 244 | 2.7 | |
| 5 | 0.4 | 1 | 0.1 | 13 | 0.2 | 12 | 0.1 | |
| 54 | 4.7 | 52 | 4.2 | 321 | 4.0 | 443 | 4.9 | |
| 60 | 5.2 | 77 | 6.2 | 422 | 5.2 | 437 | 4.8 | |
| 9 | 0.8 | 1 | 0.1 | 58 | 0.7 | 21 | 0.2 | |
| 7 | 0.6 | 10 | 0.8 | 81 | 1.0 | 107 | 1.2 | |
| 34 | 2.9 | 48 | 3.9 | 474 | 5.9 | 584 | 6.4 | |
| 28 | 2.4 | 21 | 1.7 | 74 | 0.9 | 77 | 0.8 | |
| 1,161 | 100 | 1,239 | 100 | 8,053 | 100 | 9,101 | 100 | |
| 14.4 | | 13.6 | | 100 | | 100 | | |

Table 13. Shoulder. Total replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017

| Region of admission | Class of volun | Class of volume | | | | | | | | | |
|-----------------------|----------------|-----------------|------|------|------|------|--|--|--|--|--|
| | 1- | -4 | 5- | .9 | 10- | ·14 | | | | | |
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | | | | | |
| | N | N | N | N | N | N | | | | | |
| Piedmont | 15 | 12 | 7 | 5 | 10 | 8 | | | | | |
| Aosta Valley | 2 | 0 | 0 | 2 | 0 | 0 | | | | | |
| Lombardy | 34 | 23 | 29 | 32 | 11 | 11 | | | | | |
| AP Bolzano | 7 | 5 | 2 | 1 | 0 | 2 | | | | | |
| AP Trento | 3 | 2 | 3 | 0 | 0 | 2 | | | | | |
| Veneto | 5 | 6 | 10 | 8 | 8 | 11 | | | | | |
| Friuli Venezia Giulia | 5 | 2 | 3 | 4 | 5 | 5 | | | | | |
| Liguria | 9 | 9 | 3 | 5 | 1 | 1 | | | | | |
| Emilia-Romagna | 17 | 17 | 14 | 11 | 3 | 9 | | | | | |
| Tuscany | 14 | 14 | 4 | 8 | 4 | 1 | | | | | |
| Umbria | 1 | 2 | 4 | 2 | 3 | 3 | | | | | |
| Marche | 7 | 6 | 4 | 4 | 2 | 3 | | | | | |
| Lazio | 26 | 28 | 7 | 7 | 4 | 2 | | | | | |
| Abruzzi | 9 | 8 | 6 | 3 | 1 | 0 | | | | | |
| Molise | 1 | 2 | 1 | 1 | 0 | 0 | | | | | |
| Campania | 21 | 18 | 6 | 8 | 2 | 5 | | | | | |
| Apulia | 14 | 15 | 4 | 7 | 5 | 5 | | | | | |
| Basilicata | 0 | 0 | 2 | 0 | 0 | 2 | | | | | |
| Calabria | 4 | 7 | 4 | 3 | 3 | 1 | | | | | |
| Sicily | 23 | 27 | 12 | 11 | 5 | 6 | | | | | |
| Sardinia | 10 | 7 | 0 | 1 | 0 | 1 | | | | | |
| Italy | 227 | 210 | 125 | 123 | 67 | 78 | | | | | |
| % of national volume | 40.7 | 37.0 | 22.4 | 21.7 | 12.0 | 13.8 | | | | | |

| 15 | -24 | > | 24 | Total | | | | |
|------|------|------|------|-------|------|-----|------|--|
| 2016 | 2017 | 2016 | 2017 | 20 | 16 | 20 | 2017 | |
| N | N | N | N | N | % | N | % | |
| 7 | 8 | 7 | 8 | 46 | 8.2 | 41 | 7.2 | |
| 0 | 0 | 0 | 0 | 2 | 0.4 | 2 | 0.4 | |
| 8 | 15 | 15 | 15 | 97 | 17.4 | 96 | 16.9 | |
| 0 | 1 | 0 | 0 | 9 | 1.6 | 9 | 1.6 | |
| 2 | 2 | 0 | 0 | 8 | 1.4 | 6 | 1.1 | |
| 12 | 9 | 8 | 10 | 43 | 7.7 | 44 | 7.8 | |
| 2 | 0 | 1 | 1 | 16 | 2.9 | 12 | 2.1 | |
| 0 | 0 | 1 | 0 | 14 | 2.5 | 15 | 2.6 | |
| 5 | 2 | 13 | 15 | 52 | 9.3 | 54 | 9.5 | |
| 4 | 2 | 8 | 10 | 34 | 6.1 | 35 | 6.2 | |
| 2 | 2 | 0 | 2 | 10 | 1.8 | 11 | 1.9 | |
| 0 | 3 | 3 | 1 | 16 | 2.9 | 17 | 3.0 | |
| 7 | 9 | 10 | 10 | 54 | 9.7 | 56 | 9.9 | |
| 1 | 4 | 2 | 2 | 19 | 3.4 | 17 | 3.0 | |
| 0 | 0 | 0 | 0 | 2 | 0.4 | 3 | 0.5 | |
| 1 | 2 | 4 | 5 | 34 | 6.1 | 38 | 6.7 | |
| 3 | 3 | 4 | 3 | 30 | 5.4 | 33 | 5.8 | |
| 0 | 0 | 1 | 0 | 3 | 0.5 | 2 | 0.4 | |
| 0 | 1 | 0 | 1 | 11 | 2.0 | 13 | 2.3 | |
| 3 | 3 | 4 | 6 | 47 | 8.4 | 53 | 9.3 | |
| 0 | 1 | 1 | 0 | 11 | 2.0 | 10 | 1.8 | |
| 57 | 67 | 82 | 89 | 558 | 100 | 567 | 100 | |
| 10.2 | 11.8 | 14.7 | 15.7 | 100 | | 100 | | |

Table 14. Shoulder. Partial replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017

| Region of admission | Class of volun | Class of volume | | | | | | | | | |
|-----------------------|----------------|-----------------|------|------|------|------|--|--|--|--|--|
| | 1- | -4 | 5- | -9 | 10- | 14 | | | | | |
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | | | | | |
| | N | N | N | N | N | N | | | | | |
| Piedmont | 18 | 17 | 1 | 2 | 0 | 0 | | | | | |
| Aosta Valley | 0 | 0 | 0 | 0 | 0 | 0 | | | | | |
| Lombardy | 34 | 48 | 7 | 3 | 5 | 2 | | | | | |
| AP Bolzano | 3 | 1 | 1 | 1 | 0 | 0 | | | | | |
| AP Trento | 3 | 4 | 1 | 1 | 1 | 0 | | | | | |
| Veneto | 19 | 22 | 7 | 6 | 4 | 4 | | | | | |
| Friuli Venezia Giulia | 10 | 5 | 0 | 1 | 1 | 1 | | | | | |
| Liguria | 4 | 1 | 2 | 1 | 0 | 0 | | | | | |
| Emilia-Romagna | 19 | 21 | 1 | 1 | 0 | 0 | | | | | |
| Tuscany | 19 | 20 | 3 | 1 | 0 | 0 | | | | | |
| Umbria | 6 | 4 | 1 | 2 | 0 | 0 | | | | | |
| Marche | 5 | 3 | 2 | 1 | 0 | 0 | | | | | |
| Lazio | 28 | 27 | 4 | 5 | 2 | 1 | | | | | |
| Abruzzi | 8 | 8 | 1 | 2 | 0 | 0 | | | | | |
| Molise | 3 | 1 | 0 | 0 | 0 | 0 | | | | | |
| Campania | 15 | 11 | 2 | 3 | 0 | 1 | | | | | |
| Apulia | 13 | 13 | 4 | 3 | 1 | 2 | | | | | |
| Basilicata | 4 | 1 | 0 | 0 | 0 | 0 | | | | | |
| Calabria | 3 | 2 | 0 | 1 | 0 | 0 | | | | | |
| Sicily | 12 | 22 | 2 | 2 | 0 | 0 | | | | | |
| Sardinia | 5 | 6 | 1 | 1 | 0 | 0 | | | | | |
| Italy | 231 | 237 | 40 | 37 | 14 | 11 | | | | | |
| % of national volume | 77.0 | 79.0 | 13.3 | 12.3 | 4.7 | 3.7 | | | | | |

| 15- | -24 | >2 | 24 | Total | | | | |
|------|------|------|------|-------|------|-----|------|--|
| 2016 | 2017 | 2016 | 2017 | 20 | 16 | 20 | 2017 | |
| N | N | N | N | N | % | N | % | |
| 0 | 0 | 0 | 0 | 19 | 6.3 | 19 | 6.3 | |
| 0 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0.0 | |
| 2 | 1 | 1 | 1 | 49 | 16.3 | 55 | 18.3 | |
| 0 | 0 | 0 | 0 | 4 | 1.3 | 2 | 0.7 | |
| 0 | 0 | 0 | 0 | 5 | 1.7 | 5 | 1.7 | |
| 5 | 2 | 1 | 4 | 36 | 12.0 | 38 | 12.7 | |
| 0 | 0 | 0 | 0 | 11 | 3.7 | 7 | 2.3 | |
| 0 | 0 | 0 | 0 | 6 | 2.0 | 2 | 0.7 | |
| 2 | 2 | 0 | 0 | 22 | 7.3 | 24 | 8.0 | |
| 0 | 0 | 1 | 1 | 23 | 7.7 | 22 | 7.3 | |
| 1 | 1 | 0 | 1 | 8 | 2.7 | 8 | 2.7 | |
| 0 | 0 | 0 | 0 | 7 | 2.3 | 4 | 1.3 | |
| 0 | 0 | 0 | 1 | 34 | 11.3 | 34 | 11.3 | |
| 0 | 0 | 0 | 0 | 9 | 3.0 | 10 | 3.3 | |
| 0 | 0 | 0 | 0 | 3 | 1.0 | 1 | 0.3 | |
| 1 | 0 | 0 | 0 | 18 | 6.0 | 15 | 5.0 | |
| 0 | 1 | 0 | 0 | 18 | 6.0 | 19 | 6.3 | |
| 0 | 0 | 0 | 0 | 4 | 1.3 | 1 | 0.3 | |
| 0 | 0 | 0 | 0 | 3 | 1.0 | 3 | 1.0 | |
| 0 | 0 | 0 | 0 | 14 | 4.7 | 24 | 8.0 | |
| 1 | 0 | 0 | 0 | 7 | 2.3 | 7 | 2.3 | |
| 12 | 7 | 3 | 8 | 300 | 100 | 300 | 100 | |
| 4.0 | 2.3 | 1.0 | 2.7 | 100 | | 100 | | |


Table 15. Shoulder. Percent distribution of hospital discharges by patient gender and age group, and by procedure type. 2016 and 2017

| | Total replacement (elective) | | Total repl (emerg | | Partial replacement | | Total | |
|--------------------|---------------------------------|------|----------------------|------|---------------------|------|-------|------|
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 |
| | % | % | % | % | % | % | % | % |
| Gender | | | * | | | | | |
| Male | 30.2 | 31.1 | 15.2 | 16.9 | 40.1 | 43.4 | 28.5 | 29.8 |
| Female | 69.8 | 68.9 | 84.8 | 83.1 | 59.9 | 56.6 | 71.5 | 70.2 |
| Age (male) | | | | | | | | |
| Mean age | 69.1 | 69.0 | 72.2 | 71.1 | 63.4 | 63.6 | 68.3 | 68.2 |
| Standard deviation | 9.8 | 9.7 | 9.9 | 10.2 | 12.3 | 12.5 | 10.7 | 10.6 |
| Age group | | | | | | | | |
| 0-44 | 2.0 | 2.0 | 0.8 | 1.5 | 6.9 | 6.2 | 2.9 | 2.7 |
| 45-54 | 5.8 | 6.4 | 5.1 | 5.5 | 15.5 | 14.2 | 7.7 | 7.8 |
| 55-64 | 16.6 | 17.3 | 11.5 | 16.2 | 23.9 | 29.7 | 17.5 | 19.6 |
| 65-74 | 44.2 | 43.4 | 39.1 | 33.9 | 36.8 | 31.7 | 42.1 | 39.9 |
| 75-84 | 29.9 | 29.8 | 35.6 | 37.3 | 15.1 | 15.7 | 27.5 | 27.9 |
| 85+ | 1.5 | 1.1 | 7.9 | 5.5 | 1.9 | 2.6 | 2.3 | 1.9 |
| Age (female) | | | | | | | | |
| Mean age | 72.8 | 73.0 | 75.0 | 74.7 | 70.5 | 70.1 | 73.1 | 73.1 |
| Standard deviation | 7.5 | 7.3 | 7.3 | 7.3 | 10.7 | 11.0 | 8.0 | 7.9 |
| Age group | | | | | | | | |
| 0-44 | 0.6 | 0.4 | 0.1 | 0.2 | 1.7 | 1.7 | 0.6 | 0.5 |
| 45-54 | 1.6 | 1.5 | 0.8 | 0.7 | 5.0 | 6.9 | 1.8 | 1.9 |
| 55-64 | 8.4 | 9.0 | 6.7 | 7.2 | 19.4 | 18.1 | 9.3 | 9.5 |
| 65-74 | 44.2 | 42.1 | 35.4 | 36.9 | 33.7 | 34.8 | 40.8 | 40.0 |
| 75-84 | 42.1 | 44.6 | 49.5 | 48.4 | 34.0 | 31.9 | 43.0 | 44.2 |
| 85+ | 3.0 | 2.4 | 7.7 | 6.6 | 6.2 | 6.6 | 4.5 | 3.9 |

Table 16. Shoulder. Percent distribution of hospital discharges by mode of discharge and by procedure type. 2016 and 2017

| Mode of discharge | Total replacement (elective) | | Total replacement (emergency) | | Partial replacement | | Total | |
|---|------------------------------|------|----------------------------------|------|---------------------|------|-------|------|
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 |
| | % | % | | | % | % | % | % |
| Deceased | 0.1 | 0.0 | 0.2 | 0.2 | 0.0 | 0.2 | 0.1 | 0.1 |
| Ordinary discharge | 93.5 | 92.8 | 90.6 | 90.6 | 95.2 | 94.2 | 93.2 | 92.5 |
| Discharge to a residential health care | 0.2 | 0.2 | 2.0 | 1.5 | 0.3 | 1.0 | 0.6 | 0.6 |
| Discharge to hospital at home | 0.0 | 0.0 | 0.2 | 0.2 | 0.1 | 0.0 | 0.0 | 0.0 |
| Discharge against medical advice | 0.1 | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 |
| Transfer to an acute admission unit of a different hospital | 0.4 | 0.6 | 1.2 | 1.5 | 1.2 | 1.1 | 0.7 | 0.9 |
| Transfer in the same hospital | 3.8 | 4.4 | 2.0 | 2.3 | 1.1 | 1.7 | 3.1 | 3.6 |
| Transfer to an inpatient rehabilitation hospital | 1.0 | 1.7 | 2.8 | 3.0 | 1.1 | 1.2 | 1.4 | 1.9 |
| Discharge to a nursing home | 0.8 | 0.1 | 0.6 | 0.5 | 0.8 | 0.2 | 0.8 | 0.2 |

Figure 3. Shoulder. Elective total replacement. Inter-regional mobility (attraction and escape indices) in 2016 (a) and in 2017 (b)



NOTE: For regions with low number of procedures (i.e Umbria, Molise and Basilicata) inter-regional mobility indices might be biased

Table 17. Ankle. Total replacement. Number of procedures by region of admission and procedure type. 2016-2017

| Region of admission | | Total repl | acement | |
|-----------------------|-----|------------|---------|------|
| | 20 | 16 | 20 | 17 |
| | N | % | N | % |
| Piedmont | 33 | 6.0 | 26 | 4.3 |
| Aosta Valley | 0 | 0.0 | 0 | 0.0 |
| Lombardy | 248 | 45.4 | 279 | 46.5 |
| AP Bolzano | 0 | 0.0 | 0 | 0.0 |
| AP Trento | 13 | 2.4 | 9 | 1.5 |
| Veneto | 42 | 7.7 | 53 | 8.8 |
| Friuli Venezia Giulia | 2 | 0.4 | 1 | 0.2 |
| Liguria | 1 | 0.2 | 0 | 0.0 |
| Emilia-Romagna | 131 | 24.0 | 125 | 20.8 |
| Tuscany | 7 | 1.3 | 22 | 3.7 |
| Umbria | 2 | 0.4 | 2 | 0.3 |
| Marche | 3 | 0.5 | 5 | 0.8 |
| Lazio | 30 | 5.5 | 39 | 6.5 |
| Abruzzi | 7 | 1.3 | 0 | 0.0 |
| Molise | 0 | 0.0 | 0 | 0.0 |
| Campania | 7 | 1.3 | 11 | 1.8 |
| Apulia | 5 | 0.9 | 8 | 1.3 |
| Basilicata | 1 | 0.2 | 0 | 0.0 |
| Calabria | 1 | 0.2 | 4 | 0.7 |
| Sicily | 13 | 2.4 | 16 | 2.7 |
| Sardinia | 0 | 0.0 | 0 | 0.0 |
| Italy | 546 | 100 | 600 | 100 |
| % of national volume | 100 | | 100 | |

Table 18. Ankle. Total replacement. Number of hospitals by region of admission and class of volume. 2016 and 2017

| Region of admission | Class of volume | Class of volume | | | | | | | |
|-----------------------|-----------------|-----------------|------|------|--|--|--|--|--|
| | 1 | -2 | 3- | -4 | | | | | |
| | 2016 | 2017 | 2016 | 2017 | | | | | |
| | N | N | N | N | | | | | |
| Piedmont | 5 | 7 | 0 | 1 | | | | | |
| Aosta Valley | 0 | 0 | 0 | 0 | | | | | |
| Lombardy | 12 | 16 | 3 | 1 | | | | | |
| AP Bolzano | 0 | 0 | 0 | 0 | | | | | |
| AP Trento | 1 | 1 | 0 | 0 | | | | | |
| Veneto | 10 | 5 | 1 | 1 | | | | | |
| Friuli Venezia Giulia | 2 | 1 | 0 | 0 | | | | | |
| Liguria | 1 | 0 | 0 | 0 | | | | | |
| Emilia-Romagna | 7 | 8 | 2 | 1 | | | | | |
| Tuscany | 7 | 3 | 0 | 1 | | | | | |
| Umbria | 2 | 1 | 0 | 0 | | | | | |
| Marche | 3 | 1 | 0 | 1 | | | | | |
| Lazio | 12 | 16 | 0 | 0 | | | | | |
| Abruzzi | 6 | 0 | 0 | 0 | | | | | |
| Molise | 0 | 0 | 0 | 0 | | | | | |
| Campania | 5 | 5 | 0 | 1 | | | | | |
| Apulia | 1 | 3 | 1 | 0 | | | | | |
| Basilicata | 1 | 0 | 0 | 0 | | | | | |
| Calabria | 1 | 0 | 0 | 1 | | | | | |
| Sicily | 3 | 7 | 1 | 2 | | | | | |
| Sardinia | 0 | 0 | 0 | 0 | | | | | |
| Italy | 79 | 74 | 8 | 10 | | | | | |
| % of national volume | 74.5 | 68.5 | 7.5 | 9.3 | | | | | |

| > | 4 | | Total | | | | | |
|------|------|-----|-------|-----|------|--|--|--|
| 2016 | 2017 | 20 | | 20 | 17 | | | |
| N | N | N | % | N | % | | | |
| 3 | 2 | 8 | 7.5 | 10 | 9.3 | | | |
| 0 | 0 | 0 | 0.0 | 0 | 0.0 | | | |
| 3 | 8 | 18 | 17.0 | 25 | 23.1 | | | |
| 0 | 0 | 0 | 0.0 | 0 | 0.0 | | | |
| 1 | 1 | 2 | 1.9 | 2 | 1.9 | | | |
| 4 | 5 | 15 | 14.2 | 11 | 10.2 | | | |
| 0 | 0 | 2 | 1.9 | 1 | 0.9 | | | |
| 0 | 0 | 1 | 0.9 | 0 | 0.0 | | | |
| 6 | 5 | 15 | 14.2 | 14 | 13.0 | | | |
| 0 | 1 | 7 | 6.6 | 5 | 4.6 | | | |
| 0 | 0 | 2 | 1.9 | 1 | 0.9 | | | |
| 0 | 0 | 3 | 2.8 | 2 | 1.9 | | | |
| 1 | 1 | 13 | 12.3 | 17 | 15.7 | | | |
| 0 | 0 | 6 | 5.7 | 0 | 0.0 | | | |
| 0 | 0 | 0 | 0.0 | 0 | 0.0 | | | |
| 0 | 0 | 5 | 4.7 | 6 | 5.6 | | | |
| 0 | 1 | 2 | 1.9 | 4 | 3.7 | | | |
| 0 | 0 | 1 | 0.9 | 0 | 0.0 | | | |
| 0 | 0 | 1 | 0.9 | 1 | 0.9 | | | |
| 1 | 0 | 5 | 4.7 | 9 | 8.3 | | | |
| 0 | 0 | 0 | 0.0 | 0 | 0.0 | | | |
| 19 | 24 | 106 | 100 | 108 | 100 | | | |
| 17.9 | 22.2 | 100 | | 100 | | | | |

Table 19. Ankle. Total replacement. Percent distribution of hospital discharges by patient gender and age group, and by procedure type. 2016 and 2017

| | Total replacement (elective) | | | | |
|--------------------|------------------------------|------|--|--|--|
| | 2016 | 2017 | | | |
| | % | % | | | |
| Gender | | | | | |
| Male | 57.1 | 56.6 | | | |
| Female | 42.9 | 43.4 | | | |
| Age (male) | | | | | |
| Mean age | 53.7 | 53.2 | | | |
| Standard deviation | 14.9 | 15.2 | | | |
| Age group | | | | | |
| 0-19 | 1.3 | 1.2 | | | |
| 20-39 | 15.7 | 19.5 | | | |
| 40-49 | 22.4 | 18.6 | | | |
| 50-59 | 22.8 | 21.9 | | | |
| 60-69 | 21.5 | 23.1 | | | |
| 70-79 | 14.4 | 14.8 | | | |
| 80+ | 1.9 | 0.9 | | | |
| Age (female) | | | | | |
| Mean age | 55.3 | 55.2 | | | |
| Standard deviation | 14.5 | 15.7 | | | |
| Age group | | | | | |
| 0-19 | 1.7 | 3.9 | | | |
| 20-39 | 11.1 | 10.8 | | | |
| 40-49 | 19.7 | 15.8 | | | |
| 50-59 | 27.8 | 28.6 | | | |
| 60-69 | 20.1 | 20.5 | | | |
| 70-79 | 18.4 | 17.0 | | | |
| 80+ | 1.3 | 3.5 | | | |

Table 20. Ankle. Total replacement. Percent distribution of hospital discharges by mode of discharge, by procedure type. 2016 and 2017

| Mode of discharge | Total replacen | Total replacement (elective) | |
|---|----------------|------------------------------|--|
| | 2016 | 2017 | |
| | % | % | |
| Deceased | 0.0 | 0.0 | |
| Ordinary discharge | 98.0 | 97.7 | |
| Discharge to a residential health care | 0.4 | 0.0 | |
| Discharge to hospital at home | 0.0 | 0.0 | |
| Discharge against medical advice | 0.2 | 0.0 | |
| Transfer to an acute admission unit of a different hospital | 0.0 | 0.0 | |
| Transfer in the same hospital | 1.4 | 1.5 | |
| Transfer to an inpatient rehabilitation hospital | 0.0 | 0.8 | |
| Discharge to a nursing home | 0.0 | 0.0 | |